

**PRIMARY STATEMENTS
DANGERS TO HUMANS
DOMESTIC ANIMALS**

Corrosive. May cause severe skin irritation to broken skin. Causes eye damage. Do not get on clothing. Wear goggles or face shield when handling this product. Wash after bathing vapors. Vacate poorly ventilated area. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide should not be applied to lakes, streams or ponds. Treated effluent should not be discharged into lakes, streams, ponds or public water supply. For guidance, consult the Environmental Protection Agency.

ADDITIONAL CHEMICAL HAZARDS:

IRRITATING AGENT: May irritate with direct contact. Mixing this product with acids, urine, etc. or with ammonia, acids, chemicals may release hazardous gases and mucous membranes.

Manufactured by

MARKETING ASSOC., INC.
1 St. Louis Street
Orleans, LA 70112

4394-1 EPA EST. NO.
CONTENTS

ACCEPTED
441394-1
JUL 28 1980

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the purpose of registration under

**SODIUM
HYPOCHLORITE
SOLUTION**

**"FOR SWIMMING POOL CHLORINATION
AND SANITIZING."**

**KEEP OUT OF REACH OF CHILDREN
DANGER**

FIRST AID: If on skin, wash with plenty of water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of water. Do not induce vomiting. If inhaled, get fresh air. If breathing is difficult, get medical attention. Get medical attention if affected.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

DIRECTIONS FOR SWIMMING POOL CHLORINATION

1. Test the water for pH and alkalinity. Adjust pH to 7.2-7.6 and alkalinity to 100-150 ppm. 2. Add 1 lb. of Sodium Hypochlorite Solution per 10,000 gallons of water. 3. Wait 1 hour. 4. Retest the water. 5. Repeat steps 1-4 as needed.

**DIRECTIONS FOR DISINFECTION OF POTABLE WATER
FOR HOME WELL WATER SYSTEMS**

1. Turn off the pump. 2. Add 1 lb. of Sodium Hypochlorite Solution per 10,000 gallons of water. 3. Wait 1 hour. 4. Turn on the pump. 5. Retest the water. 6. Repeat steps 1-4 as needed.

DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT

1. Wash equipment with water. 2. Add 1 lb. of Sodium Hypochlorite Solution per 10,000 gallons of water. 3. Wait 1 hour. 4. Rinse equipment with water.

NOTE:

STORAGE AND DISPOSAL:

Store in a cool, dry place. Do not mix with other chemicals. Dispose of in accordance with local regulations.