

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS &
DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contamination of feed and foodstuffs. When using this product in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation. Remove pets, birds and cover fish aquariums before spraying. Do not remain in treated area and ventilate the area after treatment is completed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and birds. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR
CHEMICAL HAZARDS**

Do not spray near or toward an open flame

DIRECTIONS FOR USE

It is a violation of Federal to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Reseal container and offer for reconditioning or disposal in approved landfill or bury in a safe place. Consult Federal, state or local disposal authorities for approved alternative procedures.

**PYRENONE[®] GENERAL PURPOSE
LIQUID INSECTICIDE**

READY TO USE.

Effective, fast acting insecticide.

For use in homes, motels, institutions, restaurants, warehouses, food processing plants, stores, office buildings.

Apply with conventional mechanical or compressed air equipment.

ACTIVE INGREDIENTS:

Pyrethrins	0.1%
*Piperonyl Butoxide, Technical	1.0 "
Petroleum Distillate	0.4 "
INERT INGREDIENTS	98.5%
	100.0%

* Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

PYRENONE Registered trademark of Fairfield American Corporation.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Harmful if swallowed. Do not spray near or toward an open flame. Avoid contamination of feed and foodstuffs

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

Manufactured By:

ARROW CHEMICAL CORPORATION

151 Horton Avenue - Lynbrook, N.Y. 11563

EPA Establishment No. 41632-NY-1 EPA Registration No. 41632 2

NET CONTENTS

GALLONS

**DIRECTIONS FOR USE
INDOOR AFFLICTION**

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. Thoroughly cleaned before using.

Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites and cadellids. Use a good quality adjusted atomizer wet spray. Spray the prey and hiding places, cracks and crevices, baseboards, and containers of stored foodstuffs. Be sure to spray chert, behind baseboards, under floorboards, and spray baseboards. Do not spray directly on foodstuffs at entry. If a face applicator is used, do not spray the face. The spray should be applied at a rate of 1/2 gallon per 1000 sq. ft.

To control carpet beetles, use a good quality adjusted atomizer wet spray. Spray the prey and hiding places, cracks and crevices, baseboards, and containers of stored foodstuffs. Be sure to spray chert, behind baseboards, under floorboards, and spray baseboards. Do not spray directly on foodstuffs at entry. If a face applicator is used, do not spray the face. The spray should be applied at a rate of 1/2 gallon per 1000 sq. ft.

To control bedbugs, use a good quality adjusted atomizer wet spray. Spray the prey and hiding places, cracks and crevices, baseboards, and containers of stored foodstuffs. Be sure to spray chert, behind baseboards, under floorboards, and spray baseboards. Do not spray directly on foodstuffs at entry. If a face applicator is used, do not spray the face. The spray should be applied at a rate of 1/2 gallon per 1000 sq. ft.

To control house flies, use a good quality adjusted atomizer wet spray. Spray the prey and hiding places, cracks and crevices, baseboards, and containers of stored foodstuffs. Be sure to spray chert, behind baseboards, under floorboards, and spray baseboards. Do not spray directly on foodstuffs at entry. If a face applicator is used, do not spray the face. The spray should be applied at a rate of 1/2 gallon per 1000 sq. ft.

To control mosquitoes, use a good quality adjusted atomizer wet spray. Spray the prey and hiding places, cracks and crevices, baseboards, and containers of stored foodstuffs. Be sure to spray chert, behind baseboards, under floorboards, and spray baseboards. Do not spray directly on foodstuffs at entry. If a face applicator is used, do not spray the face. The spray should be applied at a rate of 1/2 gallon per 1000 sq. ft.

To control ticks, use a good quality adjusted atomizer wet spray. Spray the prey and hiding places, cracks and crevices, baseboards, and containers of stored foodstuffs. Be sure to spray chert, behind baseboards, under floorboards, and spray baseboards. Do not spray directly on foodstuffs at entry. If a face applicator is used, do not spray the face. The spray should be applied at a rate of 1/2 gallon per 1000 sq. ft.

To control fleas, use a good quality adjusted atomizer wet spray. Spray the prey and hiding places, cracks and crevices, baseboards, and containers of stored foodstuffs. Be sure to spray chert, behind baseboards, under floorboards, and spray baseboards. Do not spray directly on foodstuffs at entry. If a face applicator is used, do not spray the face. The spray should be applied at a rate of 1/2 gallon per 1000 sq. ft.

To control lice, use a good quality adjusted atomizer wet spray. Spray the prey and hiding places, cracks and crevices, baseboards, and containers of stored foodstuffs. Be sure to spray chert, behind baseboards, under floorboards, and spray baseboards. Do not spray directly on foodstuffs at entry. If a face applicator is used, do not spray the face. The spray should be applied at a rate of 1/2 gallon per 1000 sq. ft.

To control earwigs, use a good quality adjusted atomizer wet spray. Spray the prey and hiding places, cracks and crevices, baseboards, and containers of stored foodstuffs. Be sure to spray chert, behind baseboards, under floorboards, and spray baseboards. Do not spray directly on foodstuffs at entry. If a face applicator is used, do not spray the face. The spray should be applied at a rate of 1/2 gallon per 1000 sq. ft.