

41632-4

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
& DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid contamination of feed and foodstuffs when using this product in food processing plants. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation. Remove pets, birds and cover fish aquariums before spraying. Do not remain in treated areas and ventilate the area after treatment is completed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR
CHEMICAL HAZARDS**

Do not spray near or toward an open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or residue that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. For containers smaller than 30 gallons triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place. For containers larger than 30 gallons, reuse container and offer for reconditioning or disposal in approved landfill or bury in a safe place. Consult Federal, state or local disposal authorities for approved alternative procedures.

**PYRENONE
FOOD PLANT SPRAY**

**A READY TO USE INSECTICIDE FOR
FOGGING AND SURFACE SPRAY**

For Use in Institutions, Hospitals, Hotels, Restaurants, Stores, Theaters, Food Processing Plants, Bakeries, Breweries, Bottling Plants, and Warehouses

Designed for Use in Edible Product Areas of Meat and Poultry Processing Plants

The insecticidal components piperonyl butoxide and pyrethrins are classified as food additives and when this product is used as directed allowable tolerance levels will not be exceeded.

ACTIVE INGREDIENTS

Pyrethrins	0.20%
*Piperonyl Butoxide, Technical	1.00%
Petroleum Distillate	98.80%
	100.00%

*Equivalent to 0.8% (butylcarbitol) (6 propylpiperonyl) ether and 0.2% related compounds.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Harmful if swallowed. Do not spray near or toward an open flame. Avoid contamination of feed and foodstuffs.

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

PYRENONE - Fairfield American Corporation Trademark
Registered

Manufactured by
ARROW CHEMICAL CORPORATION
151 Horton Avenue, Lynbrook, N.Y. 11553

EPA Registration No. 41632-4 EPA Registration No. 41632-4

NET CONTENTS GALLONS

DIRECTIONS

The use of this product in food processing plants should be controlled when the plant is not in operation. Foods should be removed or covered during treatment or food processing surface during treatment or thoroughly cleaned before using.

SPACE SPRAY: To kill flies, mosquitoes, small flies. Close doors and windows. Apply using a fogger or a fine mist. Direct portions of the fogger mist. Use 1/2 to 1 ounce per space.

To kill exposed stages of clover mites, cheese mites, saw-toothed grain beetles, weevils, granary weevils, etc. above using 2 to 3 feet of space. Keep area vacant for 2 to 3 minutes. Repeat treatment if necessary.

SURFACE SPRAY: To kill spiders, clover mites, saw-toothed grain beetles, weevils and grain mites. Direct toward insects in along baseboards, around containers of stored products where insects may hide and points of entry. Spray daily in other areas, re-spray weekly or when reinfestation occurs.

ACCEPTED
DATE 7 1960
BY [Signature]

BEST DOCUMENT AVAILABLE

41632-4

41632-4

PRECAUTIONARY STATEMENTS TO HUMANS AND DOMESTIC ANIMALS

CONTAMINATION
Avoid contamination of foodstuffs. When using this product in processing plants, food processing surfaces should be removed or thoroughly cleaned before spraying. Do not spray on or near people and animals before spraying. Do not breathe the mist.

ENVIRONMENTAL HAZARDS
Harmful to fish. Keep out of streams. Do not contaminate water. Do not spray on or near people and animals before spraying. Do not breathe the mist.

FLAMMABLE OR COMBUSTIBLE HAZARDS
Harmful toward an open flame.

USE INSTRUCTIONS
Read label to use this product in accordance with its labeling.

STORAGE AND DISPOSAL
Do not mix with water, food or feed by any means. Do not dump or empty into water. Rinse container with water. Rinse in a landfill approved for chemical waste. For containers smaller than 5 gallons, rinse (or equivalent) and recycle, reuse, or landfill or bury in a safe manner. For larger than 5 gallons, offer for recycling, reuse, or landfill or bury in a safe manner. Federal, state or local approved alternative.

PYRENONE[®] FOOD PLANT SPRAY

A READY TO USE INSECTICIDE FOR FOGGING AND SURFACE SPRAY

For Use in Institutions, Hospitals, Hotels, Restaurants, Stores, Theaters, Food Processing Plants, Bakeries, Breweries, Bottling Plants, and Warehouses.

Designed for Use in Edible Product Areas of Meat and Poultry Processing Plants.

The insecticidal components piperonyl butoxide and pyrethrins are classified as food additives and when this product is used as directed allowable tolerance levels will not be exceeded.

ACTIVE INGREDIENTS

Pyrethrins	0.20%
*Piperonyl Butoxide, Technical	1.00%
Petroleum Distillate	98.80%
	100.00%

*Equivalent to 0.8% (butylcarbitol) (S-propylpiperonyl) ether and 0.2% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

Harmful if swallowed. Do not spray near or toward an open flame. Avoid contamination of feed and foodstuffs.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PYRENONE - Fairfield American Corporation Trademark Registered.
Manufactured By
ARROW CHEMICAL CORPORATION
151 Horton Avenue Lynbrook, N.Y. 11563
EPA Registration No. 41632-4 EPA Registration No. 41632-1
NET CONTENTS **GALLONS**

DIRECTIONS FOR USE

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

SPACE SPRAY To control houseflies, fruit flies, mosquitoes, small flying moths and gnats. Close doors and windows and shut off ventilating systems. Apply using a good sprayer adjusted to deliver a fine mist. Direct spray toward upper portions of the enclosure fuming the room with mist. Use 1 to 1 ounce per 1000 cubic feet of space.

To kill exposed stages of roaches, ants, spiders, clover mites, cheese mites, confused flour beetles, saw-toothed grain beetles, red flour beetles, rice weevils, granary weevils and grain mites. Treat as above using 2 to 3 ounces per 1000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

SURFACE SPRAY To control roaches, ants, spiders, clover mites, confused flour beetles, saw-toothed grain beetles, rice weevils, granary weevils and grain mites: apply as a coarse mist directed toward insects, into cracks and crevices, along baseboards, around the base of machinery, containers of stored products and other areas where insects may hide. Spray any trails, nests and points of entry. Spray badly infested areas daily. In other areas, repeat treatment twice weekly or when reinfestation occurs.

ACCEPTED
BATCH NO. 7 1980
EPA Reg. No. 41632-4

RECIPE AVAILABLE