4/848--/ MAY 2 1 1981 Under the Federal Inserticide. Fungicide, and Rodensidde Rot. a amended, for the periods and the federal inserticide

SMITH E

ACTIVE INGREDIENT

Poly (oxyethylene (dimethyliminio) ethylene-	
(dimethyliminio) ethylene dichloride]	

This product contains 1.29 lb. of active ingredient per gallon and weighs 8.58 lb. per gallon.

KEEP OUT OF REACH OF CHIL

PRECAUTIONARY STATEMENTS ... HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Hannful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENMARONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SMITH ALGAECIDE is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbio-logical slime and other deposits.


An initial slug addition of 4.0 to 10.0 fluid ounces of SMITH ALGAECIDE per 1000 gallons of water to provide a concentration of 32 to 80 parts per million of SMITH ALGAECIDE, based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 1.0 to 10.0 fluid ounces of SMITH ALGAECIDE per 1000 gallons of water (8 to 80 parts per million of SMITH ALGAECIDE) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

STORAGE & DISPOSAL

Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Rinsate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.





and the second second