

41954-14

5/20/2003

1/4



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Antimicrobials Division 7510C
1200 Pennsylvania Avenue, N.W.
Washington, D.C. 20004

EPA Reg. Number:
41954-14

Date of Issuance:
MAY 20 2003

NOTICE OF PESTICIDE:
 x Registration
 Reregistration

Term of Issuance:

Conditional

Name of Pesticide Product:
JM Cal Hypo-TSP

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

J&M Laboratories, Inc.
658 Barret Avenue
Louisville, KY 40204

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes before your release the product for shipment:

- a. Revise the "EPA Registration Number to read, "EPA Reg. No. 41954-14".
- b. Revise the "Ingredient" Statement to read:

ACTIVE INGREDIENT
 Calcium Hypochlorite 67.70%
 INERT INGREDIENTS 32.30%
 TOTAL 100.00%

Minimum Available Chlorine 65%

Signature of Approving Official:

Robert S. Brennis, PM 32

Date:

MAY 20 2003

- c. Revise the "First Aid" Statement in accordance with PR Notice 2001-1.
- d. The "EPA Establishment Number" must be placed on the label.
- e. Under the Precautionary Statements revise "Hazardous to Human and Domestic Animals" to read: "Hazards to Human and Domestic Animals".
- f. Under the Directions for Use for "Food Egg Sanitization" revise to read: Thoroughly clean all eggs. Thoroughly mix 1 oz. of this product with 20 gallons of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130°F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable water rinse. The solution should not be reused to sanitize eggs.

3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.


A stamped copy of the label is enclosed for your records.

Handwritten signature

JM CAL HYPO-TSP

POTABLE/INDUSTRIAL WATER CHLORINATION 3" TABLETS
FOR INDUSTRIAL USE ONLY

**KILLS BACTERIA
 CONTROLS ALGAE
 DESTROYS ORGANIC CONTAMINANTS**



LABORATORIES, INC.

ACTIVE INGREDIENT: Calcium Hypochlorite..... 68%
 INERT INGREDIENTS..... 32%
 TOTAL..... 100%

MINIMUM AVAILABLE CHLORINE..... 65%

KEEP OUT OF REACH OF CHILDREN
DANGER

CONTAMINATION OR IMPROPER USE MAY CAUSE FIRE OR EXPLOSION.
ADD ONLY INTO WATER.
SEE PRECAUTIONARY STATEMENTS.

STATEMENT OF PRACTICAL TREATMENT (FIRST AID)
 IF ON SKIN: Brush off excess chemical and flush skin with water for at least 15 minutes. If irritation persists, get medical attention. IF INHALED: Remove person to fresh air. Call a physician immediately. IF IN EYES: Flush with water for at least 15 minutes. Call a physician immediately. IF SWALLOWED: Drink large quantities of water. Do not induce vomiting. Call a physician immediately.

NET WEIGHT 55 LBS (24.95 kg)

STORAGE AND DISPOSAL: Keep this product dry in its tightly closed container when not in use. Store in a cool, dry well-ventilated area. Keep away from heat or open flame. Do not reuse empty container. Place in a trash collection. Do not contaminate food or feed by storage or disposal of empty container. DISPOSAL: FOR A CONTAMINATED OR DECOMPOSING PRODUCT, DO NOT RESEAL CONTAINER IF THE PRODUCT GIVES OFF HAZARDOUS GASES, BUBBLES, OR FUMES. If possible, place in a container for hazardous waste disposal. Do not mix with any other material.

PRECAUTIONARY STATEMENTS: HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS: DANGER. Highly corrosive. Causes skin and eye damage. May be fatal if swallowed. Do not get in eyes, on skin, or on clothing. Do not handle with bare hands. Wear goggles or face shield and use rubber gloves. Only use utensils which are thoroughly clean and dry, including to nose and throat. Avoid breathing dust and fumes. Remove and wash contaminated clothing before reuse.

CHEMICAL HAZARDS: DANGER. Strong oxidizing agent. Add only to water. Contamination may start a chemical reaction. This reaction can give off heat, fire, or explosion. Do not touch. Do not get in eyes, on skin, or on clothing. Do not handle with bare hands. Wear goggles or face shield and use rubber gloves. Only use utensils which are thoroughly clean and dry, including to nose and throat. Avoid breathing dust and fumes. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, ponds, streams, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollution Discharge Elimination System (NPDES) permit and the permiting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

EMERGENCY HANDLING: In case of contamination or decomposition: 1. Do not re-seal container. 2. Place the container in an open and well-ventilated area by itself (if possible). 3. Flood with large amounts of water until the material is dissolved. 4. Place contaminated material in an approved landfill area.

SANITIZATION OF IMPOROUS FOOD CONTACT SURFACES: RINSE METHOD. - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. water to yield 200 ppm available chlorine. Immerse the surface to be sanitized in the solution for 2 minutes. Rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. The solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight. Sanitizers used in automated systems may be used for general cleaning but may not be reused for sanitizing purposes. **IMMERSION METHOD.** - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. water to yield 200 ppm available chlorine. Immerse the surface to be sanitized in the solution for 2 minutes. Rinse all surfaces thoroughly with the sanitizing solution for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight. Sanitizers used in automated systems may be used for general cleaning but may not be reused for sanitizing purposes. **FLOWPRESSURE METHOD.** - Dissassemble equipment and thoroughly clean after use. Assemble equipment in operating position prior to use. Prepare a volume of a 200 ppm available chlorine sanitizing solution equal to 110% of the volume capacity of equipment by mixing product (rate of 1 oz. product with 20 gal. water. Pump solution through system until full pressure is reached. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine. **SPRAY METHOD.** - Pre clean all surfaces after use. Use a 200 ppm available chlorine solution to control bacteria, mold or fungi and a 600 ppm solution to control bacteriophage. Prepare a 200 ppm sanitizing solution of sufficient size by thoroughly mixing product in ratio of 1 oz. product with 20 gal. water. Prepare a 600 ppm solution by thoroughly mixing product in ratio of 1 oz. product with 20 gal. water. Use spray equipment which can resist hypochlorite solutions. Always empty and rinse spray equipment with potable water after use. Thoroughly spray all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces treated with a 600 ppm solution with a 200 ppm solution.

SANITIZATION OF POREOUS FOOD CONTACT SURFACES: RINSE METHOD. - Prepare a 600 ppm solution by thoroughly mixing 3 oz. of this product with 20 gal. water. Clean surfaces in normal manner. Rinse all surfaces thoroughly with the 600 ppm solution, maintaining contact for at least 2 minutes. Prepare a 200 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. water. Rinse with the 200 ppm solution. **IMMERSION METHOD.** - Prepare a 600 ppm sanitizing solution by thoroughly mixing 3 oz. of this product with 20 gal. water. Immerse the surface to be sanitized in the solution for 2 minutes. Rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight. **IMMERSION METHOD.** - Prepare a sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. water to provide approximately 200 ppm available chlorine. Immerse the surface to be sanitized in the solution for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight. **SPRAY METHOD.** - Pre clean all surfaces after use. Prepare a 200 ppm available chlorine sanitizing solution of sufficient size by thoroughly mixing product in ratio of 1 oz. product with 20 gal. water. Use spray equipment which can resist hypochlorite solutions. Always empty and rinse spray equipment with potable water after use. Thoroughly spray all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution. Prepare a 200 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. water.

SANITIZATION OF IMPOROUS NON-FOOD CONTACT SURFACES: RINSE METHOD. - Prepare a sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 min. Do not rinse equipment with water after treatment and do not soak equipment overnight. **IMMERSION METHOD.** - Prepare a sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. water to provide approximately 200 ppm available chlorine by weight. Immerse the surface to be sanitized in the solution for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight. **SPRAY METHOD.** - Pre clean all surfaces after use. Prepare a 200 ppm available chlorine sanitizing solution of sufficient size by thoroughly mixing product in ratio of 1 oz. product with 20 gal. water. Use spray equipment which can resist hypochlorite solutions. Always empty and rinse spray equipment with potable water after use. Thoroughly spray all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution. Prepare a 200 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. water.

DISINFECTION OF NON-POROUS SURFACES: Prepare a 1000 ppm solution by thoroughly mixing 3 oz. of this product with 20 gal. of water. Clean equipment surfaces in the normal manner. Thoroughly spray all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution. Prepare a 1000 ppm sanitizing solution by thoroughly mixing 3 oz. of this product with 20 gal. water.

SANITIZATION OF POREOUS NON-FOOD CONTACT SURFACES: Prepare a 600 ppm solution by thoroughly mixing 3 oz. of this product with 20 gal. of water. Clean surfaces in normal manner. Rinse all surfaces thoroughly with the 600 ppm solution, maintaining contact for at least 2 minutes. Prepare a 200 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. water. Rinse with the 200 ppm solution.

SANITIZATION OF IMPOROUS NON-FOOD CONTACT SURFACES: Prepare a 600 ppm solution by thoroughly mixing 3 oz. of this product with 20 gal. of water. Clean surfaces in normal manner. Rinse all surfaces thoroughly with the 600 ppm solution, maintaining contact for at least 2 minutes. Prepare a 200 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 20 gal. water. Rinse with the 200 ppm solution.

SEWAGE AND WASTEWATER EFFLUENT TREATMENT: Add 1 oz. of this product to 100 gal. of wastewater. Allow to stand for 2 hours. Then add 10 gal. of water. Allow to stand for 2 hours. Then add 10 gal. of water. Allow to stand for 2 hours. Then add 10 gal. of water. Allow to stand for 2 hours.

EFFLUENT SLIME CONTROL: Apply 1 oz. of this product to 100 gal. of effluent. Allow to stand for 2 hours. Then add 10 gal. of water. Allow to stand for 2 hours. Then add 10 gal. of water. Allow to stand for 2 hours.

FILTER BEDS-SLIME CONTROL: Apply 1 oz. of this product to 100 gal. of water. Allow to stand for 2 hours. Then add 10 gal. of water. Allow to stand for 2 hours. Then add 10 gal. of water. Allow to stand for 2 hours.

DISINFECTION OF POREOUS NON-FOOD CONTACT SURFACES: Prepare a 1000 ppm solution by thoroughly mixing 3 oz. of this product with 20 gal. of water. Clean equipment surfaces in the normal manner. Thoroughly spray all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution. Prepare a 1000 ppm sanitizing solution by thoroughly mixing 3 oz. of this product with 20 gal. water.

PUBLIC WATER SYSTEMS: RESERVOIR TREATMENT: Add 1 oz. of this product to 100 gal. of water. Allow to stand for 2 hours. Then add 10 gal. of water. Allow to stand for 2 hours. Then add 10 gal. of water. Allow to stand for 2 hours.

NEW WELLS: Push the casing and seal the well. The well should stand for 24 hours.

Sanitize raw water samples obtained. Bacterial examination of the water will indicate whether further treatment is necessary. **EXISTING EQUIPMENT:** Remove equipment from service, thoroughly clean surfaces of all physical soil. Sanitize by placing 1.0 oz. of this product for each 5 cubic feet capacity (approximate) into 500-gallon available chlorine. Fill to working capacity and let stand at least 4 hours. Drain and place in service. If the residues (orange-chlorine) practical surfaces may be sprayed with solution of 1 oz. of this product for each 5 gal. water (approximately 1000 ppm available chlorine). After drying, wash with water and return to service.

COOLING TOWER/EVAPORATIVE CONDENSER WATER: **SLUG FEED METHOD:** In-line Dose: When system is noticeably fouled apply 10.000 gal. of this product per 10,000 gal. water in the system to obtain 5.10 ppm available chlorine. Repeat until control is achieved. **Sequenced Dose:** When microbial control is evident, add 2 oz. of this product per 10,000 gal. water in system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. **Bath Dosed Systems:** Must be cleaned before treatment is begun. **INTERMITTENT FEED METHOD:** In-line Dose: When system is noticeably fouled, apply 10.000 gal. of this product per 10,000 gal. water in the system to obtain 5.10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blow down. **Subsequent Dose:** When microbial control is evident, add 2 oz. of this product per 10,000 gal. water in the system to obtain 5.10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blow down. **Bath Dosed Systems:** Must be cleaned before treatment is begun. **COINTEGRATED FEED METHOD:** In-line Dose: When system is noticeably fouled, apply 10.000 gal. of this product per 10,000 gal. water in the system to obtain 5.10 ppm available chlorine. **Sequenced Dose:** When microbial control is evident, add 2 oz. of this product per 10,000 gal. water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. **Bath Dosed Systems:** Must be cleaned before treatment is begun.

LAUNDRY SANITIZERS: HOUSEHOLD LAUNDRY SANITIZERS. IN SOAKING SUDS: Thoroughly mix 1 TBSP. of this product to 10 gal. of wash water to provide 200 ppm available chlorine. Wash 5 minutes, then add soap or detergent. Immerse laundry for at least 1 1/2 minutes prior to starting the wash/rinse cycle. **IN WASHING SUDS:** Thoroughly mix 1 TBSP. of this product to 10 gal. of wash water to provide 200 ppm available chlorine. Wash 5 minutes, then add soap or detergent and start the wash/rinse cycle. **COMMERCIAL LAUNDRY SANITIZERS:** Wet fabrics or clothes should be spun dry prior to sanitization. Thoroughly mix 1 oz. of this product to 20 gallons water to obtain 200 ppm available chlorine. Promptly after turning the sanitizer, add the solution into the pre-wash prior to washing fabrics/clothes in the regular wash cycle with a good detergent. Test the level of available chlorine. If solution has been allowed to stand, Add more of this product if the available chlorine level has dropped below 200 ppm.


FARM PREMISES: Remove all animals, poultry, and feed from premises, vehicles, and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities occupied or frequented by animals or poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse in water. To disinfect, saturate all surfaces with a solution of at least 1000 ppm available chlorine. Allow 10 minutes to stand. Rinse thoroughly with water. **AGRICULTURAL USES: FOOD EGG SANITIZATION:** Thoroughly clean all eggs. Thoroughly mix 1 oz. of this product with the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable water rinse. The solution should not be reused to sanitize eggs. **FRUIT AND VEGETABLE WASHING:** Thoroughly clean all fruits and vegetables in a wash tank. Thoroughly mix 1 oz. of this product in 200 gal. of water to make a sanitizing solution of 25 ppm available chlorine. After draining the tank, submerge fruit or vegetables for 2 minutes in a second wash tank containing the recirculating sanitizing solution. Spray rinse vegetables with the sanitizing solution prior to packaging. Rinse fruit with potable water only prior to packaging.

AQUACULTURAL USES: FISH PONDS: Remove fish from ponds prior to treatment. Thoroughly mix 20 oz. of this product to 10,000 gallons of water to obtain 10 ppm available chlorine. Add more product to the water if the available chlorine level is below 1 ppm after 5 minutes. Return fish to ponds after the available chlorine level reaches zero. **FISH POND EQUIPMENT:** Thoroughly clean all equipment prior to treatment. Thoroughly mix 1 oz. of this product to 10,000 gal. of water to obtain 10 ppm available chlorine. Proceed with treatment. Thoroughly mix 1 oz. of this product to 10,000 gal. of water to obtain 10 ppm available chlorine. **COOLING TOWER/EVAPORATIVE CONDENSER WATER:** Thoroughly mix 1 oz. of this product to 10,000 gal. of water to obtain 10 ppm available chlorine. **SEQUENCED DOSE:** When microbial control is evident, add 2 oz. of this product to 10,000 gal. of water in the system to obtain 5.10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blow down. **Subsequent Dose:** When microbial control is evident, add 2 oz. of this product to 10,000 gal. of water in the system to obtain 5.10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blow down. **Bath Dosed Systems:** Must be cleaned before treatment is begun. **COINTEGRATED FEED METHOD:** In-line Dose: When system is noticeably fouled, apply 10.000 gal. of this product per 10,000 gal. water in the system to obtain 5.10 ppm available chlorine. **Sequenced Dose:** When microbial control is evident, add 2 oz. of this product per 10,000 gal. water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. **Bath Dosed Systems:** Must be cleaned before treatment is begun.

AGRICULTURAL USES: FOOD EGG SANITIZATION: Thoroughly clean all eggs. Thoroughly mix 1 oz. of this product with the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable water rinse. The solution should not be reused to sanitize eggs. **FRUIT AND VEGETABLE WASHING:** Thoroughly clean all fruits and vegetables in a wash tank. Thoroughly mix 1 oz. of this product in 200 gal. of water to make a sanitizing solution of 25 ppm available chlorine. After draining the tank, submerge fruit or vegetables for 2 minutes in a second wash tank containing the recirculating sanitizing solution. Spray rinse vegetables with the sanitizing solution prior to packaging. Rinse fruit with potable water only prior to packaging.

AQUACULTURAL USES: FISH PONDS: Remove fish from ponds prior to treatment. Thoroughly mix 20 oz. of this product to 10,000 gallons of water to obtain 10 ppm available chlorine. Add more product to the water if the available chlorine level is below 1 ppm after 5 minutes. Return fish to ponds after the available chlorine level reaches zero. **FISH POND EQUIPMENT:** Thoroughly clean all equipment prior to treatment. Thoroughly mix 1 oz. of this product to 10,000 gal. of water to obtain 10 ppm available chlorine. **COOLING TOWER/EVAPORATIVE CONDENSER WATER:** Thoroughly mix 1 oz. of this product to 10,000 gal. of water to obtain 10 ppm available chlorine. **SEQUENCED DOSE:** When microbial control is evident, add 2 oz. of this product to 10,000 gal. of water in the system to obtain 5.10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blow down. **Subsequent Dose:** When microbial control is evident, add 2 oz. of this product to 10,000 gal. of water in the system to obtain 5.10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blow down. **Bath Dosed Systems:** Must be cleaned before treatment is begun. **COINTEGRATED FEED METHOD:** In-line Dose: When system is noticeably fouled, apply 10.000 gal. of this product per 10,000 gal. water in the system to obtain 5.10 ppm available chlorine. **Sequenced Dose:** When microbial control is evident, add 2 oz. of this product per 10,000 gal. water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. **Bath Dosed Systems:** Must be cleaned before treatment is begun.

FOOD PROCESSING PLANTS: POULTRY DRINKING WATER: Spray or flush with a solution containing 1 oz. of this product for every gallon of water. Treat poultry drinking water to a dosage of 1 to 5 ppm available chlorine by adding 1 to 5 oz. of this product per 1000 gallons of water. **FISH KILLING:** Cryocera and degassed fish removed from fishing vessel are placed in a wash tank of sea water or fresh water which has been treated with enough product to produce a chlorine residual of 25 ppm as determined by a test kit. Remove fish from treated water 24 to 48 hours before filleting. After scaling the fish are again washed in a 25 ppm solution and are ready for filleting.



MANUFACTURED FOR:
J&M LABORATORIES, INC.
 658 BARRET AVENUE
 LOUISVILLE, KY 40204
 (502) 585-3176
 (800) 473-6173


EPA REG. NO. 425-00000
 EPA EST. NO. 00000000



WARNING
AVISO



Children can fall into bucket and drown. Keep children away from bucket with even a small amount of water. Los niños pueden caerse a esta cubo y ahogarse. No deja que los niños se acerquen a este cubo aunque sólo contenga una pequeña cantidad de agua.



ACCEPTED with COMMENTS
 EPA Letter Dated: MAY 20 2003
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 41954-14

4/4