

DIRECTIONS FOR USE

MUSHROOMS

To control sciard fly larvae in mushroom growing facilities. The Insect Growth Regulator DIMILIN in the mushroom growing media will prevent the development of the larval stages of the sciard fly. This effectively stops reproduction in the growing medium and prevents damage to the mushroom.

Because of its unique type of activity, do not expect immediate reductions in adult fly populations. DIMILIN does not directly affect adults, but kills the larva in the growing medium.


COMPOST TREATMENT: Apply 2.4 to 4 lbs. DIMILIN 25W per 1000 sq. feet to the compost at spawning time by thorough incorporation such as with a spawning machine. This is equivalent to 30-50 ppm active assuming a compost wet weight of 40 lbs. per cu. ft.

CASING TREATMENT: Apply 13.5 oz of DIMILIN 25W per 1000 sq. feet at the time of casing by thorough incorporation into the casing or in sufficient water to obtain a thorough drench. This is equivalent to a rate of 30 ppm active assuming a casing weight of 6700 pounds per 1000 sq. ft.

CHRYSANTHEMUMS

For control of beet armyworm, apply ½ to 1 lb. of DIMILIN 25W per acre in a dilute spray not to exceed 200 gallons of water per acre. Begin applications when larvae appear and repeat at weekly intervals as required. DIMILIN interferes with the molting process of insects and is, therefore, slow in its action. The insect dies during molting following contact, therefore, full effect will not be seen for 3 to 5 days following application.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



Mr. Nathaniel Digges
J&B Industries
211 J&B Drive
San Benito, TX 78586

Dear Mr. Digges:

Subject: Fog-20
EPA Registration No. 42050-4
Your Letter Dated May 21, 1987

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,




Phil Hutton
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

57607::Schroeder:C.Disk;KENCO:10/25/88:rw:vo:rw

CONCURRENCES

SYMBOL							
SURNAME							
DATE							



FOG-20

INSECTICIDE

EPA Reg. No. 42650-4
E-A Est. No. 42650-TX-1

An Effective Semi-Concentrated Pyrethroid Insecticide.

For Application in Specialized Mechanical, Thermal, or Compressed Air Foaming Equipment.

For Use in Bakeries, Food Plants, Restaurants, Hotels, Motels, Hospitals, Schools, Office Buildings, Food Processing or Handling Plants and Warehouses.

ACTIVE INGREDIENTS:

Pyrethrins.....	0.3%
*Eperonyl Butoxide, Technical.....	2.4%
retroleum Distillate.....	97.3%
	100%

*Equivalent to 1,027 (hexylphenyl) (4-propylphenyl) ether and 0.46% related compounds.

FOG-20 - Registered trademark of Fairfield American Corp.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
See side panel for additional precautions

ACCEPTED
NET CONTENTS
Batch No. 42050-4
DEC 27 1988

NET CONTENTS: 1.05 L (35.3 FL OZ) for the pesticide
registered under EPA Reg. No.

Manufactured by: **42050-4**
J & B INDUSTRIES
2100 J&B DRIVE • SAN BENITO, TEXAS 78586
PHONE: (512) 399-4003

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

This ready-to-use concentrate may be dispersed through a good mechanical, thermal or compressed air foaming or sprayer equipment. If used in edible product areas of Federally Inspected meat, poultry, shell egg grading and egg product establishments, it must be dispersed as an aerosol spray (no droplets or particles in diameter and R.S. less than 33 microns).

SEMI-TREATMENTS: Close doors and windows and shut off ventilating systems. Apply as a fog or fine mist, distributing the spray throughout the area. Use at a rate of 1 ounce per 100 cubic feet of space. Do not spray behind and under obstructions and onto crawling insects whenever possible. Keep area closed for 10 to 15 minutes. Vacate treated area and ventilate before re-occupancy. Repeat treatment as necessary. For wasps and hornets, spray late in the evening when insects are at rest.

The above described application will kill all possible exposed stages of house flies, fruit flies, mosquitoes, silverfish, small flying insects, gnats, crickets, cockroaches, lightbulb gnats, yellow jackets, silver chafers, stoneflies, crane flies, beetles, and those common household pests: granary weevils, rice weevils, confused flour beetles, saw toed grain beetles, spider beetles, cigarette beetles, longhorn beetles, Argentine grain moths, 15 liter room floor beetles, black head beetles, tobacco moth, meal worms, and grain weevils.

SPRAY APPLICATION: For general pest control of crawling insects, lightly spray as a coarse mist directly and close to, into cracks, crevices, under furniture, around containers of food products and other areas where the insects may be. Repeat as necessary.

SPRAY APPLICATION: Spread 1/2 oz. (14.3 ml) of this material per 1,000 cubic feet of space by using a thermal or compressed air foaming equipment. Apply as a space spray with a mechanical or thermal generator. Apply only when flying insects are present. During periods of heavy infestation several applications may be necessary. Do not make more than 10 applications.

STORAGE AND

Do not store in areas where children may have access.

RECOMMENDATIONS: Do not use in areas where children may have access. Do not use in areas where children may have access.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose of or by other approved method.

PRECAUTIONS HARMFUL TO BEES

Caution: Harmful if swallowed. Avoid breathing mist. Avoid contact.

The use of this product in establishments should be restricted to periods when the product should be removed. All food should be covered during application. If necessary, use a detergent solution of water to remove all. Do not apply directly to food. Wash hands before spraying.

PRECAUTIONS

If swallowed, contact a physician. Do not use in areas where children may have access. Do not use in areas where children may have access.

If inhaled, remove to fresh air. Apply as directed.

If on skin, wash with soap and water. Do not use in areas where children may have access.

PRECAUTIONS

This product is toxic. Do not use in areas where children may have access. Do not use in areas where children may have access.

Do not use in areas where children may have access. Do not use in areas where children may have access.