

middle panel 10F5
110.21
42056-8

T C I

Trace Chemicals, Inc.

CAPTAN-METHOXYCHLOR 75-3

Seed Protectant
(Insecticide-Fungicide)

ACCEPTED
MAR 3 1986
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 42056-8

Active Ingredients	By wt.
*Captan	70.9%
Related Derivatives.	4.1%
**Methoxychlor, Technical.	3.0%
Inert Ingredients.	22.0%
*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide. **Equivalent to 2.64% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and .36% of other isomers and related compounds.	

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

If swallowed, give a large amount of water to drink, make person vomit, and get medical attention. In case of eye contact, immediately flush eyes with plenty of clear water and get medical attention.


SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT: 50 Pounds

EPA Reg No. 42056-

EPA Est. No. 42056-IL-1

Manufactured by
Trace Chemicals, Inc.
Pekin, Illinois 61555-0518



PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Avoid inhalation of dust or spray mist. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administrator (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part II. Wash hands after use.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Open dumping is prohibited.

Do not reuse empty container.

STORAGE

Keep pesticide in original container.

Do not put concentrate or dilute into food or drink containers.

Store in a cool, dry place.

Protect from excessive heat.

PESTICIDE DISPOSAL

This product is acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinse is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

left panel, cont.


Captan-Methoxychlor 75-3 Seed Protectant is a special formulation containing captan and methoxychlor for seed treatment application. Captan-Methoxychlor 75-3 Seed Protectant has no adverse effect on seed germination or planting rate. It is used in preventing seed rot and damp-off diseases on germinating seeds and seedlings and smut on onion and hulled sorghum seeds. It also reduces losses from Cadelle, Rice Weevil, Confused Flour Beetle, Angoumois Grain Moth and Indian Meal Moth during preplanting storage of clean seed.

Do not feed treated seed to animals. Do not use treated seed for human consumption. Seeds treated with Captan-Methoxychlor 75-3 Seed Protectant, sold or shipped as feed or foodstuffs may be considered adulterated under State and Federal Laws. Treated seed must be labeled as follows: "TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL".

DIRECTIONS: Apply as a slurry treatment. The recommended amount of product should be mixed with the required amount of water for the slurry treater machine to be used. Do not allow slurry mixtures to stand overnight in the machine. Clean out equipment thoroughly between treating operations.

GLADIOLUS CORNS: For prevention of Bulb Rot, use 1-1/2 oz. in 1 gal. of water(1 lb. to 10 gals.). Soak bulbs in mixture 20 to 30 minutes.

ONION SEED PELLETING: Use 1 lb. Captan-Methoxychlor 75-3 Seed Protectant per 1 lb. of seed.



right panel, cont.

SEED	Dosage Rate	Lbs. Seed Treated/Lb. of Product
COTTON - Machine Delinted Fuzzy(not delinted)	3 oz/100 lbs.	530
BLUE GRASS	5-1/2 oz/100 lbs.	290
CABBAGE CAULIFLOWER BROCCOLI BRUSSELS SPROUTS	3-1/2 oz/100 lbs.	450
SPINACH	4 oz/100 lbs.	400
SORGHUM hulled(Kernel Saut)	3 oz/100 lbs.	530
PEPPERS	2 oz/100 lbs.	800
COWPEAS	2 oz/100 lbs. 1-1/4 oz/bu.	800
RICE	2-1/4 oz/100 lbs.	710

Conditions of Sale

TRACE CHEMICALS, INC. warrants that this material conforms to its label description and is reasonably fit for use if used strictly in accordance with label directions. Neither Trace nor Seller makes any other warranty, expressed or implied, as to MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Since storage, weather, soil or plant conditions, farming practices and other factors beyond Trace's control may adversely effect crop yield, even when this material is used in accordance with label directions, Buyer assumes all risks incident thereto. Trace's and Seller's liability shall be limited to return of the purchase price.

Trace Chemicals, Inc.
Pekin, Illinois 61555-0518