

PM 04

42057-39

12/8/99

ACCEPTED

Best Available Copy

DEC 8 1999



Under Federal Insecticide, Fungicide, and Rodenticide Act as amended, this product is a restricted use pesticide. EPA Reg. No. 42057-39

PEST MEAL



Manufactured by: **MORGRO, INC.** / Salt Lake City, UT

2 9 2

MORGRO PEST MEAL

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Scatter PEST MEAL around annual flowers, vegetables, and perennial shrubs bothered by slugs, snails, earwigs, ants, grasshoppers, cutworms, millipedes or sowbugs. Apply at rates specified in the chart below.

CROP	RATE	RETREATMENT INTERVAL	PRE-HARVEST INSTRUCTIONS
Asparagus	16 oz/850 sq ft	7 Days/Not to exceed 3 applications	1 Day
Fresh Beans, Fresh Peas	12 oz/850 sq ft	7 Days/Not to exceed 3 applications	3 Days
Pepper, Tomato, Cabbage	16 oz/850 sq ft	7 Days/Not to exceed 4 applications	3 Days
Potatoes, Radishes, Carrots, Turnips	16 oz/850 sq ft	7 Days/Not to exceed 3 applications	7 Days
Sweet Corn	16 oz/850 sq ft	3 Days/Not to exceed 8 applications	14 Days
Cucumber, Squash, Melons	8 oz/850 sq ft	7 Days/Not to exceed 6 applications	3 Days
Lettuce, Spinach	16 oz/850 sq ft	7 Days/Not to exceed 3 applications	14 Days

FOR ADULT STRAWBERRY ROOT WEEVIL:

Apply 1 teaspoon of PEST MEAL around the base of Strawberry plants, Rhododendrons, Camellias, Primroses, and Azaleas. Broadcast lightly on soil surface using 1 lb. per 850 square feet. (DO NOT APPLY DIRECTLY TO PLANTS.) Apply in April or May only, or when weevil is in pupae stage. Repeat treatment in 2 to 3 weeks if hatching weevils continue to appear.

NON-WARRANTY: Manufacturer or Seller makes no warranty, expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described and recommended on the label.

This package is sold by weight, not volume. Some settling of contents may have occurred during shipment or handling.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: (READ BEFORE USING)

May be fatal if swallowed or inhaled. DO NOT BREATHE DUST OR VAPORS. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. KEEP OUT OF THE REACH OF CHILDREN AND PETS. This pesticide may be fatal to children and dogs or other pets if eaten. Keep children and pets out of treated area. DO NOT USE INSIDE THE HOUSE. This product may be attractive to dogs. Keep dogs away from treated areas. Avoid contamination of feed and foodstuffs. DO NOT APPLY DIRECTLY TO EDIBLE CROPS. Use and store under conditions where there will not be a hazard.

STATEMENT OF PRACTICAL TREATMENT

If in eyes or on skin: Flush eyes thoroughly with water; wash skin thoroughly with soap and water.

If swallowed: Induce vomiting by administration of syrup of ipecac or under medical supervision, call a physician.

If inhaled: Remove from exposure and seek medical attention

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use (PAM), opiates or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS

This product is toxic to birds and wildlife. Birds feeding in treated areas may be killed. Do not apply to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.