

193



Manufactured by MORGRO CHEMICAL COMPANY
P.O. Box 151048 - Salt Lake City, Utah 84115

ACCEPTED
MAY 31 1988

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for this pesticide registered under EPA Reg. No. 42057-92

BEST AVAILABLE

293

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May irritate eyes, nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. In case of contact, flush skin or eyes with plenty of water. For eyes, get medical attention.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish in adjacent areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on label.

Chemigation: Do not apply this product through any type of irrigation system.

Physical or Chemical Hazards: Keep Away From Fire Or Sparks.

Safety Rules: It is recommended to wear a long sleeve shirt, pants, and gloves during application. If clothing becomes contaminated, remove and clean after use.

STORAGE AND DISPOSAL

STORAGE: Keep product in original container. Store in a cool, dry place. Keep container closed when not in use. Store container in a locked storage area.

DISPOSAL: Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Morgro BENOMYL SYSTEMIC FUNGICIDE is highly effective against fungus diseases on roses, flowers, certain fruits and vegetables, ornamental plants and turf, providing both curative (eradicant) action and protective action. The addition of a surfactant to the spray mixture increases surface acceptance of spray on hard-to-wet plants such as roses.

FOR ROSES, FLOWERS & ORNAMENTALS: Controls Powdery Mildew, Black Spot of Rose, (Botrytis) Gray Mold. Use 1 level tablespoon in 2 gallons water. For control of Black Spot of Rose, use one level tablespoon in 1 gallon of water. Mix thoroughly. Apply when disease first appears and repeat at 10-14 day intervals during growing season. Shorten interval during humid or rainy weather.

FOR TURF USE: Spray on lawn areas as directed below. Apply when disease first appears and repeat at 10-14 day intervals as needed.

DOLLAR SPOT-Mix 1 oz. (6 level tablespoons) Benomyl in 2-5 gallons water and apply evenly over 1,000 sq.ft. LARGE BROWN PATCH, FUSARIUM PATCH-Mix 1 oz. in 1½-2 gallons water and apply evenly to 500 sq.ft. Apply when disease appears and repeat at 10-14 day intervals. If conditions favor disease development, reduce interval 5-7 days.

FOR TOMATOES: Controls (Botrytis) Gray Mold, Leaf Mold, White Mold, (Cercospora) Leaf Spot, Phoma Leaf Spot-Mix 1 to 2 level tablespoons Benomyl in 2 gallons water. Apply when disease first appears and repeat at 7-14 day intervals as needed.

FOR STRAWBERRIES: Controls (Botrytis) Gray Mold, Powdery Mildew, Leaf Scorch, Leaf Blight, Leaf Spot-Mix 1 tablespoon Benomyl in 2 gallons water. Apply at early bloom and full bloom and repeat at 10-14 day intervals as needed.

FOR CANEBERRIES: (Raspberries, Blackberries, Boysenberries, Loganberries, Dewberries): Controls (Botrytis) Gray Mold, Powdery Mildew, Penicillium Rot-Mix 1½ tablespoons Benomyl in 2 gallons water. Apply at early bloom and at full bloom; make up to 3 additional applications at 14-day intervals as needed. Do not apply within 3 days of harvest.

FOR CUCUMBERS, MELONS, PUMPKINS, SUMMER & WINTER SQUASH: Controls Target Spot (Cucumbers), Gummy Stem Blight, Powdery Mildew, Anthracnose-Mix 1 level tablespoon with 2 gallons water and apply when stems begin to run or when disease first appears. Repeat at 7-14 day intervals as needed.

FOR BEANS: Controls White Rot, (Botrytis) Gray Mold-Mix 1 level tablespoon Benomyl with 2 gallons water. Apply at early (25-50%) bloom and repeat at full bloom. Do not apply within 14 days of harvest (28 days for lima beans); do not use where crop is grown only for forage purposes.

NON-WARRANTY: Manufacturer or Seller makes no warranty expressed or implied concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

EPA Est. No. 42057-UT-1
Lot No.

EPA Reg. No. 42057-92

(UPC Code)

