

ACCEPTED
 MAR 22 1982
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 42057-96

BEST AVAILABLE COPY

COMPARA
 PMS III
 PREMIUM WEED & FEED

COMPARATM PMS III

Premium Weed & Feed

Contains proven TRIMEC[®] Herbicide for effective control
 of most broadleaf weeds. 22-3-3 plus Iron.

ACTIVE INGREDIENTS:	*2,4-dichlorophenoxyacetic acid	0.664%
	**2-(2-methyl-4-chlorophenoxy) propionic acid	0.309%
	***samba (3,6-dichloro-o-anisic acid)	0.065%
INERT INGREDIENTS:	Relative to herbicidal content	99.945%
	TOTAL	100.000%

Equivalent to: *1.191 lbs. per acre, **0.533 lbs. per acre and ***0.112 lbs. per acre.

TRIMEC[®] is a registered trademark
 of PBI/Gordon Corporation,
 U.S. Patent No. 3,264,188.
 EPA Reg. No. 42057-98
 EPA Est. No. 42057-UT-1

CAUTION
 KEEP OUT OF REACH OF CHILDREN
 SEE OTHER CAUTIONS ON BACK PANEL

NET WEIGHT: 21 lbs. (9.52 Kg.) FEEDS 5,250 SQ. FT. (487.7 sq. meters)

Manufactured by
 MORGRO CHEMICAL COMPANY


P.O. Box 151048 • 145 West Central Avenue • Salt Lake City, Utah 84115

P41 23
 P9 184
 42057-96

022  TG


FREIGHT SHIPPING BAG
Meeting requirements of
APPLICABLE FREIGHT CLASSIFICATION
Developed by
HEMCO COMPANY, INC.
VANCOUVER, WASH.

MORGRO



TM


MORGRO



BEST AVAILABLE COPY

294

COMPARE™ PLUS III



MORGRO

COMPARE™ PLUS III

Premium Weed & Feed

22-3-3 plus Iron

MORGRO COMPARE™ PLUS III IS A SPECIAL FORMULATION OF PREMIUM QUALITY LAWN FERTILIZERS BLENDED WITH PROVEN TRIMEC® HERBICIDE FOR EFFECTIVE CONTROL OF MOST BROADLEAF WEEDS.

- Contains two types of Nitrogen fertilizers—one to produce rapid greening and another, slow-release Nitrogen source to provide a steady, controlled feeding for up to 8 weeks.
- Safe to use on all types of turf grasses. Will not burn when used as directed.
- Contains 2% Iron to prevent and correct yellowing due to iron deficiency, plus 6% Sulfur to reduce alkali and correct pH balance in western soils.
- COMPARE™ Plus III effectively controls the following weeds when used as directed:

Bedstraw	Dandelion	Lespedeza	Poison Ivy	Spurge
Black Medic	Dock	Mallow	Poison Oak	Thistle
Buckhorn	Ground Ivy	Morning Glory	Purslane	Wild Carrot
Burdock	Heart-All	Oxalis	Ragweed	Wild Garlic
Chicory	Henbit	Peppergress	Sheep Sorrel	Wild Lettuce
Chickweed	Knockweed	Pigweed	Shepherd's Purse	Wild Onion
Clover	Lambquarter	Plantain	Speedwell	Yarrow

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	22%
15% Water Soluble Nitrogen	
7% Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	3%
Sulfur (S)	6%

BEST AVAILABLE COPY

374

MORGRO

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Morgro COMPARE™ Plus III can be used on Blue Grass, Fescue, Bermuda, and Zoysia lawns. Apply Morgro COMPARE™ Plus III with a broadcast or cyclone spreader set to deliver 4 pounds per 1,000 sq. ft. WATER LAWN THOROUGHLY BEFORE APPLICATION. After application, sweep particles off walks and driveways to avoid staining. Do not water, mow or rake lawn for 48 hours after application. For best results apply Morgro COMPARE™ Plus III in spring or early fall when weeds are actively growing. COMPARE™ Plus III will not burn grasses when used as directed. Re-apply Morgro COMPARE™ Plus III or another Morgro lawn fertilizer 6 to 8 weeks after application of this product.

SPREADER SETTINGS

Apply according to spreader manufacturer's recommended settings. When using a wheeled, drop-type spreader, overlap wheel tracks to assure even coverage.

SPREADER TYPE

Scotts Standard, Automatic or
Precision Flow Control Models
Cyclone Spreader

SETTING

8
3

SUGGESTED FERTILIZER APPLICATION PROGRAM

Lawn research demonstrates that turf grasses need adequate quantities of plant food to maintain good health and color. Morgro recommends the following application programs:

GOOD - Fertilize twice: early spring and in early fall.

BETTER - Fertilize three times: early spring, late spring or early summer, and early fall.

BEST - Fertilize four times: early spring, late spring or early summer, late summer or early fall, and late fall.

If lawn becomes infested with Crabgrass or Spurge, apply Morgro CRABGRASS AND SPURGE KILLER (includes fertilizer) in early spring or late fall. If dandelions or other perennial broadleaf weeds appear, apply Morgro COMPARE™ Plus III Weed & Feed or Morgro WEEDIT II SPRAY in early spring and as needed for continued control.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals: CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from area. Do not contaminate water used for irrigation or domestic purposes.

Application Precautions: Do not use where desirable clovers are present or allow product to wind drift on flowers, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. Do not apply in areas underlain by roots of desirable trees and shrubs.

STORAGE & DISPOSAL

Do not store near pesticides or seeds. Do not reuse empty container. Wrap and put in trash collection.

NON-WARRANTY: Manufacturer or Seller makes no warranty, expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described and recommended on the label.

Manufactured by **MORGRO CHEMICAL COMPANY**
P.O. Box 151048 • 145 West Central Avenue • Salt Lake City, Utah 84115

Product of USA

© 1981 Morgro Chemical Company

All rights reserved

Lot No.

BEST AVAILABLE COPY

7/12/81

ACCEPTED

MAR 22 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 42057-96

BEST AVAILABLE COPY

COMPARA PBI

COMPARETM PBI

Premium Weed & Feed

Contains proven TRIMEC[®] Herbicide for effective control of most broadleaf weeds. 22-3-3 plus Iron.

ACTIVE INGREDIENTS	**2,4-dichlorophenoxyacetic acid	0.684%	TRIMEC [®] is a registered trademark of PBI/Gordon Corporation, U.S. Patent No. 3,284,186.
	**2-(2-methyl-4-chlorophenoxy) propionic acid	0.306%	
	***D-camba (3,6-dichloro-o-anisic acid)	0.056%	
INERT INGREDIENTS	Relative to herbicidal content	98.945%	EPA Reg. No. 42057-96 EPA Est. No. 42057-UT-1
	TOTAL	100.000%	

Equivalent to **1191 lbs. per acre, ***0.533 lbs. per acre, and ****0.112 lbs. per acre.

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON BACK PANEL


NET WEIGHT: 21 lbs. (9.52 Kg.) FEEDS 5,250 SQ. FT. (487.7 sq. meters)

Manufactured by
MORGRO CHEMICAL COMPANY
P.O. Box 151048 - 145 West Central Avenue - Salt Lake City, Utah 84115

022  TG


FREIGHT SHIPPING BAG
Meeting requirements of
APPLICABLE FREIGHT CLASSIFICATION
Manufactured by
BERRY COMPANY, INC.
VANCOUVER, WASH.


MORGRO



TM

MORGRO





MORGRO

COMPARE™ PLUS III

Premium Weed & Feed

22-3-3 plus Iron

MORGRO COMPARE™ PLUS III IS A SPECIAL FORMULATION OF PREMIUM QUALITY LAWN FERTILIZERS BLENDED WITH PROVEN TRIMEC® HERBICIDE FOR EFFECTIVE CONTROL OF MOST BROADLEAF WEEDS.


- Contains two types of Nitrogen fertilizers—one to produce rapid greening and another, slow-release nitrogen source to provide a steady, controlled feeding for up to 8 weeks.
- Safe to use on all types of turf grasses. Will not burn when used as directed.
- Contains 2% Iron to prevent and correct yellowing due to iron deficiency, plus 6% Sulfur to reduce alkali and correct pH balance in western soils
- COMPARE™ Plus III effectively controls the following weeds when used as directed:

Bedstraw	Dandelion	Lespedeza	Poison Ivy	Spurge
Black Medic	Dock	Mallow	Poison Oak	Thistle
Buckhorn	Ground Ivy	Morning Glory	Purslane	Wild Carrot
Burdock	Heal All	Oxalis	Ragweed	Wild Garlic
Chicory	Henbit	Peppergress	Sheep Sorrel	Wild Lettuce
Chickweed	Knockweed	Pigweed	Shepherd's Purse	Wild Onion
Clover	Lambquarter	Plantain	Speedwell	Yarrow

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	22%
15% Water soluble Nitrogen	
7% Water insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3%
Sulfur Polash (K ₂ O)	3%
Sulfur (S)	6%
Iron (Fe)	2%

COMPARE™
LAWN



BEST AVAILABLE COPY

MORGO

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Morgro COMPARE™ Plus III can be used on Blue Grass, Fescue, Bermuda, and Zoysia lawns.

Apply Morgro COMPARE™ Plus III with a broadcast or cyclone type spreader set to deliver 4 pounds per 1,000 sq. ft. WATER LAWN THOROUGHLY BEFORE APPLICATION.

After application, sweep particles off walks and driveways to avoid staining. Do not water, mow or rake lawn for 48 hours after application.

For best results apply Morgro COMPARE™ Plus III in spring or early fall when weeds are actively growing. COMPARE™ Plus III will not burn grasses when used as directed.

Re-apply Morgro COMPARE™ Plus III or another Morgro lawn fertilizer 6 to 8 weeks after application of this product.

SPREADER SETTINGS

Apply according to spreader manufacturer's recommended settings. When using a wheeled, drop-type spreader, overlap wheel tracks to assure even coverage.

SPREADER TYPE

Scotts Standard, Automatic or
Precision, Flow Control Models
Cyclone Spreader

SETTING

8
3

SUGGESTED FERTILIZER APPLICATION PROGRAM

Lawn research demonstrates that turf grasses need adequate quantities of plant food to maintain good health and color. Morgro recommends the following application programs.

GOOD - Fertilize twice, early spring and in early fall.

BETTER - Fertilize three times, early spring, late spring or early summer, and early fall.

BEST - Fertilize four times, early spring, late spring or early summer, late summer or early fall, and late fall.

If lawn becomes infested with Crabgrass or Spurge, apply Morgro CRABGRASS AND SPURGE KILLER (includes fertilizer) in early spring or late fall. If dandelions or other perennial broadleaf weeds appear, apply Morgro COMPARE™ Plus III Weed & Feed or Morgro WEEDIT II SPRAY in early spring and as needed for continued control.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals. CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from area. Do not contaminate water used for irrigation or domestic purposes.

Application Precautions: Do not use where desirable clovers are present or allow product to wind drift on flowers, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. Do not apply in areas unshaded by roots of desirable trees and shrubs.

STORAGE & DISPOSAL

Do not store near pesticides or seeds. Do not reuse empty container. Wrap and put in trash collection.

NO WARRANTY: Manufacturer or Seller makes no warranty, expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described and recommended on the label.

Manufactured by **MORGO CHEMICAL COMPANY**
P.O. Box 151048 • 145 West Central Avenue • Salt Lake City, Utah 84115

Product of USA

© 1981 Morgro Chemical Company

All rights reserved.

Lot No.