

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**DANGER:** Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

**PHYSICAL AND CHEMICAL HAZARDS:**  
**STRONG OXIDIZING AGENT:** Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc., or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

manufactured by

**SURE CHEMICAL CORPORATION  
P.O. BOX 395  
Route 140  
Upton, MA 01568**

**EPA REG. NO. 42263-3 EPA EST. NO.**

**NET CONTENTS:**

**SCC-500 CHLOR  
SANITIZER**

**"FOR SWIMMING POOL CHLORINATION  
AND SANITIZING."**

ACTIVE INGREDIENT  
SODIUM HYPOCHLORITE 12.5%  
INERT INGREDIENTS 87.5%

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**FIRST AID:** If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel.

**DIRECTIONS FOR USE  
GENERAL CLASSIFICATION**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT  
IN A MANNER INCONSISTENT WITH ITS LABELING**

**DIRECTIONS FOR SWIMMING POOL CHLORINATION**

For a new pool or for spring start-up, superchlorinate with one (1) pint (16 flu. ozs.) of sodium hypochlorite solution for each 3000 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.

For Pool Maintenance

- A) Adjust pool water pH to 7.6 - 7.2 range and maintain.
- B) Add manually or by a feeder device this sodium hypochlorite solution at a rate to maintain an available chlorine residual of 0.6 to 1.0 ppm. One-fourth pint (4 flu. ozs.) of this sodium hypochlorite solution for each 4000 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.6 to 1.0 ppm available chlorine will depend on temperature and number of swimmers. Use test kit to make certain the pH and chlorine residual are in the proper range.

**DIRECTIONS FOR DISINFECTION OF POTABLE WATER  
FOR HOME WELL WATER SYSTEMS**

Dilute this sodium hypochlorite solution in the ratio of one part sodium hypochlorite solution to 11 parts softened water. Mix sodium hypochlorite solution and water thoroughly and begin feeding of solution with a hypochlorinator (metering pump). Maintain a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm throughout the distribution system, as determined by a DPD chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations. Check water frequently with a DPD chlorine test kit.

\*Contact your local health department for further details.

**DIRECTIONS FOR SANITIZING FOOD PROCESSING  
OR DAIRY EQUIPMENT**

Clean equipment in the normal manner. Just before using, rinse all surfaces thoroughly with this sodium hypochlorite solution containing 200 ppm available chlorine. Maintain contact with disinfectant for a minimum of two minutes. Do not rinse with water after treatment with sodium hypochlorite solution. Do not soak overnight. Addition of one fluid oz. (2 fl. ozs.) of this sodium hypochlorite solution per 5 gallons of water will provide approximately 200 ppm available chlorine by weight.

**NOTE:** This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

**STORAGE AND DISPOSAL:** Store in a cool, dry area away from direct sunlight. In case of spill, flood area with large quantities of water. Rinse empty container thoroughly with water and either return to manufacturer or discard by placing in trash collection or burying in an approved landfill. Product or rinseate that cannot be used, should be diluted with water and disposed of in a sanitary sewer. Do not contaminate food, or feed by storage, disposal or cleaning of equipment.

ACCEPTED  
42263-G  
OCT 4 1979  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO.

**FORMATION**

er soluble liquid to be sprayed undiluted  
neral herbicide for control of brush and  
particularly useful for control of perenn  
and may not become apparent until the  
he of the weeds by moisture. The degree  
ith the amount of chemical applied, soil

lative, non-flammable, and non-corrosive  
with aluminum spray nozzles.

**IONS**

**DISPOSAL**

by storage or disposal.  
C). Store in a cool dry place away from  
aled when not in use.  
re toxic. Improper disposal of excess  
ation of Federal Law. If these wastes  
o label instructions, contact your state  
cy of the Hazardous Waste represen-  
ce for guidance.  
alent). Then offer for recycling or recon-  
sanitary landfill, or by other procedures

st before or during the period of active  
hfall can be expected for soil activation.  
mproved if vegetation is removed before  
rozen.  
niformly apply the product. Use a fixed-  
o a constant speed and rate of delivery.  
e careful to obtain an even distribution  
all areas, a tank-type hand sprayer may

**CONTROL**

riod of time on noncropland areas such  
RIGHT-OF-WAYS, PETROLEUM TANK  
EAS, and INDUSTRIAL PLANT SITES:  
eed Killer per acre to control ANNUAL  
ss, cheatgrass, crabgrass, foxtail, lamb-  
weed, ryegrass, turkey mullein, and wild  
mergence of annuals, rates as low as  
s and grasses in low rainfall areas and  
areas.  
eed Killer per acre to control PEREN-  
er, bahiagrass, bluegrass, broomsedge,  
purpletop, quackgrass, redtop, smooth  
or seasonal rainfall, rates as low as 20  
eds and grasses.

(Continued on right panel)

**BRO-KIL III  
WATER SOLUBLE LIQUID**

Active Ingredient:  
Lithium salt of bromacil (5-bromo-3-sec-butyl-6-methyluracil) ..... 7.5%  
Inert Ingredients: ..... 92.5%

U.S. Patents 3,235,357 & 3,352,862  
EPA Reg. No. 30948-32 EPA Est. No. 30948-TN-1

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING!** Harmful or fatal if swallowed. Combustible.  
Causes eye irritation. May irritate nose, throat, and skin.  
Do not get in eyes. Avoid contact with skin and clothing. Avoid breathing  
spray mist. Keep away from heat and open flame. Wash thoroughly after  
handling.  
In case of contact, flush eyes or skin with plenty of water; for eyes, get  
medical attention.

**IMPORTANT**

Injury to or loss of desirable trees or other plants may result from failure  
to observe the following: Do not apply, drain, or flush equipment on or near  
desirable trees or other plants or on areas where their roots may extend,  
or in locations where the chemical may be washed or moved into contact  
with their roots. Do not use on lawns, walks, driveways, tennis courts, or  
similar areas. Prevent drift of spray to desirable plants. Do not contaminate  
domestic waters. Keep from contact with fertilizers, insecticides, fungicides,  
and seeds. Thoroughly clean all traces of 7.5% Bromacil Liquid Weed Killer  
from application equipment immediately after use. Flush tank, pump, hoses,  
and boom with several changes of water after removing nozzle tips and  
screens (clean these parts separately).

Product will freeze at temperatures below 32° F.  
Keep Container Closed.

In Accordance with EPA Labeling Rule 40.2,  
Based on Draft Labeling Method  
NET CONTENTS \_\_\_\_\_ GALLONS

Manufactured By  
**BIONOMICAL CHEMICALS AND SERVICES, INC.**  
1003 Pineville Road  
Chattanooga, TN 37405

2052

(Continued from left panel)

Apply 16 to 32 gals. 7.5% Bromacil Liquid Weed Killer per acre to control  
JOHNSONGRASS; use at the same rate for OTHER HARD-TO-KILL PERENNIAL  
WEEDS and GRASSES such as bermudagrass, bouncingbet, bracken fern,  
dallisgrass, dogbane, horsetail, nutsedge, saltgrass, and vaseygrass. Where limited  
rainfall (usually less than 4 inches) occurs during the active growth period, such  
as some areas of the West, 7.5% Bromacil Liquid Weed Killer will not provide  
satisfactory control of hard-to-kill, deep-rooted perennial weeds such as  
johnsongrass.

Use the higher levels of the dosage ranges on adsorptive soils (usually those high  
in organic matter or carbon).

Retreatment—Apply 4 to 8 gals. 7.5% Bromacil Liquid Weed Killer per acre when  
annual weeds and grasses reappear on sites where weed growth has been  
controlled.

For small areas—1 gal. per 1,000 sq. ft. is approximately 44 gals. per acre.

**BRUSH CONTROL**

To control undesirable woody plants on noncropland areas such as RAILROAD  
RIGHT OF WAYS, STORAGE AREAS, INDUSTRIAL PLANT SITES, and DRAINAGE  
DITCHES:

Apply in spring or summer as a broadcast or basal (soil) treatment; for use on  
drainage ditches, apply as a basal (soil) treatment only.

NOTE: For effective brush control and prevention of damage to desirable vegeta-  
tion, do not apply to brush standing in water, do not use water from treated ditches  
for irrigation; do not use in irrigation ditches nor on right-of-ways or other sites where  
marketable timber or other desirable trees or shrubs are immediately adjacent to  
the treated areas.

Broadcast Treatment—Apply 6 to 13 gals. per acre to control oak, pine, sweet  
gum, and willow, apply 48 to 96 gals. per acre to control brush such as American  
elm, cottonwood, hackberry, sumac, and winged elm. Use the higher levels of  
the dosage ranges on adsorptive soils (those high in organic matter or carbon).

Basal (Soil) Treatment—Apply at the rate of 2/3 to 1-1/3 fl. oz. per stem 2" to 4"  
in basal diameter; wet base of stem to run-off. Treatment controls woody plants  
such as cottonwood, hackberry, maple, oak, poplar, red bud, sweet gum, wild cherry,  
willow, and winged elm.

**NOTICE OF WARRANTY**

Seller warrants that this product conforms to the chemical description on the  
label thereof and is reasonably fit for purposes stated on such label only when  
used in accordance with directions under normal use conditions. It is impossi-  
ble to eliminate all risks inherently associated with use of this product. Ineffec-  
tiveness or other unintended consequences may result because of such fac-  
tors as weather conditions, presence of other materials, or the manner of use  
or application, all of which are beyond the control of the seller. In no case shall  
the seller be liable for consequential, special, or indirect damages resulting  
from the use or handling of this product. All such risks shall be assumed by  
the buyer. Seller makes no warranty of merchantability or fitness for a particular  
purpose nor any other express or implied warranty except as stated above.

NOTICE TO BUYER: Purchase of this material does not confer any rights under  
patents of countries outside of the United States.