



Quality Through Research and Development

FOOD PLANT FOGGING INSECTICIDE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use against flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths.

Close windows and doors and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist of fog. In Official Meat and Poultry Establishments use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 5 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Ventilate the treated area and ventilate before reoccupying. Repeat treatment as necessary. For use in other than Official Meat and Poultry Establishments a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, chorioidea, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, mealworms, grain mites, and cadellies. Use a good sprayer adjusted to deliver a fine mist spray. Direct the spray into hiding places, cracks and crevices, under objects, around containers of stored foods, around the base of machinery, behind shelves and drawers, first and second spray bookcases. For ants spray trails, nests and colonies. If surface application only is to be used, spray floors, walls, and other surfaces at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make a liberal application to floor and baseboards. Spray directly into cracks, crevices, and under areas of shelving. Repeat treatment as necessary.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Make beds apart from walls and joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray treated areas, pet beds, and the area around pet beds and drawers, along and behind baseboards, moldings, window and door frames, and localized areas of carpet and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with approved acaricides is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles and thoroughly brush thoroughly, and air for 24 hours in sunlight. Apply spray liberally to empty chests, closets, trunks, and other storage containers. Direct the spray into cracks, joints, and crevices. After a time infested articles may be treated lightly keeping in mind at least three feet from fabric to avoid staining. Store clothes in moth light containers or repeat treatment monthly.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy areas, stagnant pools or ponds and shore lines. Apply at a rate of 1/2 fluid ounce per acre (0.2 fluid ounce per 1000 square feet).

Using truck mounted conventional misting equipment traveling at 5 m.p.h. to treat a 300 foot swath, calibrate equipment to deliver 0.23 fluid ounces per minute.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist

ACTIVE INGREDIENTS:

Pyrethrin	0.5%
Piperonyl Butoxide Technical	90%
Petroleum Distillate	0.5%
	100.0%

Equivalent to 4.0% (butylcarbityl) (5-propylpiperonyl) ether and 1.0% related compounds.

PYRENONE — Registered trademark of Fairfield Ammann Corporation

CAUTION KEEP OUT OF REACH OF CHILDREN

MFG. BY EPA REG. NO. 42458-12 EPA EST. NO. 42458-PA-01

QUALITY RESEARCH LABS, INC.
1000 FORD ST. S.W. ATLANTA, GA 30310

BEST DOCUMENT AVAILABLE

BEST DOCUMENT

CONTAINING
Triplicate
for
disposal
by

QRL

Apply with Equipment Designed and Adjusted to Deliver a True Approx. Min.

Pyrethrins	0.5%
Piperonyl Butoxide Technical	50%
Petroleum Distillate	

*Equivalent to 4-(4-butylcarbonyl) (5-propylperonyl) ether and 1,1,1-trichloro-2,2,2-trifluoroethane related compounds

PYRENONE -- Registered trademark of Fairfield American Corporation

PYRENONE -- Registered trademark of Fairfield American Corporation

CAUTION KEEP OUT OF REACH OF CHILDREN

MFG. BY: EPA REG. NO.: 42458-12 EPA EST. NO.: 42458-PA-01

QUALITY RESEARCH LABS, INC.

1366 FORD RD. #100
EVERETT LABS, INC.
CORNWELL HEIGHTS, PA 19024

PRECAUTIONARY STATEMENTS
- HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Avoid breathing vapors. Avoid contact with eyes. In case of contact, immediately flush eyes or skin with lots of water. Obtain medical attention if irritation persists. If swallowed, do not induce vomiting. Call a physician immediately.

Do not only directly feed the animals; feed them with a variety of different feeds and feed stores to avoid monotony and overfeeding. Do not give them fat. Turn out the animals. Allow them to graze in feed processing plants. Wash all equipment, tanks, troughs, etc. before exposing them to the animals, with special care. At feed processing surfaces and utensils should be disinfected by treatment with 10% caustic soda solution.

[illegible]

CONTAINER DISPOSAL:

Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in sanitary landfill, or by other state and local procedures.

EATC

BEST DOCUMENT AVAILABLE

1062

42458