


Reg # 42463-33

PM - 16

10/3/89

191



DIRECTIONS

It is a Violation of Federal Law To Use This Product in A Manner Inconsistent With its Labeling.
Apply This Product Only As Specified On This Label.

FOR PROTECTION OF STORED GRAINS (wheat, oats, corn, rye, barley, grain sorghum, and field or garden seeds) against confused flour beetles, rice weevils, Granary weevils, Saw-toothed grain beetle, Flat grain beetle, Red flour beetle, Rusty grain beetle, Lesser grain borers and Indian meal moths - Apply as the grain is being loaded or turned into final storage at the rate of 1/2 pint of "D-H" Malathion 57% E.C. in 2 to 5 gallons of water per 1,000 bushels.

To protect stored grains from attack by Indian meal moths, apply to the surface of clean or uninfested grain 1/2 pint of "D-H" Malathion 57% E.C. in 1 to 2 gallons of water per 1,000 square feet of grain surface area. Apply immediately after grain is loaded into storage and repeat application if necessary.

For Residual Wall, Floor and Machinery spray in Grain Elevator, in treating truck beds and box cars, before loading grain. Apply 1/2 gallon of "D-H" Malathion 57% E.C. per 25 gallons of water making thorough application. Before applying spray, clean elevators, box cars, etc. thoroughly. Remove and burn all sweepings and debris.

BEEF CATTLE: Lice, Ticks, Hornfly - Mix 1 to 2 gallons per 100 gallons of water. Apply as a complete coverage spray. Do not apply to dairy animals or calves under one month of age.

SWINE: Lice, Ticks - Mix 1 gallon per 100 gallons of water.

POULTRY: Northern Fowl Mite, Poultry Lice - Mix 2 tablespoons per one gallon of water. Apply one gallon of mixture per 100 to 150 birds.

POULTRY HOUSES: Northern Fowl Mite, Chicken Red Mite, Poultry Lice - Mix 4 tablespoons per gallon of water. Apply 3 gallons of mixture per 100 sq. ft. to walls, ceilings, roosts, nests and adjacent areas and taking care to force the spray into all cracks and crevices. For control of Poultry Ticks use at rate of 7 ounces per gallon of water.

MOSQUITOS: Outdoors - Use 2% to 5% Malathion fog or space spray. Dilute 1 to 25 parts of water, fuel oil or diesel oil. **MOSQUITO LARVAE** in standing water - use one pint per acre. Mix in sufficient water or oil to obtain even coverage. Contamination of shallow, fish bearing waters at application rates of 8 ounces per acre will kill fish.

CABBAGE, COLLARDS, BROCCOLI, KALE, MUSTARD, TURNIPS: Aphids, imported Cabbage Worms, Cabbage Loopers, Harlequin Bug - Use at the rate of 2 pints per acre in sufficient water for good coverage. Repeat as necessary. Do not apply within 7 days of harvest.

CUCUMBERS, EGGPLANTS, PEPPERS, SQUASH: Aphids, Spider Mites - Use 1 1/2 pints per acre in sufficient water for good coverage. Make thorough full coverage sprays. Repeat as necessary. Do not apply to cucurbits unless plants are dry. Do not apply within 3 days of harvest.

BARLEY, OATS, RYE, WHEAT: Greenbugs, English Grain Aphids, Army Worm - Use at the rate of 2 pints per acre in sufficient water for good coverage. Do not apply within 7 days of harvest.

TOBACCO: Aphids - Use at the rate of 1 1/2 pints per acre.

D-H 57% MALATHION

SAMPLE LABEL

ACTIVE INGREDIENTS:

Malathion * 57%

INERT INGREDIENTS 43%

TOTAL 100%

*0, 0-dimethyl dithiophosphate of diethyl mercaptosuccinate (Compound 4049)

CAUTION KEEP OUT OF REACH OF CHILDREN

See Side Panel For Other Caution
Manufactured By
DALY-HERRING CO.
KINSTON, N. C. 28501

EPA REG. NO. 42463-33
EPA EST. NO. 42463-NC-1
(330889)

Net Contents: One Gallon



Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

CAUTION KEEP OUT OF THE REACH OF CHILDREN

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: Remove from exposure. Keep patient quiet. Get medical attention.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Rinse eyes with water for 15 minutes. Get medical attention.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

ENVIRONMENTAL HAZARDS—This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

Highly toxic to fish. Do not contaminate any body of water, by direct application, cleaning of equipment, or disposal of wastes and containers.

Storage and Disposal: Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry area away from water, feed and foodstuffs.

CONTAINER DISPOSAL—Metal Containers—Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Glass Containers—Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state and local procedures.

Pesticide Disposal—Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.