



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

DANGER: Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS:
STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc., or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

manufactured by

**BIG A POOL SERVICE
533 N.E. 11th Avenue
Ft. Lauderdale, FL 33301**

**EPA REG. NO. 42468-1 EPA EST. NO.
NET CONTENTS**

**SODIUM
HYPOCHLORITE
SOLUTION**

**"FOR SWIMMING POOL CHLORINATION
AND SANITIZING."**

ACTIVE INGREDIENT..... 9.2%
SODIUM HYPOCHLORITE
INERT INGREDIENTS..... 90.8%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel

**DIRECTIONS
GENERAL CHLORINATION**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH THE LABEL DIRECTIONS.

DIRECTIONS FOR SWIMMING POOL CHLORINATION:
For a new pool or for spring start-up, add this sodium hypochlorite solution for equivalent to 5 ppm available chlorine by weight to the pool water.

For Pool Maintenance:
A) Adjust pool water pH to 7.6 - 7.8.
B) Add manually or by a feeder device to maintain an available chlorine residual of this sodium hypochlorite solution of 1.0 ppm available chlorine by weight. 1.0 ppm available chlorine will depart from the pool water within 24 hours. Use test kit to make certain the pH.

DIRECTIONS FOR DISINFECTING

Disinfect this sodium hypochlorite solution to 9 parts softened water. Mix thoroughly and begin feeding of solution to maintain a free available chlorine residual throughout the distribution system, as required by the National Interim Primary Drinking Water Act. Contact your local Health Department with a DPD chlorine test kit.

DIRECTIONS FOR SANITIZING FOOD SERVICE EQUIPMENT:
Clean equipment in the normal manner. Then contact with this sodium hypochlorite solution for 15 minutes. Rinse thoroughly with water. After treatment with sodium hypochlorite, one fluid oz. (2 tablespoons) of this sodium hypochlorite solution provides approximately 200 ppm available chlorine.

NOTE: This product degrades with age. As necessary, to obtain the required level of available chlorine, it may be necessary to add more product. **STORAGE AND DISPOSAL:** Store in original container, away from direct sunlight. In case of spill, fill empty container thoroughly with water by placing it in a collection container. Do not use. Do not pour into sewer. Do not contaminate food or feed by storage or disposal.

37

CERTIFIED

APR 1980

2468-1

FEDERAL INSECTICIDE
AND RODENTICIDE ACT
EPA PESTICIDE REGISTRATION NO.

STATEMENTS TO HUMANS DOMESTIC ANIMALS

Irritative, may cause severe skin irritation, broken skin. Causes eye damage. Do not get on clothing. Wear goggles or face shield when handling this product. Wash after use. Vacate poorly ventilated areas. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish, birds, and aquatic life. Do not discharge into lakes, streams, ponds or public water supply. For guidance, consult the Environmental Protection Agency.

CHEMICAL HAZARDS:

WARNING AGENT: Mix only with water. Do not mix with urine, etc., or with ammonia, acids, alkalis, or other chemicals. Mixing may release hazardous gases and mucous membranes.

Manufactured by
POOL SERVICE
E. 11th Avenue
Jerdale, FL 33301

EPA EST. NO.

CONTENTS:

SODIUM HYPOCHLORITE SOLUTION

"FOR SWIMMING POOL CHLORINATION AND SANITIZING."

ACTIVE INGREDIENT: SODIUM HYPOCHLORITE 9.2%
INERT INGREDIENTS 90.8%

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT WITH ITS LABELING

DIRECTIONS FOR SWIMMING POOL CHLORINATION

For a new pool or for spring start-up, superchlorinate with one (1) pint (16 fl. oz.) of this sodium hypochlorite solution for each 2500 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.

For Pool Maintenance:

A) Adjust pool water pH to 7.6 - 7.2 range and maintain.

B) Add manually or by a feeder device this sodium hypochlorite solution at a rate to maintain an available chlorine residual of 0.6 to 1.0 ppm. One-half pint (8 fl. oz.) of this sodium hypochlorite solution for each 6000 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.6 to 1.0 ppm available chlorine will depend upon temperature and number of swimmers. Use test kit to make certain the pH and chlorine residual are in the proper range.

DIRECTIONS FOR DISINFECTION OF POTABLE WATER FOR HOME WELL WATER SYSTEMS

Dilute this sodium hypochlorite solution in the ratio of one part sodium hypochlorite solution to 9 parts softened water. Mix sodium hypochlorite solution and water thoroughly and begin feeding of solution with a hypochlorinator (metering pump). Maintain a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm throughout the distribution system, as determined by a DPD chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations*. Check water frequently with a DPD chlorine test kit.

*Contact your local Health Department for further details.

DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT

Clean equipment in the normal manner. Just before using, rinse all surfaces thoroughly with this sodium hypochlorite solution containing 200 ppm available chlorine. Maintain contact with disinfectant for a minimum of two minutes. Do not rinse with water after treatment with sodium hypochlorite solution. Do not soak overnight. Addition of one (1) qt. (2 fl. oz.) of this sodium hypochlorite solution per 4 gallons of water will provide approximately 200 ppm available chlorine by weight.

NOTE: The product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

STORAGE AND DISPOSAL: Store in a cool, dry area away from direct sunlight. In case of spill, flood area with large quantities of water. Rinse empty container thoroughly with water and either return to manufacturer or discard by placing in trash collection or burying in an approved landfill. Product or rinsate that cannot be used should be diluted with water and disposed of in a sanitary sewer. Do not contaminate food or feed by storage, disposal or cleaning of equipment.