

42519-3, PM-23, 1/8

Similar to 56534-53 -41

LUXEMBOURG PAMOL

MSMA 6

MSMA Liquid Plus Surfactant

For Selective Post-Emergent Weed Control

ACTIVE INGREDIENT:


Monosodium Acid Methanearsonate*.....47.6%

INERT INGREDIENTS.....52.4%

TOTAL.....100.0%

Total Arsenic (as elemental) all in water soluble form.....22.03%

*Product contains 6lbs MSMA per gallon.



KEEP OUT OF REACH OF CHILDREN
CAUTION

See Side Panel for Additional Precautionary Statements.

EPA Registration No. 42519-3
EPA Establishment No. 42519-IS-1

Net Contents: 2.5 gallons
9.45 liters

Manufactured By:

Pamol, Inc.
P.O. Box 13
Tel Aviv, Israel

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin and clothing. Avoid breathing (dust) or spray mist. Wash hands and face before eating or smoking. Wash thoroughly after handling. Remove contaminated clothing and wash before reuse. Do not contaminate feed and food stuffs. Keep children and domestic animals off treated areas until this material has been washed into the soil. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a Physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Immediately flush eyes with plenty of water and get medical attention.

IF ON SKIN: Immediately wash skin with plenty of soap and water while removing contaminated clothing and shoes. If irritation develops, seek medical attention.

IF INHALED: Remove victim to fresh air.

DO NOT enter treated areas without protective clothing until sprays have dried.

Protective Clothing Requirements:

Wear protective clothing when handling or applying this product including long pants, long-sleeve shirt, and impermeable gloves and boots. Mixer-loaders should include an apron and full-face shield when handling or mixing concentrate. Flagmen should be fully protected during spray operations or mechanical flagmen used. Pilots and ground spray rig applicators should wear a mask or respirator approved by the Mining Enforcement and Safety Administration and the National Institute for Occupational Safety and Health.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

DIRECTIONS FOR USE:

BEST AVAILABLE COPY

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Farm Worker Safety Statements

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must inform workers of areas or fields that may not be entered without specific protective clothing until sprays have dried, and

appropriate actions to take in case of accidental exposure, as described under Precautionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. CAUTION. Area treated with MSMA / on [date of application]. Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, flush eyes or skin with plenty of water. Call physician if irritation persists. Remove and wash contaminated clothing before reuse.

BEST AVAILABLE COPY

4/8 filed 4/13/85

GENERAL INFORMATION

Pamol MSMA 6 Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. Its phytotoxic properties are quickly inactivated on contact with soil. It is a combination of herbicide and surfactant. It is unnecessary to add any surfactant to the spray solution. Best results are obtained on young actively growing weeds at air temperatures above 70°F.

MIXING INSTRUCTIONS

Pamol MSMA 6 Herbicide is completely water soluble. Fill the spray equipment tank about half full with water and add the required amount of herbicide with agitation. Finish filling the tank with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period. Although Pamol MSMA 6 Herbicide is only moderately corrosive, do not use in galvanized or aluminum equipment.

Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes. Do not feed foliage to livestock or graze treated areas.

COTTON

Pamol MSMA Herbicide is useful as a preplant and/or directed application in cotton for post-emergent control of weeds and grasses, such as johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, sandbur and puncturevine. It should be applied at the rate of 1/3 gallon per acre.

Mix 3 quarts of Pamol MSMA 6 Herbicide in 90 gallons of water. Do not add any surfactant to the spray solution. Apply spray solution at 40 gallons per treated acre. Weed control is most effective at temperatures above 70°F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre of cotton.

PREPLANT APPLICATION

If planting of cotton has been delayed and weeds have emerged, use one preplant application. Cotton may be planted immediately following the preplant application.

POST-EMERGENT APPLICATION

In cotton previously treated with one application of DSMA, a directed application of Pamol MSMA 6 Herbicide may be applied, if needed.

BEST AVAILABLE COPY

5/8

Rec'd 6/2/85

In cotton not treated with DSMA, two post-emergent directed applications of Pamol MSMA 6 Herbicide may be applied.

DIRECTED APPLICATIONS Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage, but to give good coverage of the weeds and grasses. If regrowth occurs, make a second application about 1 to 3 weeks after the first. Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following directed application, but the cotton plant will develop normally.

Make directed applications only when cotton is 3 inches high to first bloom. Do not apply after first bloom.

CITRUS - BEARING AND NON-BEARING (Except Florida)

Pamol MSMA 6 Herbicide is useful as a directed application in citrus orchards, such as orange, grapefruit, tangerine, lemon and lime orchards. It should be applied at the rate of 1/3 to 2/3 gallons per acre.

Mix Pamol MSMA 6 Herbicide at the rate of 1/3 gallon in 50 gallons of water. Apply as a directed spray in interspaces and around base of trees. Spray unwanted vegetation to just short of runoff. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spray solution to contact fruit, leaves, stems or bark. Use a shield, if necessary, for nursery plantings or young trees.

AGRICULTURAL PLANTINGS

Pamol MSMA 6 Herbicide is useful as a directed application in non-bearing vineyards and in non-bearing deciduous fruit and nut orchards, such as apple, peach, pear, plum, prune, apricot, cherry, almond and walnut orchards. It should be applied at the rate of 1/3 to 2/3 gallons per acre.

Mix Pamol MSMA 6 Herbicide at the rate of 1/3 gallon in 50 gallons of water. Apply as a directed spray in interspaces and around base of trees or vines. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year. Do not allow spray solution to contact leaves, stems or bark. Use a shield, if necessary, for nursery plantings or young trees. Do not apply around trees or vines from which fruit will be harvested within one year of treatment.

NON-CROP

Pamol MSMA 6 Herbicide is useful for control of johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, sandbur, puncturevine and certain other weeds on drainage ditchbanks, rights-of-way, fence rows, storage yards, and similar non-crop areas.

BEST AVAILABLE COPY

6/8 rec'd 6/13/85

Mix Pamol MSMA 6 Herbicide at a rate of 6 to 12 pints in 100 gallons of water. Spray unwanted vegetation at a rate of about 50 gallons of spray solution per acre. Use spray equipment that gives good low volume coverage. If regrowth occurs, reapply as required.

LAWN AND ORNAMENTAL TURF

Pamol MSMA 6 Herbicide is useful for control of dallisgrass, sandbur, bahiagrass, nutsedge, crabgrass, chickweed and woodsorrel with little or no injury to tolerant lawn grasses. On new lawns, do not treat until after three mowings. Tolerant grasses may be temporarily discolored. Zoysia, bluegrasses and bermuda are quite tolerant. Do not use on St. Augustine, or centipede.

Mow lawns 1 to 1½ inches high before treatment. Mix 1 1/3 fluid ounces (8 teaspoons) of Pamol MSMA Herbicide in 2½ gallons of water and apply to an area of 1,000 sq. ft. Spray thoroughly to wet all undesirable plants. Repeat applications, 10 to 14 days apart, may be needed for good control.

Do not apply with hose-end applicators.

BEST AVAILABLE COPY

✓ STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE INSTRUCTIONS

Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. High heat may form volatile arsenic compounds. Do not stack over cases high. Move containers by handlas or in cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed, seed, fungicides, insecticides and fertilizer to avoid contamination. Store in original containers only. If the contents are leaking or material is spilled follow these steps while wearing protective clothing:

1. Contain spill, absorb with an inert material such as sand or saw dust.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Place a leaking container in a plastic tub and transfer contents as soon as possible to an empty original container.
6. Do not allow run-off to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result.

ALL SUCH RISKS ARE ASSUMED BY THE BUYER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. Seller makes no other warranties, express or implied, including FITNESS OR MERCHANTABILITY. In no case shall the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by Seller and is accepted as such by the Buyer.

When such use and handling is not in strict accordance with directions given hereon.

BEST AVAILABLE COPY

Similar to 50534-53 ²⁰⁰ -41

LUXEMBOURG PAMOL

MSMA 6

MSMA Liquid Plus Surfactant

For Selective Post-Emergent Weed Control

ACTIVE INGREDIENT:

Monosodium Acid Methanearsonate*.....47.6%

INERT INGREDIENTS.....52.4%

TOTAL.....100.0%

Total Arsenic (as elemental) all in water soluble form.....22.03%

*Product contains 6lbs MSMA per gallon.

ACCEPTED
DEC 23 1985
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 42519-3

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side Panel for Additional Precautionary Statements.

EPA Registration No. 42519-3
EPA Establishment No. 42519-IS-1

Net Contents: 2.5 gallons
9.45 liters

Manufactured By:

Pamol, Inc.
P.O. Box 13
Tel Aviv, Israel

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin and clothing. Avoid breathing (dust) or spray mist. Wash hands and face before eating or smoking. Wash thoroughly after handling. Remove contaminated clothing and wash before reuse. Do not contaminate feed and food stuffs. Keep children and domestic animals off treated areas until this material has been washed into the soil. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a Physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Immediately flush eyes with plenty of water and get medical attention.

IF ON SKIN: Immediately wash skin with plenty of soap and water while removing contaminated clothing and shoes. If irritation develops, seek medical attention.

IF INHALED: Remove victim to fresh air.

DO NOT enter treated areas without protective clothing until sprays have dried.

Protective Clothing Requirements:

Wear protective clothing when handling or applying this product including long pants, long-sleeve shirt, and impermeable gloves and boots. Mixer-loaders should include an apron and full-face shield when handling or mixing concentrate. Flagmen should be fully protected during spray operations or mechanical flagmen used. Pilots and ground spray rig applicators should wear a mask or respirator approved by the Mining Enforcement and Safety Administration and the National Institute for Occupational Safety and Health.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

✓ **Farm Worker Safety Statements**

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must inform workers of areas or fields that may not be entered without specific protective clothing until sprays have dried, and

appropriate actions to take in case of accidental exposure, as described under Precautionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. CAUTION. Area treated with MSMA / on [date of application]. Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, flush eyes or skin with plenty of water. Call physician if irritation persists. Remove and wash contaminated clothing before reuse.

Handwritten: 11/24/85

GENERAL INFORMATION

Pamol MSMA 6 Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. Its phytotoxic properties are quickly inactivated on contact with soil. It is a combination of herbicide and surfactant. It is unnecessary to add any surfactant to the spray solution. Best results are obtained on young actively growing weeds at air temperatures above 70°F.

MIXING INSTRUCTIONS

Pamol MSMA 6 Herbicide is completely water soluble. Fill the spray equipment tank about half full with water and add the required amount of herbicide with agitation. Finish filling the tank with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period. Although Pamol MSMA 6 Herbicide is only moderately corrosive, do not use in galvanized or aluminum equipment.

Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes. Do not feed foliage to livestock or graze treated areas.

COTTON

Pamol MSMA Herbicide is useful as a preplant and/or directed application in cotton for post-emergent control of weeds and grasses, such as johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, sandbur and puncturevine. It should be applied at the rate of 1/3 gallon per acre.

Mix 3 quarts of Pamol MSMA 6 Herbicide in 90 gallons of water. Do not add any surfactant to the spray solution. Apply spray solution at 40 gallons per treated acre. Weed control is most effective at temperatures above 70°F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre of cotton.

PREPLANT APPLICATION

If planting of cotton has been delayed and weeds have emerged, use one preplant application. Cotton may be planted immediately following the preplant application.

POST-EMERGENT APPLICATION

In cotton previously treated with one application of DSMA, a directed application of Pamol MSMA 6 Herbicide may be applied, if needed.

Rec'd 4/13/85

In cotton not treated with DSMA, two post-emergent directed applications of Pamol MSMA 6 Herbicide may be applied.

DIRECTED APPLICATIONS Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage, but to give good coverage of the weeds and grasses. If regrowth occurs, make a second application about 1 to 3 weeks after the first. Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following directed application, but the cotton plant will develop normally.

Make directed applications only when cotton is 3 inches high to first bloom. Do not apply after first bloom.

CITRUS - BEARING AND NON-BEARING (Except Florida)

Pamol MSMA 6 Herbicide is useful as a directed application in citrus orchards, such as orange, grapefruit, tangerine, lemon and lime orchards. It should be applied at the rate of 1/3 to 2/3 gallons per acre.

Mix Pamol MSMA 6 Herbicide at the rate of 1/3 gallon in 50 gallons of water. Apply as a directed spray in interspaces and around base of trees. Spray unwanted vegetation to just short of runoff. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spray solution to contact fruit, leaves, stems or bark. Use a shield, if necessary, for nursery plantings or young trees.

AGRICULTURAL PLANTINGS

Pamol MSMA 6 Herbicide is useful as a directed application in non-bearing vineyards and in non-bearing deciduous fruit and nut orchards, such as apple, peach, pear, plum, prune, apricot, cherry, almond and walnut orchards. It should be applied at the rate of 1/3 to 2/3 gallons per acre.

Mix Pamol MSMA 6 Herbicide at the rate of 1/3 gallon in 50 gallons of water. Apply as a directed spray in interspaces and around base of trees or vines. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year. Do not allow spray solution to contact leaves, stems or bark. Use a shield, if necessary, for nursery plantings or young trees. Do not apply around trees or vines from which fruit will be harvested within one year of treatment.

NON-CROP

Pamol MSMA 6 Herbicide is useful for control of johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, sandbur, puncturevine and certain other weeds on drainage ditchbanks, rights-of-way, fence rows, storage yards, and similar non-crop areas.

4/2
Rec'd 6/13/85

Mix Pamol MSMA 6 Herbicide at a rate of 6 to 12 pints in 100 gallons of water. Spray unwanted vegetation at a rate of about 50 gallons of spray solution per acre. Use spray equipment that gives good low volume coverage. If regrowth occurs, reapply as required.

✓ LAWN AND ORNAMENTAL TURF

Pamol MSMA 6 Herbicide is useful for control of dallisgrass, sandbur, bahiagrass, nutsedge, crabgrass, chickweed and woodsorrel with little or no injury to tolerant lawn grasses. On new lawns, do not treat until after three mowings. Tolerant grasses may be temporarily discolored. Zoysia, bluegrasses and bermuda are quite tolerant. Do not use on St. Augustine, or centipede.

(Mow lawns 1 to 1½ inches high before treatment. Mix 1 1/3 fluid ounces (8 teaspoons) of Pamol MSMA Herbicide in 2½ gallons of water and apply to an area of 1,000 sq. ft. Spray thoroughly to wet all undesirable plants. Repeat applications, 10 to 14 days apart, may be needed for good control.

(Do not apply with hose-end applicators.

Ind 6/13/85

✓ STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE INSTRUCTIONS

Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. High heat may form volatile arsenic compounds. Do not stack over cases high. Move containers by handles or in cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture sources. Avoid storage near water supplies, food, feed, seed, fungicides, insecticides and fertilizer to avoid contamination. Store in original containers only. If the contents are leaking or material is spilled follow these steps while wearing protective clothing:

1. Contain spill, absorb with an inert material such as sand or saw dust.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Place a leaking container in a plastic tub and transfer contents as soon as possible to an empty original container.
6. Do not allow run-off to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Rec'd 6/13/55

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result.

ALL SUCH RISKS ARE ASSUMED BY THE BUYER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. Seller makes no other warranties, express or implied, including FITNESS OR MERCHANTABILITY. In no case shall the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by Seller and is accepted as such by the Buyer.

When such use and/or handling is not in strict accordance with directions given hereon.