

42519-22

6/21/2001

ACCEPTED 185

(Front Panel)

**ProPhyt™**

JUN 21 2001  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 42519-22

A systemic fungicide for the control of downy mildew, purple blotch, late blight, brown rot  
 and root rot in vegetables, field crops, vineyards and orchards

**ACTIVE INGREDIENT:**

Potassium Phosphite\* ..... 54.50%

**OTHER INGREDIENTS:** ..... 45.50%**TOTAL** ..... 100.00%

\*Phosphorous acid equivalent: 34.30% (4.2 lbs / gal.).

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**FIRST AID**

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 – 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

**See back panel for additional precautionary statements.**

EPA Reg. No.: 42519-22

Net Contents: 1 or 2.5 gallons

EPA Establishment No.: 42519-ISR-001

**Manufactured by:**

Luxembourg Industries (Pamol) Ltd.

P.O. Box 13

Tel Aviv 61000, ISRAEL

Batch No. XXXX-XX

FOR CHEMICAL SPILL, LEAK,  
 FIRE OR EXPOSURE CALL  
 TOLL FREE: 1-800-424-9300

**AGRICULTURAL CHEMICAL**  
 DO NOT SHIP OR STORE WITH FOODS, FEEDS,  
 DRUGS OR CLOTHING

(Back Panel)

### PRECAUTIONARY STATEMENTS

**Hazard to humans and domestic animals: Caution.** Harmful if swallowed or if absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):** Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks. Follow manufacturer's instructions for cleaning / maintaining PPE. Keep and wash PPE separately from other laundry.

**User Safety Requirements:** Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards:** Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when disposing of equipment washwater.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 4 hours.

PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is coveralls, waterproof gloves and shoes plus socks.

**General Information:**

ProPhyt™ is a systemic product, which contains potassium and phosphate. The phosphonic anion is effective in prevention and control of downy mildew and *Phytophthora*, by activation of the plants' natural resistance mechanism. ProPhyt™ is intended for use as part of an Integrated Pest Management (IPM) system. When used in conjunction with good cultural management practices and as part of a complete disease control program, ProPhyt™ will control downy mildew, purple blotch, late blight, brown rot and root rot in field crops, vineyards and orchards. In order to achieve maximum results in controlling the disease, spraying should be carried out before the appearance of the disease or when disease is first observed. Applications should be initiated when environmental conditions are favorable for disease development. The preharvest interval is 0 days for this product.

**Mixing and Application Instructions:**

Apply in sufficient water to ensure thorough coverage. Fill spray tank halfway with clean water. Add the required amount of ProPhyt™ to the tank as indicated in the table below. While agitating, add the remaining amount of water. Apply solution directly to foliage, unless directed otherwise in the application rate table. For optimum control, thorough coverage is required.

**Compatibility:**

ProPhyt™ is believed to be compatible with most commonly used pesticides. Consult specific product labels for additional information. It is always advisable to conduct a tank spray compatibility test when you plan to mix this product with other products. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, and then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank.

It is recommended that tank-mix combinations be used on a small number of plants before treating large areas, as crop sensitivity to these mixtures may vary.

ProPhyt™ has been evaluated for phytotoxicity on a large variety of crops under various normal field conditions. However, testing all crop varieties, in all mixtures and combinations is not feasible. We therefore recommend testing for phytotoxicity a small portion of the area to be treated, prior to treating the entire area.

### Application Rate:

Begin spraying when environmental conditions are favorable for the development of disease or as indicated below. Do not apply above the recommended rates.

Crop	Disease	Rate/Acre	Conc.	Remarks
<b>Cucurbits, such as:</b> Cucumber, melon, pumpkin, squash (in open field and under coverage)	Downy mildew	3-6 <sup>1</sup> pints	-	Make preventive foliar applications at one-week intervals with a spray volume of 40-100 gal/acre, depending on size of plants.
<b>Bulb vegetables, such as:</b> Garlic, onion, shallot	Downy mildew Purple Blotch	3-6 <sup>1</sup> pints	-	Make preventive foliar applications at one-week intervals with a spray volume of 20-40 <sup>2</sup> gal/acre depending on size of plants. <sup>3</sup>
<b>Brassica vegetables, such as:</b> Broccoli, cabbage, cauliflower	Downy mildew	3-6 <sup>1</sup> pints	-	Make preventive foliar applications at 7 to 10 day intervals. Thorough coverage is required.
<b>Fruiting vegetables, such as:</b> Potato, tomato, tomatillo	Late Blight ( <i>Phytophthora infestans</i> )	3-6 <sup>1</sup> pints	-	Make preventive foliar applications at 2 to 4 leaf stage of growth or immediately after trans-planting into the field. Apply at 5 to 10 day intervals. <sup>3</sup> Thorough coverage is required.
<b>Leafy vegetables, such as:</b> Lettuce, spinach	Downy mildew	3-6 <sup>1</sup> pints	-	Make preventive foliar applications at 7 to 10 day intervals with a spray volume of 40-100 gal/acre. <sup>3</sup> Begin spraying at 2 to 4 leaf stage of growth.
<b>Grapes, such as:</b> Carignane, French Colombard, Cabernet Sauvignon, Superior, Thompson, Parlet	Downy mildew	-	0.3%	Designed for prevention of downy mildew on the foliage. During the sensitive period, spray before the appearance of the disease, every 7-10 days. <sup>3</sup>
<b>Avocado</b>	Root rot ( <i>Phytophthora</i> )	-	0.3%	Make four preventive foliar applications with spray volumes of 200-300 gal/acre: twice (at 10 to 14 day intervals) in each of two root growth seasons (usually May and July). Avoid applying during period of foliar growth.

Crop	Disease	Rate/Acre	Conc.	Remarks
Citrus, such as: Grapefruit, kumquat, lemon, lime, orange	Brown rot ( <i>Phytophthora</i> )	-	0.25%	Make preventive foliar applications during spring and fall. Spray to wet (about 200-500 gal/acre).
	Root rot ( <i>Phytophthora</i> )	-	0.3%	Make preventive foliar applications when conditions favor disease development, usually three times annually (March-April, May-June, and Sept.-Oct.). Spray to wet (about 200-500 gal/acre).
Citrus seedlings	Root rot ( <i>Phytophthora</i> )	-	2.5%	Make preventive applications of 7 fl oz (200 ml) per plant (in 2 gallons soil). Apply every two months, during the growth period of the seedlings. Do not combine with other products.

**Notes:**

- 1 Use higher rate when conditions favor disease development and/or with increased disease pressure.
- 2 Use a low spray volume, so that the material **does not** wash off or drip to the ground.
- 3 For prevention of downy mildew and late blight in bulb, fruiting and leafy vegetables, and grapes, it is recommended to combine ProPhyt™ treatment with additional protective products, specifically with dithiocarbamates, by tank-mix or by alternating treatments.

**Important:**

Do not apply on plants when they are under water stress, or during severe weather conditions such as high or very low temperatures.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a dry, cool place.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Warranty Limitations and Disclaimer**

Luxembourg warrants that at the time of delivery, the product will conform to its chemical description on the label, that it will pass without objection in the trade under the contract description, that seller will convey good title thereto, and that such product will be delivered free from any lawful security interest, lien or encumbrance.

This is the only warranty made on this product. Luxembourg EXPRESSLY DISCLAIMS ANY IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE AND, EXCEPT AS SET FORTH IN THE ABOVE PARAGRAPH, ANY OTHER EXPRESS OR IMPLIED WARRANTIES. Buyer acknowledges the use of its own independent skill and expertise in the selection and use of the product and does not rely on any oral or written statements or representations.