



required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in parenthesis following crop.

✓ **FELLED LOGS:** Bark Beetles (Scolytid species) - 2 gals. per 100 gals.; Bark Beetles (Buprestids, C. rambyceids, Sarcophoridae) - 4 gals. per 100 gals. Apply in late Spring to felled logs. Spray logs thoroughly to the point of spray run-off.

**SUCCULENT BEANS, EXCEPT LIMAS:** Mexican Bean Beetles - 1 qt. per acre. Do not apply after pods begin to form. Do not feed treated threshings to live stock.

**BROCCOLI (7), CABBAGE (7):** Cabbage Looper, Imported Cabbage Worm, Diamondback Moth Larvae, Cross-striped Cabbage-worm, Flea Beetles, Harlequin Bugs, Stink Bugs - 3 to 4 pts. per acre.

✓ **CAULIFLOWER:** Cabbage Loopers, Imported Cabbage Worm, Diamondback Moth Larvae, Cross-striped Cabbage-worm - 1 to 1½ qts. per acre; Flea Beetles, Harlequin Bugs, Stink Bugs - 1½ qts. per acre. Do not apply after edible parts begin to form. Close trim outer leaves prior to marketing.

✓ **ORNAMENTALS, GREENHOUSE & OUT OF DOORS:** Aphids, Cyclamen Mite, Whitefly - 1 qt. per 100 gals. On chrysanthemums, best results will be obtained if applied before plants flower. Do not use on Birch Trees. Do not apply to "Bonnafon Deluxe", "Fred Shoemith" and "White Knight" chrysanthemums as injury may result.

✓ **PEACH NURSERY STOCK:** Peach Tree Borer - Use 1 gal. per 40 gals. of water. Mix thoroughly. Immerse the bundles of trees so that the roots and crowns are covered well above the grafting bud scar. If trees are not planted immediately, allow to dry before returning them to the storage cellar. Wear rubber gloves during the dripping operation.

**CUCUMBERS; SQUASH, SUMMER AND WINTER (0):** Aphids, Cucumber Beetle, Pickleworm, Melonworm, Squash Beetle, Squash Bug, Squash Vine Borer, Striped Cabbage Flea Beetles - 1 to 2 qts. per acre. For vine borer control, apply weekly to flower buds, stems and vines beginning when moths first appear.

**PEPPERS (1):** Flea Beetles, Green Peach Aphid, Pepper maggot - 1 qt. per acre.

**EGGPLANTS (1):** Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Green Stink Bug, Whitefly - 1 qt. per acre.

**LETTUCE, HEAD (14):** Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbage Worm - 3 to 4 pts. per acre. Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not feed crop refuse to live stock.

**MELONS (0):** Aphids, Cucumber Beetles, Melonworm, Pickleworm, Squash Beetle, Squash Vine Borer, Squash Bug - 1 to 2 qts. per acre.

**POTATOES (0):** Aphids, Colorado Potato Beetle, Plant Bugs (including Shield Shaped Bugs), Potato Flea Beetle, Potato Leafhopper, Potato Tuberworm, Southern Armyworm 1 to 2 qts. per acre; European Corn Borer, Potato Psyllid - 1½ to 2 qts. per acre; False Chinch Bug - 2 qts. per acre; Three-lined Potato Beetle - 1 qt. per acre.

**SHADE TREES:** Aphids - 1 qt. per 100 gals. Do not use on Birch Trees.

**ELM TREES:** Elm Leaf Beetle - **FOR USE IN NEW ENGLAND AREA ONLY.** Use 2 qts. per 100 gallons. Apply when 50% of the overwintering eggs have hatched. Repeat as necessary to maintain control.

**TAXUS:** Taxus Bud Mite - 1 qt. per 100 gals. Apply 3 to 5 applications as required beginning in mid-May. A thorough treatment of foliage, twigs and bark is essential.

**TOBACCO (5):** Flea Beetle; Hornworm, Budworm, Green Peach Aphid, Cabbage Looper - **SEED BED:** 1 qt. per 100 gallons. Apply about 6 gallons per 100 square yards. **FIELD:** 1 to 2 qts. per acre. Apply in sufficient water for uniform coverage. Stink Bugs - 2 to 3 pints per 50 gal.

**TOMATOES (Field and Greenhouse) (1):** Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Whitefly - 1 qt. per acre. Hornworms - 1 to 2 qts. per acre. Tomato Fruitworm, Yellow-striped Armyworm - 2 qts. per acre - Stink Bugs - 1½ to 2 qts. per acre. Cabbage Looper - 1½ qts. per acre.

Highly toxic to fish and wildlife. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Dealers Should Sell in Original Packages Only.

**DO NOT USE IN UNDILUTED FORM**

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

Do not store at temperatures below 10° F.

**WARNING** - Flammable. Keep away from Heat and Open Flame. Keep Container Closed. Leaking Packages Should Be Removed to A Safe Place.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of