

APR 02 1991

Carol F. DiRenzo  
Agrolinz, Inc.  
1755 Kirby Parkway, Suite 300  
Memphis, TN 38120

Dear Ms. DiRenzo:

Subject: Label Amendment - Revised Label Text  
SEE 2,4-D LV6 Low Volatile Herbicide  
EPA Registration No. 42545-38  
Your submission dated March 11, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following provisions:

- 1) Reinstate the aerial application spray volume rate under GENERAL INFORMATION as 3-5 gallons as per the accepted label (accepted on November 19, 1986). The aerial application spray volume rate of 1-5 gallons is unacceptable. Please refer to the previous Agency letter regarding SEE 2,4-D Low Volatile Ester Herbicide dated March 28, 1991 for the reasoning behind this policy.
- 2) Reinstate the recommended amount of water under CONIFER RELEASE, HERBACEOUS WEED CONTROL, and SITE PREPARATION as 8-25 gallons as per the accepted label (accepted on November 19, 1986). The deletion of the lower limit for the recommended amount of water is unacceptable.
- 3) Under WILD GARLIC AND WILD ONION CONTROL, correct the sentence "DO NOT graze dairy animals nor cut forage for hay within 7 days of application" to read "Do not graze dairy cattle within 7 days of application. Do not apply this product within 30 days of cutting grass for hay. Remove meat animals from treated areas 3 days prior to slaughter."

CONCURRENCES							
SYMBOL	H7505C						
SURNAME	D. KENNY						
DATE	4/1/91						

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

2 9 15

- 4) Add the appropriate net contents statement to the front panel.

A stamped copy is enclosed for your records. Please submit five (5) final printed copies for the referenced label, incorporating the above changes.

Please note that all future label amendment submissions must include five (5) copies of the sample label.

Sincerely yours,

Joanne I. Miller  
Product Manager (23)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosure

CONCURRENCES							
SYMBOL	H7505C						
SURNAME	D. KENNY						
DATE	4/1/91						

EPA Form 1320-1A (1/90) Printed on Recycled Paper OFFICIAL FILE COPY

3 7 15

ACCEPTED  
with COMMENTS  
EPA Letter Dated

1

AGROLINZ, INC.

2,4-D LV6

LOW VOLATILE HERBICIDE

APR 02 1991  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
42545-38

ACTIVE INGREDIENT:

Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid.....89.5%

INERT INGREDIENTS:.....10.5%

\* Equivalent to 59.4% of 2,4-Dichlorophenoxyacetic acid or 5.64 lb./gal.

Isomer specific by AOAC Method.

EPA Reg. No. 42545-38

EPA Est. No. 42545-MO-1

KEEP OUT OF REACH OF CHILDREN

**CAUTION - CAUCION**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Get medical attention. Contains petroleum distillates.

IF ON SKIN: Wash contaminated areas thoroughly with soap and water. Do not reuse contaminated clothing until washed. Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of clean water for 10 - 15 minutes. Get medical attention.

NOTE TO ATTENDING PHYSICIAN: May cause chemical pneumonitis if aspirated. If lavage is performed, suggest endotracheal and/or esophagosopic control.

PRECAUTIONARY STATEMENTS

**CAUTION - CAUCION**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. When handling this product, wear chemical resistant gloves.

BEST AVAILABLE COPY

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water except as specified on this label. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**PHYSICAL AND CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**AGRICULTURAL CHEMICAL**

Do not ship or store with foods, feeds, drugs or clothing.

**FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE  
CALL CHEMTREC (800) 424-9300**

Before buying or using this product, read "Warranty Limitations and Disclaimer" elsewhere on this label. IF terms are not acceptable, return **unopened** package at once to seller for full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under the Warranty Limitations and Disclaimer.

**WARRANTY LIMITATIONS AND DISCLAIMER**

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the DIRECTIONS FOR USE when used under normal conditions. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL. Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller. When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract,

**BEST AVAILABLE COPY**

negligence, strict liability, or other legal theories), buyer or user must promptly notify seller, in writing, of any claims to be eligible to receive either remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of seller will be one of the following, at the election of the seller:

1. Refund of purchase price paid by buyer or user for product bought or
2. Replacement of amount of product used.

The seller will not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations and Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations and Disclaimer in any manner.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.


#### REENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried. Protective clothing should include hat or other suitable head covering, long sleeved shirt and long legged trousers or a coverall type garment, shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. The front panel PRECAUTIONARY STATEMENTS should be read to workers as well as the instruction not to enter until sprays have dried. When oral warnings are given, warnings shall be given in a language customarily understood by the workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION: Area treated with 2,4-D on (date of application). Do not enter without appropriate protective clothing until sprays have dried. (Insert here Statements of Practical Treatment as on front panel.)



## STORAGE AND DISPOSAL

**STORAGE:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Do not store near heat or open flame. Reclose all partially used containers by thoroughly tightening screw cap. Absorb any spill with a suitable clay absorbent and dispose of as indicated under "Pesticide Disposal".

For safety and prevention of unauthorized use, all pesticides should be stored in locked facilities.

To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification.

Opened, partially used pesticides should be stored in original labeled containers when possible. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of the new container.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

## CONTAINER DISPOSAL

**METAL CONTAINERS:** Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**PLASTIC CONTAINERS:** Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## GENERAL INFORMATION

Performance of this product may be affected by local conditions, crop varieties, and application method. User should consult local Extension Service, Agricultural Experiment, or University Weed Specialists, and state regulatory agencies for recommendations in your area.

**BEST AVAILABLE COPY**

Best results are obtained when product is applied to young succulent weeds that are actively growing. The lower recommended rates will be satisfactory on susceptible annual weeds. For perennial weeds and conditions such as the very dry areas of the western states, where control is difficult, the higher recommended rates should be used.

When product is used for weed control in crops, the growth stage of the crop must be considered.

Some plants and weeds, especially woody varieties, are difficult to control and may require repeat applications.

Application rates should be 3 to 5 gallons of total spray by air or 5 to 25 gallons by ground equipment unless otherwise directed. In either case, use the same amount of 2,4-D recommended per acre. On crops, use of this product with oil, surfactants, or other adjuvants may reduce selectivity and cause crop damage.

Aerial applications should be used only when there is no danger of drift to susceptible crops. Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. Although this product is a low volatile formulation, at temperatures above 90°F vapors may damage susceptible crops growing nearby.

Because coarse sprays are less likely to drift than fine, do not use equipment (such as hollow cone small orifice nozzles) or conditions (such as high pressure) that produce such sprays.

Product should not be allowed to come into contact with desirable, susceptible plants such as beans, cotton, fruit trees, grapes, legumes, ornamentals, peas, tomatoes, and other vegetables. Product should not be used in greenhouses. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Users should note that herbicide treatment of public water requires a permit from appropriate state agencies in most states. Your state Conservation Department, or Game and Fish Commission will aid you in securing a permit in your state.

If stored below freezing, efficacy is not affected if product is warmed to 40°F and agitated before using.

Spray equipment used to apply 2,4-D should not be used for any other purpose until thoroughly cleaned.

**Spray Preparation:** Add the recommended amount of product to approximately 1/2 the volume of water to be used for spraying. Agitate well, then add the remainder of the water. Continue

BEST AVAILABLE COPY

agitation during application until spray tank is empty.

**Use in Liquid Nitrogen Fertilizer:** Product may be combined with liquid nitrogen fertilizer suitable for foliar application on corn, grass, pastures, or small grains in one operation. Use product according to directions on this label for those crops. Use liquid nitrogen fertilizer at rates recommended by supplier or Extension Service Specialist. Mix the product and fertilizer according to the following instructions:

Fill the spray tank approximately 1/2 full with the liquid nitrogen fertilizer. Add the product while agitating the tank. Add the remainder of the fertilizer while continuing to agitate. Apply immediately, maintaining agitation during application until tank is empty. DO NOT APPLY DURING COLD (NEAR FREEZING) WEATHER. Spray mixture must be used immediately and may not be stored.

**NOTE:** (1) If good, continuous agitation is not maintained, separation of the spray mixture and/or clogging of the nozzles is likely to occur.

**NOTE:** (2) If user's spray program includes frequent application of 2,4-D in liquid fertilizer, consideration should be given to using SEE® 2,4-D which is specially designed and formulated for such use.

#### WHERE TO USE

This product is used to control broadleaf weeds in cereal crops, corn, and sorghum; weeds, and brush in rangeland, pastures, rights-of-way, and similar noncrop uses; and for aquatic weed control.



# **WEEDS CONTROLLED**

When used properly, product will kill or control the following, in addition to many other noxious plants susceptible to 2,4-D:

Alligatorweed	Ironweed
Arrowhead	Jimson weed
Artichoke	Lambsquarters
Bindweed	Locoweed
(hedge, field	Mexican weed
and European)	Morningglory
Bitter wintercress	Mustard
Boxelder	Parrotfeather
Buckhorn	Pennywort
Bull thistle	Pigweed
Bulrush	Plantain
Burdock	Pokeweed
Bur ragweed	Povertyweed
Buttercup	Puncturevine
Canada thistle	Purslane
Catnip	Rush
Chickweed	Russian thistle
Chickory	Sagebrush
Cocklebur	Shepherdspurse
Coffeebean	Sowthistle
Common ragweed	Stinkweed
Creeping jenny	Sumac
Cudweed	Sunflower
Curly dock	Swinecress
Curly indigo	Velvetleaf
Cutleaf evening primrose	Vetch - narrow leaf
Dandelion	Virginia creeper
Dock	Waterhyacinth
Dogfennel	Waterlily
Duckweed	Waterprimrose
Elderberry	Wild lettuce
Giant ragweed	Wild radish
Goldenrod	Willow
Ground ivy	
Hemp	
Hoary cress	
Honeysuckle	
Indigo	

## **LESS SUSCEPTIBLE WEEDS**

Kochia	Smartweed
Hairy vetch	Wild garlic
Pigweed (hybrid)	Wild onion
Poison ivy	

## CROPS

**Small grains (barley, oats, wheat, rye), not underseeded with a legume:** See table for recommended use rates. Spray when weeds are small after grain begins tillering but before boot stage (usually 4 to 8 inches tall). Do not apply before the tiller stage nor from early boot through milk stage. To control large weeds that will interfere with harvest or to suppress perennial weeds, preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well.

**Spring Planted Oats:** Use 1/3 pint per acre in sufficient water to give good coverage. Apply after the fully tillered stage, except during the boot to dough stage.

**Fall Planted Oats:** Apply 1/6 to 5/6 pints per acre after full tillering but before early boot stage. Some difficult weeds may require the higher rates of 1/2 to 5/6 pints per acre for maximum control, but injury may result. Do not spray during or immediately following cold weather.

**Note:** Oats are less tolerant to 2,4-D than wheat or barley and more likely to be injured. Do not forage or graze treated grain fields within 14 days after treatment with 2,4-D. Do not feed treated straw to livestock.

**Wheat and Barley:** Control of wild garlic and wild onion. For improved control of difficult weeds including wild garlic and wild onion, apply 1/3 to 1-1/3 pints of product per acre. Since these rates may injure the crop, do not use unless possible crop damage is acceptable. For the higher rates on spring wheat and barley, consult your local State Agricultural Experiment Station or Extension Service Weed Specialist for recommendations or suggestions to fit local conditions.

### **Control of Wild Garlic in Stubble Grain Fields:**

Following the harvest of small grains, wild garlic often produces new fall growth. This should be sprayed with 1-1/3 to 2 quarts of product per acre. This is a useful practice as one part of the wild garlic control program. Do not forage for 14 days following application. Do not plant any crop for three months after treatment.

**Corn:** See table for recommended use rates.

**Preemergent:** Apply product to emerged weeds from 3 to 5 days after planting but before corn emerges. Do not use on very light, sandy soils. Use the higher rates on heavy soils. Plant corn as deep as practical. Product will not control weeds which have not emerged.

**Post Emergent:** Best results are usually obtained when weeds are small and corn is 4 to 18 inches tall. When corn is over 8 inches tall, use drop nozzles to keep spray off corn foliage as much as possible. Do not apply from tasseling to dough stage. If corn is growing rapidly and temperature and soil moisture is high, use 1/3 pint per acre to reduce possibility of crop damage. Delay cultivation for 8 to 10 days to prevent stalk breakage due to temporary brittleness caused by 2,4-D. Application rates of up to 2/3 pint per acre may be used to control some hard to control weeds. However, the possibility of injury to the corn is increased.

Since the tolerance to 2,4-D of individual hybrids varies, consult your seed supplier, local Extension Service, Agricultural Experiment Station, or University Weed Specialist for information.

**Pre-Harvest:** After the hard dough or denting stage, apply 2/3 to 1-1/3 pints of product per acre by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower, velvetleaf, and vines that interfere with harvesting. Do not forage or feed corn fodder to livestock for 7 days following application.

**Post-Harvest:** Following the harvest of corn, wild garlic often produces new fall growth. This should be sprayed with 1-1/3 to 2 quarts of product per acre. This is a useful practice as one part of a wild garlic control program. Do not forage for 7 days following application. Do not plant any crop for three months after treatment.

**Sorghum (Milo):** See table for recommended rate.

Apply sorghum when crop is 4 to 12 inches high with secondary roots well established. Use drop nozzles when crop is over 10 inches high. Do not apply from flowering to dough stage. Rates of up to 2/3 pint per acre may be used to control some hard to control weeds. However, the chance of crop injury is increased with the higher rates. Do not use with oil. Use lower rate if conditions of high temperature and high soil moisture exist.

# RECOMMENDED RATES OF PRODUCT PER ACRE\*\*

Crop (See Detailed Instructions Above)	Rate, Average Conditions	Rate, Dry Conditions as in Western States*
<u>Small Grains (wheat, barley, rye):</u>		
Annual weeds	1/3 to 2/3 pts.	2/3 to 1-1/3 pts.
Perennial weeds	2/3 pt.	5/6 to 1-1/3 pts.
Preharvest	2/3 to 1-1/3 pts.	

BEST AVAILABLE COPY

---

**Oats:**

Spring

1/3 pint

Fall

1/3 to 1/2 pint

---

**Corn:**

Preemergent

2/3 to 1-1/3 qts.

Postemergent

1/3 pint

1/3 to 1/2 pt.

Preharvest

2/3 to 1-1/3 pts.

---

**Sorghum (Milo):**

Postemergent

1/3 pint

1/3 to 1/2 pt.

---

\*Arizona, Idaho, Montana, Oregon, Utah, Washington, Wyoming

\*\*If band treatment is used, base the dosage rate on the actual area sprayed.

**Sugarcane:** Use 1-1/3 pints per acre as a preemergent application to control already emerged weeds before canes appear or 2-2/3 pints per acre as a blanket spray after can emerges and through layby. Consult local Agricultural Experiment or Extension Service Weed Specialists on specific use of this product or in combination with any other product.

**Ornamental Turf:** Use 2/3 to 2 pints of product in enough water to give good coverage to one acre on established stands of perennial grasses, depending on type of weeds and stage of growth. Do not use on creeping grasses such as bentgrass except for spot spraying. Newly seeded turf should not be treated until after the second mowing and the lower dosage rate should be used.

**Grass Seed Crops:** Apply 2/3 to 2-2/3 pints of product per acre in the spring or fall to control broadleaf weeds in grass being grown for seed. Do not apply from early boot to milk stage. Spray seedling grass only after the five leaf stage, using 1/2 to 2/3 pint per acre to control small seedling weeds. After the grass is well established, higher rates of up to 2-2/3 pints per acre can be used to control hard to control annual or perennial weeds. For best results, apply when soil moisture is adequate for good growth. Do not use on bentgrass unless injury can be tolerated. Do not graze dairy cattle within 7 days of application. Do not apply this product within 30 days of cutting grass for hay. Remove meat animals from treated areas 3 days prior to slaughter.

**Fallow Land:** On established perennial species such as Canada thistle and field bindweed, apply up to 4 pints of product per acre. For annual broadleaf weeds, apply 1-1/3 to 2-2/3 pints per acre. Do not plant any crop for 3 months after treatment.

**Established Pastures and Rangelands:** Use 2/3 to 2-2/3 pints of product in sufficient water to give good coverage to one acre

BEST AVAILABLE COPY

depending on type of weeds and stage of growth. Use only on established stands of perennial grasses. Do not graze dairy cattle within 7 days of application. Do not apply this product within 30 days of cutting grass for hay. Remove meat animals from treated areas 3 days prior to slaughter.

**Wild Garlic and Wild Onion Control:** Apply 2-2/3 to 4 pints of product per acre making three applications, fall-spring-fall or spring-fall-spring, starting in the late fall or early spring. DO NOT graze dairy animals nor cut forage for hay within 7 days of application.

**General Weed Control:** (airfields, roadsides, vacant lots, drainage ditch banks, fence rows, industrial sites and similar areas):

Use 1-1/3 to 4 pints of product per acre. Usually 2-2/3 pints per acre will give adequate control. Do not use on herbaceous ground covers or creeping grass such as bentgrass. Legumes will usually be damaged or killed. Deep-rooted perennials may require repeat applications. Do not use on freshly seeded turf until grass is well established. Delay seeding for 30 days.

**Woody Plant Control:** To control woody plants susceptible to 2,4-D, such as alder, buckbrush, elderberry, sumac, and willow on non-crop areas, use 1-1/3 to 2 quarts of product per acre in 100 gallons of water. Wet all parts of the plants thoroughly, including stem and foliage, to the point of runoff. Higher volumes of up to 400 gallons per acre are necessary where the brush is very dense and over 6 to 8 feet high. Applications are more effective when made on actively growing plants. Treatment should not be made during time of severe drought or in early fall when leaves lose their green color. Hard to control species may require re-treatment next season.

#### USES IN FOREST MANAGEMENT

**Conifer Release:** For control of alder, apply 1 to 1-1/3 quarts of product per acre in up to 25 gallons of water, and apply as a foliage spray between mid-May and mid-June.

For the control of susceptible brush species such as *Ceanothus* spp., chinquapin, madrone, manzanita, oak and tanoak. Apply 2 quarts of product per acre just before or during bud break of the conifer. To increase performance, add 2 to 4 quarts of diesel, fuel oil, kerosene, or a suitable approved agricultural surfactant at recommended label rate.

After northern conifers, jack pine, red pine, black spruce, and white spruce cease growth and "harden off" in late summer, a spray of 1 to 2 quarts of product in up to 25 gallons of water per acre may be applied by air to control certain competing hardwood species such as alder, aspen, birch, hazel and willow. Since this

treatment may cause occasional conifer injury, do not use if such injury cannot be tolerated. Consult your regional or extension forester or state herbicide specialist for recommendations to fit local conditions.

**Dormant Application (other than pine):** For the control of susceptible deciduous brush species such as alder, cascara, cherry, poplar and serviceberry, apply up to 2 quarts of product per acre in sufficient diesel, fuel oil or kerosene for good coverage. Application may be made by ground or air and should be made before conifer bud break.

**Pine Only:** Make application while pine buds are still dormant. Apply 1-1/3 quarts of product per acre in sufficient water for good coverage by air or ground equipment. Do not use this application unless some pine injury is acceptable. Use of diesel, kerosene, or other oil, or addition of surfactants to spray mix may cause unacceptable pine injury.

**Herbaceous Weed Control:** To control over-wintering susceptible weeds such as false dandelion, klamath weed, plantain, and tansy ragwort apply 2/3 to 2 quarts of product per acre in up to 25 gallons of water. Make application at rates and timing indicated above if pines are present.

For control of hazel brush and similar species in the Lake States area, apply 1-1/3 quarts of product per acre in up to 25 gallons of water, when new shoot growth of Hazel is complete.

**Site Preparation: (As Budbreak Spray)** - For control of alder prior to planting seedlings, apply 1-1/3 to 2-2/3 quarts of product per acre in up to 25 gallons of water, after alder budbreak but before foliage is 1/4 full size. Application may be made by air or ground. If desired, diesel, fuel oil or kerosene may be substituted for water as a diluent.

**(As Foliage Spray)** - For control of alder prior to planting seedlings, apply 1-1/3 quarts of product per acre in up to 25 gallons of water, after most alder leaves are full size. To increase penetration, 2 to 4 quarts per acre of diesel, fuel oil, kerosene, or a suitable approved agriculture surfactant at recommended label rates, may be added to the spray mixture.

#### **AQUATIC APPLICATIONS**

**For Aquatic Weeds in Lakes, Ponds, Drainage Ditches, and Marshes:** Use 1-2/3 to 3 pints of product in 50 to 100 gallons of water per acre. Spray to wet foliage thoroughly. Application should be made when leaves are fully developed, above water line, and plants are actively growing. Your State Conservation Department or Game and Fish Commission will assist you in determining the best time and rate for application under local conditions.

**BEST AVAILABLE COPY**

**DO NOT APPLY** to more than 1/3 to 1/2 of a lake or pond in any one month because excessive decaying vegetation may deplete oxygen content of water and kill fish.

Perennial and other hard to control weeds may require a repeat application to give adequate control.

|SEE® is a registered trademark of Agrolinz, Inc.

Manufactured by:

**AGROLINZ, INC.**

1755 Kirby Parkway, Suite 300

Memphis, TN U.S.A. 38120-4393

Phone (901) 756-4422 - Fax (901) 757-9367