



Reg # 42545-49

PM-22

file 1917

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

MAR 18 1994

J.R. FISHER
AGROLINZ INC.
6525 N. QUAIL HOLLOW RD., STE. 107
MEMPHIS, TN. 38120

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 9/21/93 Response to PR Notice 93-7
EPA Reg. No. 42545-49
CUPROXAT FLOWABLE COPPER FUNGICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Candela Ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

3917

AGROLINZ, INC.

CUPROXAT

FLOWABLE COPPER FUNGICIDE

NET CONTENTS: 1 GALLON

ACCEPTED
with COMMENTS
In EPA Letter Dated

18 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

42545-49

ACTIVE INGREDIENT:

Basic Copper Sulfate.....27.1%

INERT INGREDIENT:72.9%

Total.....100.0%

Equivalent to 1.6 pounds per gallon or 14.9% metallic copper.

KEEP OUT OF REACH OF CHILDREN

CAUTION - CAUCION

~~PRECAUCION AL USUARIO: Si usted no lee ingles, no use este
producto hasta que la etiqueta le hayo sido
explicada ampliamente.~~

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Before buying or using this product, read "Warranty Limitations and Disclaimer" elsewhere on this label. If terms are not acceptable, return unopened package at once to seller for full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under the Warranty Limitations and Disclaimer.

EPA Reg. No. 42545-49

EPA Est. No. 33526-AT-02

Manufactured By:

AGROLINZ, INC.

6525 Quail Hollow, Suite 107

Memphis, TN U.S.A. 38120-1309

Phone (901) 756-4422 - Fax: (901) 757-9367

4917

PRECAUTIONARY STATEMENTS
CAUTION - CAUCION
HAZARDS TO HUMANS (& DOMESTIC ANIMALS)

Harmful if swallowed or absorbed through the skin. May cause skin sensitization in certain individuals. Causes eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

~~Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.~~

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Waterproof gloves
3. Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high

water marks. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters.

**FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE
CALL TOLL FREE 1-800-424-9300
AGRICULTURAL CHEMICAL**

Do not ship or store with foods, feeds, drugs or clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

1. Coveralls
2. Waterproof gloves
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENT

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides 40 CFR, 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children, pets and other unprotected persons out of treated areas until sprays have dried.

REENTRY STATEMENT

~~Do not enter treated areas without protective clothing until sprays have dried. Protective clothing means, at least: a hat or other suitable head covering, a long-sleeved shirt and long-legged trousers or a coverall-type garment (all of closely woven fabric covering the body including the arms and legs), shoes and socks. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.~~

~~Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. The front panel PRECAUTIONARY STATEMENTS should be read to workers as well as the instruction not to enter until sprays have dried. When oral warnings are given, warnings shall be given in a language customarily understood by the workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:~~

~~CAUTION: Area treated with Basic Copper Sulfate on (date of application). Do not enter without appropriate protective clothing until sprays have dried. (Insert here Statements of Practical Treatment as on front panel).~~

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Store tightly closed container, out of reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Use Cuproxat as noted below. Cuproxat is adaptable to spraying from all types of spray equipment. Depending on the equipment used and the specific crop (amount of foliage to be covered), the volume applied per acre will differ. For best results, thorough coverage is essential. For dilute, high volume sprays: use from 25 to 100 gallons of water per acre (GPA) for most vegetable and row crops, 400 to 800 GPA for fruit orchards, and up to 1500 GPA as may be required for large citrus groves. For concentrate ground sprays, apply from 5 to 20 GPA for most field crops; 25 to 100 GPA for fruit and nut crops. For aerial spraying, 3 to 15 GPA are commonly used. No additional surfactants are needed. Add Cuproxat slowly to a spray tank partially filled with water. Spreader-stickers, insecticides, nutrients, etc., should be added last. Cuproxat is compatible with many commercially formulated insecticides and fungicides. Since the number of combinations of pesticide mixtures is very large, it is advisable to check for compatibility and apply mixtures as soon as possible. Observe all cautions and limitations on labels of all products used in mixtures. The following specific instructions are based on general applications. The recommendations of State Agricultural Extension Services should be closely followed as to timing, frequency and number of sprays per season.

HOMEOWNER USE: One (1) level teaspoon of Cuproxat per gallon of water is equivalent to one (1) pint of Cuproxat per 100 gallons of water. Make application according to specific instructions given below for those crops where rates are given in pints/hundred gallons. (For example: 2 pints Cuproxat per 100 gallons are equivalent to 2 level teaspoons Cuproxat per gallon.)

Use hydrated lime at the rate of 1 teaspoon to each gallon of spray solution for each lb/100 gallons recommended. (For example: A recommended rate of 4 lbs of hydrated lime/100 gallons of spray would be equivalent to 4 teaspoons per gallon of spray solution prepared.)

Spray both the upper and lower leaf surfaces to the point of runoff. Use the highest label rates when weather conditions conducive to severe disease pressure exist.

CHEMIGATION INSTRUCTIONS

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (sheet) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

IF IRRIGATION SYSTEM IS CONNECTED TO A PUBLIC WATER SYSTEM, THE FOLLOWING SAFETY DEVICES MUST BE IN PLACE IN ADDITION TO THE REQUIREMENTS LISTED ABOVE:

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical

break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

DIRECTIONS FOR CHEMIGATION USE

It is recommended that the pesticide supply tank be equipped for continuous agitation by either recirculation or a mechanical agitator.

Mixing Instructions:

Fill the supply tank with approximately one half of the amount of water to be used for dilution. With agitation add any emulsifiable concentrate to be used, including any oils. Agitate thoroughly. Next add any spreader-sticker or other adjuvant and agitate thoroughly. Then add the Cuproxat and any other flowable. Agitate thoroughly. Finally, add any wettable powder or dry flowable and agitate thoroughly. With agitation add the remainder of the water to be used for application. Continue agitation during application until supply tank is empty. Observe all precautions and limitations on the labels of all products used in the mixture.

Application:

For fixed position irrigation systems such as center pivot, big gun, etc., the pesticide should be applied towards the end of the irrigation period. Exact timing will depend on the desired pesticide application rate and calibration of the system being used. For moving systems, the pesticide should be applied continuously. In all cases, careful attention should be paid to thorough coverage of the crop during application.

Do not apply when wind speed favors drift beyond the area intended for treatment.

RECOMMENDED USES

ALMONDS: Brown Rot Blossom and Twig Blight (Monilinia): Use 2-2/3 to 4 pints Cuproxat for each 100 gallons of spray mixture during the early bloom stage (popcorn). A second application in late dormancy, before foliage buds swell, may be necessary when frequent rainfall occurs. Do not spray after 50% bloom or when trees are in leaf.

Bacterial Blast (Pseudomonas): Use 4 to 5-1/3 pints Cuproxat for each 100 gallons of spray mixture at dormant to early pink bud. For blast control in sprinkler irrigated orchards or where disease is severe, apply 2 to 4 Cuproxat sprays at the rate of 2/3

pint of product per 100 gallons of spray mixture. Apply at 2-week post-bloom intervals or just before sprinkling. Slight leaf injury may occur from post-bloom spray.

APPLES: Anthracnose: Use 4 to 5-1/3 pints Cuproxat for each 100 gallons of spray mixture. For red varieties, apply to foliage before or after harvest. For yellow varieties, apply to foliage after harvest once every 2 to 3 years as necessary.

Bitter Rot, Black Rot, Blotch, Powdery Mildew: 1/2 to 1-1/2 pints of Cuproxat plus 3 to 5 pounds hydrated lime for each 100 gallons of spray mixture. Begin at petal fall and repeat through fourth cover spray.

Bulls-eye Rot: Use 4 pints Cuproxat with a suitable spray oil for each 100 gallons of spray mixture. Apply after harvest.

Fire Blight: 1/3 to 1/2 pints Cuproxat for each 100 gallons of spray mixture; 2 pints of Cuproxat/acre in concentrate sprayers. Begin at 10 percent bloom. Repeat at 5 day intervals until late bloom is over.

CAUTION: Phytotoxicity may occur from late application. (Discontinue use when green-tip is one-half inch.)

APRICOTS: Coryneum Blight (Shot Hole) and Blossom Brown Rot: Apply at popcorn to full bloom using 2-2/3 to 5-1/3 pints Cuproxat for each 100 gallons of spray mixture as a full cover spray. To avoid spray injury, do not apply after bloom.

Scab: Apply 2-2/3 to 4 pints Cuproxat for each 100 gallons of spray mixture. For additional recommendations, consult state extension service specialists.

ARBORVITAE: Blight, Leaf Spots: See Ornamental Plants and Trees.

ASTERS: Leaf Spots, Blights: See Ornamental Plants and Trees.

AVOCADOS (FLORIDA): Scab: Apply at a rate of 2-2/3 pints Cuproxat in 100 gallons spray mixture. When bloom buds begin to swell, make the first 5 applications at four (4) week intervals. Continue applications at monthly intervals for 5 to 6 additional applications. Do not apply later than 120 days after bloom. Consult state agricultural experiment stations for additional recommendations.

BEANS: Anthracnose, Angular and other Leaf Spots, Downy Mildew, Bacterial Blight (Halo and Common): Use 1-1/3 to 4 pints per acre. First application when plants are six inches high. Apply on 7 to 14 day schedule, depending on local conditions.

BEETS: Leaf Blight and Spot Downy Mildew: Begin sprays of 2-2/3 to 6-2/3 pint./acre when diseases appear; repeat every 7 to 10 days.

BLACKBERRIES: (Santiams, Logons, Boysens, Marions, Auroras, Cascades, Chehalems and Thornless Evergreens): **Leaf and Cane Spot:** Use 2 pints for each 100 gallons of spray mixture. Spray first when leaf buds swell, again when flowers begin to open.

Anthracnose: Use 2 pints of Cuproxat for each 100 gallons of spray mixture. Spray first when leaf buds begin to open, again when flower buds turn white. Repeat at 10-14 day intervals. Post-harvest, after pruning but before fall rains, apply 5-1/3 pints of Cuproxat with 1 quart superior-type oil for each 100 gallons of spray mixture. Dormant spray in spring after training, apply 5-1/3 pints with 1 quart superior-type oil for each 100 gallons of spray mixture.

BOXWOOD: Wilt: See Ornamental Plants and Trees.

BROCCOLI, BRUSSELS SPROUT, CABBAGE, AND CAULIFLOWER: Downy Mildew: In planted beds, apply 2/3 to 1-1/3 pints/acre in a minimum of 25 gallons of spray mixture per acre before diseases appear. Repeat at 7 to 10 day intervals, as needed. (CABBAGE ONLY), **Black Rot** (*Xanthomonas*) & **Black Leaf Spot** (*Alternaria*). Apply at 2/3 pints per acre in a minimum of 25 gallons of spray mixture per acre at 7 to 10 day intervals. (Caution: A slight reddening of older leaves may occur on broccoli and a slight flecking of wrapper leaves may occur on cabbage at the 2/3 pint rate). For control of diseases of these crops, begin applications after transplants are set in the field, or shortly after emergence of field-seeded crops or when conditions favor disease development.

CARNATIONS: Botrytis Blight, Leaf Spots, Leaf Spots, Leaf Gall: See Ornamental Plants and Trees.

CARROTS: Leaf Spot, Carrot Blight (Cercospora): When disease threatens, apply 2-2/3 pints per acre at 7 to 14 day intervals depending on disease severity.

CELERY: Early, Late, and Bacterial Blights: Apply as soon as plants are established in the field at 2-2/3 pints per acre, then every 5-7 days, depending on disease severity and weather. One to two quarts of a suitable agricultural spray oil per acre may be used as spreader-sticker.

CHERRY: Dead Bud: Apply 8 pints Cuproxat plus 1 pint superior-type oil for each 100 gallons of spray mixture in October (before heavy fall rains) and again in January. In orchards where the disease is severe, a spray should also be applied in August.

Brown Rot: 2-2/3 to 4 pints Cuproxat for each 100 gallons of spray mixture. Apply in red bud to popcorn and late bloom stages.

Leaf Spots: Use 2-2/3 to 4 pints Cuproxat plus 4.0 pounds of hydrated lime for each

100 gallons of spray mixture. Spray at petal fall and make 1 or 2 harvest applications.

Shot Hole: 2-2/3 to 4 pints Cuproxat for each 100 gallons of spray mixture. Apply as a dormant spray.

CHRYSANTHEMUMS: Botrytis Blight, Downy Mildew, Leaf Spots: See Ornamental Plants and Trees.

CITRUS: Melanose, Scab, Greasy Spot and Pink Pitting: Apply as pre-bloom and post-bloom sprays. Use 1 to 2 pints Cuproxat for each 100 gallons of spray mixture, depending on disease severity. May be used in concentrate sprays at equivalent rates. For aerial application, use 8 pints Cuproxat per 10 gallons of spray mixture per acre.

Brown Rot: Use 2/3 to 1-1/3 pints Cuproxat for each 100 gallons of spray mixture as dilute spray or at equivalent rate as concentrate spray, depending on severity of disease. Make a single application in fall before or just after first heavy rains. Apply also to bare ground on foot beyond skirt. (California only: In areas subject to copper injury, add 1/3 to 1 pound of high quality lime per 2/3 pints of Cuproxat.)

CRANBERRY: Fruit Rot: Apply at 10-2/3 pints per acre beginning in late bloom. One or two additional applications made at 10 to 12 day intervals may be required, depending on disease pressure. Follow the advice of the state agricultural extension service.

CUCURBITS: (Cantaloupe, Cucumbers, Honeydew Melons, Muskmelons, Pumpkins, Squash, Watermelons): Angular Leaf Spot, Anthracnose, Bacterial Wilt, Downy and Powdery Mildew, Gummy Stem Blight, other Leaf Spots and Scab: Apply 2 to 2-2/3 pints/acre. Start spraying when plants emerge or begin to vine, or before disease appears. Repeat at 7 to 10 day intervals. Spray to protect crop at all stages of development. Ground application of this product after planting and before plant emergence may help decrease infestation of Angular Leaf Spot.

CURRENTS AND GOOSEBERRY: Anthracnose, Leaf Spot: Apply 1-1/3 pints Cuproxat for each 100 gallons of spray mixture at leaf opening and repeat in 10 to 14 day intervals.

DAHLIAS: Botrytis Blight, Leaf Spots: See Ornamental Plants and Trees.

DOGWOOD: Anthracnose, Leaf Spots: See Ornamental Plants and Trees.

EGGPLANT: Anthracnose, Downy Mildew, Early (Alternaria) Blight, Phomopsis Blight, Leaf Spots: Beginning in plant bed or in field before disease appears, apply 2-2/3 pints/acre. Repeat at 7 to 10 day intervals.

GARDENIAS: Leaf Spots: See Ornamental Plants and Trees.

GERANIUMS: Botrytis Blight, Downy Mildew: See Ornamental Plants and Trees.

GLADIOLUS: Botrytis Blight, Downy Mildew, Leaf Spots: See Ornamental Plants and Trees.

GRAPEFRUIT: See Citrus.

GRAPES: Anthracnose: Use 2-2/3 pints Cuproxat for each 100 gallons of spray mixture on dormant vines in late spring. Repeat every 10 to 14 days after leaves form.

Downy Mildew, Black Rot, Powdery Mildew, Bitter Rot: Apply 2-2/3 to 3 pints Cuproxat for each 100 gallons of spray mixture plus 2-6 pounds hydrated lime. Spray when first leaves appear, repeat before blossom open, after petal fall and every two weeks thereafter.

For Downy Mildew control only: Apply 2-2/3 to 3 pints Cuproxat for each 100 gallons of spray mixture for the last two late summer applications following early season application of another fungicide. For exact timing, follow local recommendation for downy mildew infection.

Dead Arm: Use 3 pints Cuproxat plus 16 pounds of hydrated lime for each 100 gallons of spray mixture. Apply as soon as green tips show. **CAUTION:** Slight to severe foliage injury may occur on copper-sensitive varieties such as Concord, Delaware, Niagara and Rosette.)

HOLLYHOCKS: Leaf Spots, Leaf Blights: See Ornamental Plants and Trees.

LETTUCE: Anthracnose, Leaf Spots (*Cercospora*), Downy Mildew: Spray 1-1/3 to 2-2/3 pints/acre before diseases appear. Repeat at 7 to 10 day intervals.

LILAC: Bacterial Blights: See Ornamental Plants and Trees.

LILIES: Botrytis Blight: See Ornamental Plants and Trees.

KUMQUAT, LEMON, LIME: See Citrus.

MANGO (FLORIDA): Anthracnose: Apply monthly after fruit set until harvest at 2-2/3 pints Cuproxat per 100 gallons. Consult Extension Service for recommendations.

MAPLES: Leaf Spots: See Ornamental Plants and Trees.

MARIGOLDS: Leaf Spots: See Ornamental Plants and Trees.

NASTURTIUM: Anthracnose, Leaf Spots: See Ornamental Plants and Trees.

NECTARINES: See Peaches.

OAKS: Leaf Blister, (Taphrina): See Ornamental Plants and Trees.

OLIVES: Peacock Spot: Make first application at 2-2/3 to 4 pints Cuproxat for each 100 gallons of spray mixture, before winter rains fall. A second application in early spring should be made if disease is severe.

ONIONS: Downy Mildew, Leaf Spots: Use 2-2/3 pints/acre. Begin as soon as plants grow to 4 to 6 inches; repeat every 7 days.

ORANGES: See Citrus.

ORNAMENTAL PLANTS AND TREES: Apply 1-1/3 pints Cuproxat for each 100 gallons of spray mixture before diseases appear. Repeat every 7 to 10 days and after each rain.

PALMS: False Smut: See Ornamental Plants and Trees.

PANSIES: Downy Mildew: See Ornamental Plants and Trees.

PEACHES AND NECTARINES: Bacterial Disease (Blight) of Peaches, Leaf Curl, Shot Hole: For dormant application, use 2-2/3 to 5-1/3 pints Cuproxat for each 100 gallons of spray mixture during pink or red bud to popcorn stages. Do not apply during blossom stages or when tree is in leaf. Post-bloom (AL, AR, FL, GA, LA, MS, NC, OK, SC AND TX), apply 1/3 pint Cuproxat for each 100 gallons of spray mixture at first and second cover sprays. Do not use at rates above those recommended. **CAUTION:** Slight defoliation and spotting of leaves may occur from use in cover sprays.

PEANUTS: Leaf Spot (Cercospora): Begin spraying 40-45 days after planting or when first disease symptoms appear. Make ground or aerial application at 2 to 4 pints per acre. For aerial application, use 3-10 gallons of water. Continue applications at 10-14 day intervals. Use sufficient water to get adequate coverage. Cuproxat may be blended with flowable sulfur products.

PEARS: Fire Blight: (West of Mississippi River) Apply 1/3 pint Cuproxat for each 100 gallons of spray mixture or 1-1/3 pints Cuproxat per acre at 5 day intervals throughout bloom period. Excessive dosages may cause fruit russet.

Leaf Blight, Scab: Use 1 pint Cuproxat plus 2-1/2 pounds hydrated lime for each 100 gallons of spray mixture. Spray at the first three cover sprays.

Bull's Eye Rot: Apply 3 pints Cuproxat for each 100 gallons of spray mixture plus a suitable spray oil before harvest. Do not use on Anjou, Comice or Seckel varieties. Limits per acre: 2 pints for Fire Blight, 5-1/3 pints pre-harvest, 20 pints post-harvest.

PEAS: Powdery Mildew: Begin spray treatment before disease symptoms first appear. Use at 2-4 pints per acre according to disease severity. Repeat applications at weekly intervals.

PECANS: Scab: Apply 2 pints Cuproxat for each 100 gallons of spray mixture as foliar spray. Begin applications at end break (2-3 inches long). Make second application at pre-pollination. Thereafter, repeat at 2-4 week intervals as needed to maintain control.

PEONIES: Botrytis Blight, Leaf Spots: See Ornamental Plants and Trees.

PEPPERS: Anthracnose, Bacterial Spot, Early and Late Blight, Cercospora Leaf Spot (Frogeye Spot), Downy Mildew: Spray in plant bed before disease appears. Continue in field at 7 to 10 day intervals, especially during fruiting stage. Use 2-2/3 to 4 pints/acre.

PHLOX: Leaf Spots: See Ornamental Plants and Trees.

PINES: Brown Spot, Needle Blight: See Ornamental Plants and Trees.

PLUMS, PRUNES: Dormant spray for Brown Rot: Apply 5 pints Cuproxat plus 1-1/2 gallons spreader for each 100 gallons of spray mixture in dormant season. LIMITS: 25 pints/acre.

Black Knot: Apply 2 to 3 pints Cuproxat plus 8 to 12 pounds hydrated lime for each 100 gallons of spray mixture at green tip stage. Repeat at shuckshed and in two cover sprays if disease is serious.

Blossom Blight, Leaf Spots: Apply 1-1/2 to 2-1/2 pints Cuproxat plus 6 to 10 pounds hydrated lime for each 100 gallons of spray mixture; spray just before blossoms open; repeat at shuck split and again in 2 to 3 weeks. Over-spraying may injure tender foliage of some Japanese plums.

POTATOES: Early and Late Blight: Apply at 7 day intervals starting when plants emerge until two weeks before harvest. Follow local recommendations for application timing (weather observation). Use 1-1/3 to 2-1/3 pints per acre in those locations where disease is light and up to 4 to 5-1/3 pints per acre where disease is more severe.

RHODODENDRONS: Leaf Blight, Leaf Spots: See Ornamental Plants and Trees.

ROSES: Anthracnose, Black Spot, Downy and Powdery Mildew, Leaf Spots, Stem Canker: See Ornamental Plants and Trees.

STOCK: Leaf Spots: See Ornamental Plants and Trees.

STRAWBERRIES: Downy Mildew, Leaf Blight, Scorch or Spot: After leaves form, spray 2-2/3 to 3 pints Cuproxat in 100 gallons of water per acre at 10 to 14 day intervals.

Anthracnose: Use 2-2/3 to 3 pints per acre plus 2 to 3 pounds hydrated lime/100 gallons pre-blossom. Post-harvest; spray foliage with 3 pints per acre plus 6 pounds hydrated lime per 100 gallons.

SUGAR BEETS: (Cercospora) Leaf Spot: Start spray when disease threatens and continue for 4 to 5 applications. Spray every 7-14 days depending on weather conditions at 2-2/3 to 6-2/3 pints per acre depending on disease severity. Addition of a suitable agricultural spray oil is recommended at 2 quarts per acre.

SWEET PEAS: Downy Mildew, Leaf Blights, Leaf Spots: See Peas.

TANGERINE, TANGELO: See Citrus.

TOMATOES: Early Blight, Late Blight: Start spraying before disease appears. Apply 2-2/3 to 4 pints per acre at 7 to 10 day intervals.

Bacterial Speck: Apply 2-2/3 pints per acre at 10-30 day intervals beginning when the disease threatens. Use more frequent applications when disease pressure is high.

Anthracnose, Bacterial Spot: When disease threatens, apply 2-2/3 to 5-1/3 pints per acre at 7 to 10 day intervals, more frequently when disease is severe. May be combined with maneb or mancozeb. When used in combination with maneb or mancozeb, do not apply within 5 days of harvest.

TULIPS: Botyrtis Blight: See Ornamental Plants and Trees.

VIOLETS: Downy Mildew: See Ornamental Plants and Trees.

VIRGINIA CREEPER: Downy Mildew: See Ornamental Plants and Trees.

WALNUT: Walnut Blight: Apply first spray at early pre-bloom when catkins are partially expanded. Make three additional applications during bloom and early nutlet stages at 7 to 10 day intervals. Additional applications may be necessary when frequent rainfall occurs. Apply 2-2/3 to 4 pints Cuproxat for each 100 gallons of spray mixture.

17 9 17

WARRANTY LIMITATIONS AND DISCLAIMER

Agrolinz, Inc., warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the DIRECTIONS FOR USE when used under normal conditions. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL. Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.) under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses of damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify Agrolinz, Inc., in writing, of any claims to be eligible to receive either remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of Agrolinz, Inc., or any other seller will be one of the following, at the election of Agrolinz, Inc.:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

The seller may not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations and Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent or the seller is not authorized to vary or exceed the terms of this Warranty Limitations and Disclaimer in any manner.