

# Safer™

## INSECTICIDAL SOAP

### CONCENTRATE

KILLS APHIDS, MEALYBUGS, WHITEFLY, MITES AND OTHER PESTS ON TREES, SHRUBS, FLOWERS, VEGETABLES, FRUIT TREES, HOUSEPLANTS AND GREENHOUSES.

Active Ingredients	By Wt.	<b>Keep Out of Reach of Children</b>
Potassium salts of fatty acids	49.0%	
Inert Ingredients	51.0%	<b>CAUTION</b>
TOTAL	100.0%	

Net Contents 32 FL. OZ.

### Safer INSECTICIDAL SOAP

Mix as directed on label with water to use on plants and shrubs. Do not use on new transplants or newly rooted cuttings. Do not apply to plants in full sun.

For control of ADELGIDS, AFHIDS, CRICKETS, EARWIGS, GRASSHOPPERS, LACEBUGS, MEALYBUGS, PLANT BUGS, PSYLLIDS, ROSE SLUG, SAWFLY, SCALES, SPIDER MITES, TENT CATERPILLARS, THRIPS, WHITEFLY and WOOLLY APHIDS on ornamental plants and vegetable gardens and around the home. Mix 2.5 FL. OZ. (5 TBS.) PER GAL. OF WATER. Apply with insecticide pump or sprayer. Be sure to thoroughly wet all surfaces of infested leaves and branches. For aphids and mealybugs spray again after 2 or 3 days. For other pests spray weekly for 2 or 3 weeks or as needed.

**INDOOR PLANTS**—Use as directed above. Be sure that spray penetrates dense foliage to wet all infested surfaces. Do not use on Creole of Thorns, Testaceous scale, and other pests on plants and delicate ferns before making for local application. For ROSE MEALYBUGS on African violet, mix with solution of 1 lbs. per quart of tepid water.

**FLOWERS AND BEEDING PLANTS**—Use as directed above to control aphids, mealybugs, scale and plant bugs. For whitefly, mix 2.5 fl. oz. per gallon of water. Do not use on plants in full bloom. Do not use on gardenias, rhododendrons, sweet peas, and other plants which are sensitive to the active ingredient.

**ORNAMENTAL TREES AND SHRUBS**—Use as directed above. To control aphids, scale and sawfly spray at the peak of leafhopper and whitefly infestation. Do not use on plants which are sensitive to the active ingredient. For tent caterpillars, mix 2.5 fl. oz. per gallon of water. Do not use on plants which are sensitive to the active ingredient. Do not use on plants which are sensitive to the active ingredient.

**FRUIT AND NUT TREES**—Use as directed above on APPLES and PEARS to control aphids, woolly alydids, scale, crawlers, mites, pear psylla and pear slug up to day of harvest; on CITRUS to control aphids, cottony cushion scale, scales and mites up to day of harvest; on ALMONDS, FILBERTS and WALNUTS to control aphids and mites up to day of harvest; on APRICOTS, NECTARINES and PEACHES to control aphids and mites up to day of harvest; on GRAPES to control aphids and leafhoppers up to day of harvest. Avoid spraying when trees are in full bloom.

**VEGETABLES**—Use as directed above on TOMATOES, PEPPERS and EGGPLANT to control aphids, spider mites and whitefly up to day of harvest; on COLE CROPS to control aphids and leafhoppers up to day of harvest; and on CUCUMBERS, SQUASH and MELONS to control aphids, spider mites, squash bug and whitefly up to day of harvest; BEANS and PEAS to control aphids, leafhoppers, grasshoppers and spider mites up to day of harvest.

**GREENHOUSES**—Use as directed above according to crop type. For greenhouse whitefly control, dilute with 2.5 tablespoons of concentrate per gallon of water.

SAFER INC. will not accept liability for damage or injury resulting from misuse.



**Safer, Inc.**

60 WILLIAM STREET  
WELLESLEY, MA 02181

1986, Safer, Inc.

#### PRECAUTIONARY STATEMENTS

Hazards to humans and domestic animals. **CAUTION:** Harmful if swallowed. Avoid contamination of food by this product. May cause irritation, avoid eye contact. First Aid: In case of eye contact, flush with plenty of water. Get medical attention if irritation persists. Environmental Hazards: Keep out of streams, ditches and ponds. Physical or Chemical Hazards: Do not use in warm areas. Do not use in hot sun.

#### STORAGE AND DISPOSAL

Store in cool, dry place. Do not store in areas where children and pets can reach. Do not use in areas where children and pets can reach. Do not use in areas where children and pets can reach. Do not use in areas where children and pets can reach.

EPA REG. NO. 42897-01

**BEST AVAILABLE COPY**

286971

2/3

# Safer™

## INSECTICIDAL SOAP FOR ROSES & FLOWERS

### CONCENTRATE




CONTAINS POTASSIUM SALTS OF FATTY ACIDS  
EFFECTIVE AGAINST APHIDS, EARWIGS AND FLEAS

Active Ingredients	By Wt	<b>Keep Out of Reach of Children CAUTION</b> See side panel for additional precautionary statements
Potassium salts of fatty acids	49.0%	
Inert Ingredients	51.0%	
<b>TOTAL</b>	<b>100.0%</b>	

Net Contents 16 FL. OZ.

SAFER, INC. will not  
accept liability for  
damage or injury  
resulting from misuse



**Safer, Inc.**  
67 WILLIAM STREET  
WELLESLEY, MASSACHUSETTS 02156

© 1986, Safer, Inc.

# Safer™

## INSECTICIDAL SOAP FOR ROSES & FLOWERS

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


Always use fertilizer and solution. Use soft water whenever possible. For roses or new transplants. Do not spray top parts in full sun.

For control of APHIDS, ROSE SLUG, SPIDER MITES, WHITEFLY, LEAFHOPPERS and EARWIGS on roses, flowers and bedding plants: Mix 1 FL. OZ. (5 TABLESPOONS) PER GALLON OF WATER. Apply when first signs of damage are apparent. For APHIDS, ROSE SLUG, WHITEFLY and LEAFHOPPERS spray directly on insects and thoroughly wet foliage. For APHIDS spray again in 2 or 3 days. For other pests spray weekly or as needed. For MITES thoroughly spray underside of leaves where MITES feed and respray weekly for 2 or 3 weeks. For EARWIGS mix 2 tablespoons per gallon of water and spray directly on insects. Do not use on bleeding-heart, crown of thorns, gardenias or sweetpeas. Do not use on azaleas, begonias, camellias, or impatiens without first testing for varietal sensitivity.

### PRECAUTIONARY STATEMENTS

Hazardous to humans and animals.  
**CAUTION:** Harmful to aquatic life. Avoid contact with fish, snails, and shellfish. May cause permanent eye damage. Avoid contact with eyes.  
First Aid: In case of eye contact, flush with plenty of water. Get medical attention if irritation persists.  
Environmental Hazards: This product may be extremely toxic to fish, birds, and mammals.  
**STORAGE AND DISPOSAL:**  
Store in cool place and keep tightly sealed. For suggestions on safe product use, see instructions on label. Recycle when possible. Do not discard in trash. Do not reuse empty containers. Recycle through local recycling program.  
EPA REG. No. 42553-01

EPACORPORATION



4/26/86


3/11

USE IT 'TIL YOU'RE READY TO HARVEST.

# Safer™

## INSECTICIDAL SOAP FOR FRUITS & VEGETABLES

### CONCENTRATE




Active Ingredients: Potassium salts of fatty acids 49.0%  
Inert Ingredients: 51.0%  
TOTAL: 100%

**Keep Out of Reach of Children**  
**CAUTION**  
See side panel for additional precautionary statements.

Net Contents 16 FL. OZ.

SAFER, INC. will not accept liability for damage or injury resulting from misuse.



Safer, Inc.

6 W. 48th Street  
New York, N.Y. 10019

© 1986 Safer, Inc.

### Safer™

#### INSECTICIDAL SOAP FOR FRUITS & VEGETABLES

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

Always use freshly mixed solution. Use soft water whenever possible. Do not use on new transplants or when plants are stressed by drought. Unless stated otherwise MIX 2.5 FL. OZ. (5 TBS.) PER GAL. OF WATER. Do not apply to plants in full sun.

**FRUIT TREES:** Spray when insects first appear or to prevent damage. Apply to achieve thorough coverage of infested foliage and/or branches. Repeat at weekly intervals as needed. May be used up to day of harvest. Avoid spraying when transplants are in full bloom. Use as directed on **APPLES** and **PEARS** to control aphids, woolly aches, fruit scale, crawler mites, pear psylla and pear thrips. Use as directed on **CHERRIES** to control aphids, cherry curculionids, leaf-miner, cherry thrips and borers. Use as directed on **APRICOTS**, **NECTARINES** and **PEACHES** to control aphids and borers. Use as directed on **GRAPEVINES** to control aphids and leafhoppers.

**VEGETABLES:** Apply when insects first appear or when damage appears. Spray to wet infested plant surfaces especially the underside of leaves for **lettuce whittly** and **leafhoppers**. For **aphids** spray again in 2 days. For other pests apply at weekly intervals as necessary. May be used up to day of harvest. One gallon of spray covers 250 sq. ft. Use as directed on **BEANS** and **PEAS** to control **aphids**, **leafhoppers** and **harlequin bug**; on **CUCURBITS** to control **aphids**, **leafhoppers**, **spider mites**, **whittly** and **squash bug**; on **TOMATOES**, **PEPPERS** and **EGGPLANT** to control **aphids**, **spider mites** and **whittly**.

**PRECAUTIONARY STATEMENTS**

Hazardous to humans by ingestion.  
**CAUTION:** Harmful if swallowed. Avoid contact with eyes and avoid skin. May cause eye irritation. For eye contact, flush with plenty of water. Get medical attention if irritation persists.  
**First Aid:** In case of eye contact, flush with plenty of water. Get medical attention if irritation persists.  
**Environmental Hazards:** None known.  
**Physical or Chemical Hazards:** None known.

**STORAGE AND DISPOSAL**

Store in cool place and away from heat. Do not store in plastic. Do not use product, securely wrap up in original container, and dispose of in accordance with local, state and federal regulations. Do not discharge into water.  
EPA REG. No. 42697-C

ACCEPTED

MAR 6 1987

**BEST AVAILABLE COPY**

42697-1