

Hazards to Humans and Domestic Animals

CAUTION Harmful if swallowed
Environmental Hazards Keep out of lakes, streams or ponds
DISPOSAL Do not reuse empty container. Rinse clean with water and dispose of in trash collection.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


Directions continued on right side panel

EPA Reg. No. 42697
EPA Est. No. 907 TN-01



Safer Agro-Chem Inc.
370 Brooks Rd. East
Memphis, TN 38116

Safer Agro-Chem's Insecticidal Soap FOR AFRICAN VIOLETS



READY TO USE
Controls Aphids,
Mealybugs, Whitefly,
Thrips and Spider Mites.

Active ingredients	By Wt
Potassium salts of fatty acids	15%
Inert ingredients	98.5%
Total	100.0%

Keep out of reach of children

CAUTION

See side panel for additional cautions

Net Contents
8 fl. oz.

To Control APHIDS, MITES, WHITEFLY and THRIPS on the foliage, spray upper and lower leaf surfaces thoroughly or dip the entire basal portion of the plant in spray solution while holding the soil in place to avoid spillage. Repeat treatment in 7 to 10 days.

Note: Store and use spray at room temperature. Protect blossoms, or rinse with water after treatment to avoid spotting. Do not allow excess spray solution or water to sit in center of plant. Keep plants out of direct sun while foliage is wet.

To Control ROOT MEALYBUGS, be sure soil is moist, then pour ready to use spray solution evenly over the soil surface until drainage begins. Leave excess solution in watering saucer to be re-absorbed by the soil. Repeat in 7 to 10 days.

Safer Agro-Chem Inc. nor the seller accept liability resulting from misuse.

1984 SAFER AGRO-CHEM INC

