

### Safer™ Fungicide-Miticide

**SAFER CARES:** Safer is an effective, yet different, type of fungicide-miticide. Safer's scientists have formulated an easy to use product for controlling diseases and mites. If you care about your houseplants, garden, family and pets, use Safer products with confidence.

**PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS**

**CAUTION:** Harmful if swallowed.

**ENVIRONMENTAL HAZARDS:** Do not apply directly to water or contaminate water when cleaning equipment or disposing of waste.

**DISPOSAL:** Do not reuse container or contaminate water or food by storage. Dispose of in trash collection.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Buyer assumes all responsibility for safety and use not in accordance with directions. Avoid contact with plants until spray has dried.

Do not use on Boston fern, spinach, apricots, filberts, walnuts and viburnum. Do not apply when temperature exceeds 85°F. Do not apply for at least 4 weeks following application of an oil spray or within 4 days of other pesticide application. Spray plants thoroughly to wet upper and lower surfaces of foliage. Apply at 7 days intervals when mites are present. Store in cool place.

# Safer™ Garden Fungicide-Miticide



**READY TO USE**

**CONTROLS POWDERY MILDEW, BLACK SPOT, SCAB BROWN  
CANKER, LEAF SPOT AND RUST  
PLUS MITES ON FRUITS, VEGETABLES, FLOWERS & FOLIAGE**

Active Ingredients	By Wt.	Keep out of reach of children
Sulfur	0.4%	<b>CAUTION</b>
Inert Ingredients	99.6%	See back panel for additional cautions.

Net Contents 32 fl. oz.

**FLOWERS, FOLIAGE PLANTS AND ORNAMENTAL:** Spray at 7 to 10 day intervals or as needed to control powdery mildew, leaf spot, black spot, rust and mites.

**ROSES:** Spray at 7 to 10 day intervals to control powdery mildew, leaf spot, rust and mites. Be sure to treat new growth and buds. Avoid spraying when plants are in full sun or temperature exceeds 85°F.

**FRUITS AND VEGETABLES:** May be used up to day before harvest.

**BEANS AND PEAS:** to control powdery mildew and mites. Spray at first sign of disease or mite infestation, and repeat at seven day intervals as necessary.

**TOMATOES** to control russet mite spray when fruit begins to set and repeat as necessary.

**STRAWBERRIES** to control powdery mildew spray at first sign of infection and repeat at three week intervals. Do not apply to strawberry fruit that will be used for canning.

**ACCEPTED**

**MAY 1 9 1986**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 42617-17



EPA Reg. No. 42617-17  
EPA Est. No. 9377H-01  
Safer, Inc. 60 William St.  
Woburn, MA 02151

