

# Safer

FOR OUTDOORS

## GARDEN FUNGICIDE

READY TO USE

CONTROLS POWDERY MILDEW, BLACK SPOT, SCAB, BROWN CANKER, LEAF SPOT & RUST, ON FRUITS, VEGETABLES, FLOWERS & FOLIAGE.

ACTIVE INGREDIENTS	BY WT.
SULFUR	0.4%
INERT INGREDIENTS	99.6%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET CONTENTS 24 FL. OZ.

6945-1118

### PRECAUTIONARY STATEMENTS

Hazards to humans:

**CAUTION:** Harmful if swallowed.

Environmental Hazards: Do not apply directly to water or contaminate water when cleaning equipment or disposing of waste.

### STORAGE & DISPOSAL

Store at room temperature & protect from freezing. Do not reuse container or contaminate water or food by storage. Dispose of in trash collection.

**USE CAUTIONS:** Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not use on Boston Fern, spinach, apricots, figs, walnuts and viburnum. Do not apply when temperature exceeds 85°F. or in full sun. Do not apply for at least 4 weeks following application of oil spray or within 4 days of other pesticide application.

**RE-ENTRY:** Avoid contact with plants until spray has dried.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Spray plants thoroughly to wet the upper and lower surfaces of foliage. Repeat as necessary for mite/leaf control.

**FLOWERS, FOLIAGE PLANTS AND ORNAMENTALS:** Spray at 7 to 10 day intervals or as needed to control powdery mildew, leaf spot, black spot, and rust.

**ROSES:** Spray at 7 to 10 day intervals to control powdery mildew, black spot, and rust. Be sure to treat new growth and buds.

**FRUITS AND VEGETABLES:** May be used up to day before harvest.

**BEANS AND PEAS:** to control powdery mildew.

Spray at first sign of disease infestation, and repeat at 7 day intervals as necessary.

**STRAWBERRIES:** To control powdery mildew spray at first sign of infection and repeat at 3 week intervals. Do not apply to strawberry fruit that will be used for canning.

SAFER INC., will not accept liability for damage or injury resulting from misuse.

EPA REG. No. 42897-17

EPA Est. No. 327-TX-01, 489-TX-03, 429-CA-001  
Superscript corresponds to first letter of lot no. on bottom of container.

**Safer, Inc.**

180 WILLS AVENUE  
NEWTON, MA 02459

Product Code 2045



©1989, Safer, Inc.

Safer® is a registered trademark of Safer, Inc.

8045-01188

DATE JUN 2 1990  
NOTICE LABEL NO PER PR NOT

6/18/90

Full packet

4/26/17-17  
Page 12  
P421

SHAKE  
WELL




READY TO USE

CONTROLS POWDERY MILDEW, BLACK SPOT, SCAB, BROWN CANKER,  
LEAF SPOT & RUST, ON FRUITS, VEGETABLES, FLOWERS & FOLIAGE.

Active Ingredients	By Wt.	<b>Keep Out of Reach of Children</b>
Sulfur	0.4%	
Inert Ingredients	99.6%	<b>CAUTION</b>
TOTAL	100.0%	

See side panel for additional  
precautionary statements.

Net Contents 64 FL. OZ.



**Safer, Inc.**

60 WILLIAM STREET  
WELLESLEY, MA 02181

© 1987, Safer, Inc.  
Safer™ is a trademark of Safer, Inc.  
6046F-107

**Safer™**

GARDEN FUNGICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Spray plants thoroughly to wet the upper and lower surfaces of foliage. Repeat as necessary for mildew control.

**DIRECTIONS FOR USE**

**FLOWERS, FOLIAGE PLANTS AND ORNAMENTALS:** Spray at 7 to 10 day intervals or as needed to control *powdery mildew*, *leaf spot*, *black spot*, and *rust*.

**ROSES:** Spray at 7 to 10 day intervals to control *powdery mildew*, *black spot*, and *rust*. Be sure to treat new growth and buds.

**FRUITS AND VEGETABLES:** May be used up to day before harvest.

**BEANS AND PEAS:** to control *powdery mildew*. Spray at first sign of disease infestation, and repeat at 7 day intervals as necessary.

**STRAWBERRIES:** To control *powdery mildew* spray at first sign of infection and repeat at 3 week intervals. Do not apply to strawberry fruit that will be used for canning.

PRODUCT NO. 2046

6046B-107

**PRECAUTIONARY STATEMENTS**

Hazards to humans: **CAUTION:** Harmful if swallowed.  
Environmental hazards: Do not apply directly to water or contaminate water when cleaning equipment or disposing of waste.

**STORAGE & DISPOSAL**

Store at room temperature & protect from freezing. Do not reuse container; it contaminate water or food by storage. Dispose of in trash collection.

**USE CAUTIONS:** Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not use on Boston Fern, spinach, apricots, filberts, walnuts and viburnum. Do not apply when temperature exceeds 85°F. or in full sun. Do not apply for at least 4 weeks following application of oil spray or within 4 days of other pesticide application.

**RE-ENTRY:** Avoid contact with plants until spray has dried.

EPA REG. No. 42697-17 EPA EST. NO. 907-TN-01 © 1987  
Superscript corresponds to first letter of lot number on bottom of container

# Safer

FOR OUTDOORS

## GARDEN FUNGICIDE

READY TO USE

CONTROLS POWDERY MILDEW, BLACK SPOT, SCAB, BROWN CANKER, LEAF SPOT & RUST, ON FRUITS, VEGETABLES, FLOWERS & FOLIAGE.

ACTIVE INGREDIENTS 0.04%  
INERT INGREDIENTS 99.96%  
TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET CONTENTS 24 FL. OZ.

6045-1108

DATE LABEL NOT VALID  
JUN 21 1990

**PRECAUTIONARY STATEMENTS**

Hazards to humans  
**CAUTION: Harmful if swallowed.**  
Environmental hazards: Do not allow to contaminate water when cleaning equipment.

**STORAGE** (in 19)

Store at room temperature & pressure. Do not reuse container or contents. Dispose of in trash.

**USE CAUTIONS:** Some plants are susceptible under certain climatic conditions. Do not spray apricots, filberts, walnuts when temperature exceeds 85°F for at least 4 weeks following application. 4 days of other pesticides.

**RE-ENTRY:** Avoid contact with plants.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Repeat as necessary.

**FLOWERS, FOLIAGE PLANTS AND TREES:** Spray at 7 to 10 day intervals or as needed.

**ROSES:** Spray at 7 to 10 day intervals to control black spot, and rust. Be sure to spray foliage.

**FRUITS AND VEGETABLES:** Spray at 7 to 10 day intervals to control leaf spot, black spot, and rust. Be sure to spray foliage before harvest.

**BEANS AND PEAS:** to control black spot, and rust. Spray at first sign of disease at 7 day intervals as needed.

**STRAWBERRIES:** To control powdery mildew, spray at 7 to 10 day intervals and repeat at 3 week intervals. Fruit that will be used for processing should be sprayed.

SAFER INC. will not accept liability for damage resulting from use of this product.

EPA REG No 42697-17

EPA Est No 907 TN-01 409 TN 03 4091 CA-001  
Superscript corresponds to first letter of label on bottom of container.

**Safer, Inc.**

189 WELLS AVENUE  
NEWTON, MA 02459

Product Code 2045

1989, Safer is a registered trademark