

PM 21

42697-17

9-16-97

10/5

EPA #42697-17  
Page 1 of 5

**SAFER® BRAND  
GARDEN FUNGICIDE READY-TO-USE**

**Controls  
POWDERY MILDEW, BLACK SPOT, LEAF SPOT and RUST  
on  
Fruits, Vegetables, Flowers and Foliage**

<b>ACTIVE INGREDIENTS</b>	<b>BY WT.</b>
Sulfur.....	0.40%
<b>INERT INGREDIENTS</b> .....	<b>99.60%</b>
<b>TOTAL</b> .....	<b>100.00%</b>

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel for Precautionary Statements  
and First Aid (Statement of Practical Treatment).

**NET CONTENTS 32 FL. OZ.**

ACCEPTED

9/16/97

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 42697-17

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals:**

**CAUTION:** Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**First Aid (Statement of Practical Treatment):**

**IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

**In Case of Emergency, call 1-800-228-5635, ext. 170**

**Environmental Hazards:** Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**WHAT IT IS:** Safer® Brand Garden Fungicide is specially formulated with sulfur which inhibits the attack of fungal diseases by creating an environment which is not optimal for disease growth.

**WHAT IT CONTROLS:** Powdery mildew, black spot, leaf spot and rust

**WHERE TO USE IT:** Roses, Flowers, Foliage Plants, Ornamentals, Beans, Peas, Broccoli, Cabbage, Cauliflower, Collards, Kale, Peppers, Strawberries, Apples, Pears, Citrus, Potatoes, Cucumbers, Squash, Grapes, Blackberries, Blueberries, Raspberries, Loganberries, Currants, Melons and Tomatoes

**HOW TO USE IT:** Spray at first sign of infection. Spray plants thoroughly to wet the upper and lower surfaces of foliage.

**ROSES, FLOWERS, FOLIAGE PLANTS AND ORNAMENTALS (INCLUDING ALDER, ASTER, AZALEA, BUTTONBUSH, CARNATION, CATALPA, CHRYSANTHEMUM, DAHLIA, DELPHINIUM, HONEYSUCKLE, HORSE CHESTNUT, HYDRANGEA, LILAC, LINDEN, MATRIMONY VINE, PHLOX, RHODODENDRON, SNAPDRAGON, STAGHORN SUMAC, SWEETPEA, SYCAMORE, TRUMPET VINE, WITCH HAZEL, AND ZINNIA):** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew, leaf spot, black spot, and rust. Be sure to treat new growth and buds.

**BEANS, PEAS, BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE AND PEPPERS:** Apply at first sign of disease infestation. Spray at 7 day intervals, following rain, or as

needed to control powdery mildew. Can be used up to the day before harvest.

**STRAWBERRIES:** Apply at first sign of disease infestation. Repeat at 3 week intervals to control powdery mildew. Do not use on strawberry fruit that will be used for canning. Can be used up to the day before harvest.

**APPLES & PEARS:** Apply at pre-bloom stage or after petal fall as needed to control powdery mildew. **NOTE:** Some apple & pear varieties are sulfur sensitive; test for varietal sensitivity prior to full scale treatment.

**CITRUS:** Do not apply within 4 weeks following use of an oil spray.

**POTATOES:** Apply before disease infestation appears and repeat as necessary to control powdery mildew.

**CUCUMBERS & SQUASH:** Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants.

**GRAPES:** For control of powdery mildew, apply when new shoots are 6-10 inches. Repeat before blooms open and at 14 day intervals as needed. **NOTE:** Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord & other labrusca varieties.

**BLACKBERRIES, BLUEBERRIES, RASPBERRIES, LOGANBERRIES AND CURRANTS:** Apply each week starting at first bloom color to control powdery mildew. Continue to spray at 10 to 14 day intervals until fruit is set, then repeat as needed or following rain.

**MELONS:** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals following rain, or as needed to control powdery mildew. **Note:** Melon foliage is extremely sensitive to sulfur burn. See paragraph on Use Cautions.

**TOMATOES:** Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants.

**USE CAUTIONS:** Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not use on Boston fern, spinach, apricots, filberts, walnuts and viburnum. Do not apply when plants are in full sun or when temperature exceeds 85°F. Do not apply for at least 4 weeks following application of oil spray. Avoid contact with plants until spray has dried.

**STORAGE AND DISPOSAL**

**Storage:** Store at room temperature, in original container only, out of reach of children and animals. Keep from freezing.

**Disposal:** Securely wrap original container in several layers of newspaper and discard in trash.

Ringer Corporation or Safer®, Inc., will not accept liability for injury or damage resulting from misuse.

If you are not completely satisfied with this product, or for consumer information, call 1-612-703-3300 weekdays 9-5 Central time to arrange for a refund of the purchase price or replacement of the product. Proof of purchase is required.

EPA #42697-17

Page 4 of 5

Distributed by:

Safer<sup>®</sup>, Inc., a wholly owned subsidiary of Ringer Corporation

9555 James Avenue South, Suite 200

Blotmington, MN 55431

(612) 703-3300

EPA Reg. No. 42697-17

H

K

EPA Est. No. 44616-MO-001

11600-MN-001

Circled or superscript letter corresponds to first letter of lot number on bottom or back of container.

© Copyright Ringer Corporation, 1997

4075

612-887-1383 RINGER CORP. 5045  
EPA #42697-17

Page 5 of 5

**Optional Marketing claims:**

**Home ■ Lawn ■ Garden Solutions™ for Pest Control since 1976**

**Effective**

**Proven effective through University testing**

**Only Ready-to-Use formula on the market**

**Water-based**

**Made with elemental sulfur**

**University Tested**

**Use as a preventative treatment.**

**Satisfaction guaranteed. See back panel.**

**Use up to day before harvest**

**Always shake well before using**

**Si no usted entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)**

**The Right Solution™. Safer® Garden Fungicide is used for prevention and control of fungal diseases, but is gentle on flowers, plants and produce. It treats foliage with the naturally occurring essential element sulfur, to create an environment where damaging fungi cannot live.**

**Its non-burning, easy to use spray formulation can be used on flowers, fruits and vegetables up until the day before harvest.**