

42697-17

08-07-2007

D 318106
5814613

1/6



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:
42697-17

Date of Issuance:
AUG 7 2007

Terms of Issuance:
Unconditional

Name of Pesticide Product:
Safer Garden Fungicide Ready-to-Use

NOTICE OF PESTICIDE:
 Registration
XX Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Safer, Inc.
69 North Locust Street
Lititz, PA 17543

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document(s), EPA has reregistered the product listed above using the label accepted by the Agency on September 17, 2002. This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Signature of Approving Official:

Mary Waller, Product Manager (21)
Registration Division, Fungicide Branch

Date:

8/7/2007

EPA Reg. No. 042697-00017
Draft Label Amendment
June 4, 2002
Page 1 of 5
MASTER LABEL- FRONT PANEL (PAGE #1 of 1)

2/6

ACCEPTED
SEP 17 2002
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 42697-17

SAFER® BRAND GARDEN FUNGICIDE READY-TO-USE

Controls
POWDERY MILDEW, BLACK SPOT, LEAF SPOT and RUST
On
Fruits, Vegetables, Flowers and Foliage

[Additional Brand Names]


Safer® Brand Garden Fungicide For Flowers, Fruit, and Vegetables Ready-To-Use

ACTIVE INGREDIENT:	BY WT.
Sulfur.....	0.40%
OTHER INGREDIENTS:.....	99.60%
TOTAL.....	100.00%

KEEP OUT OF REACH OF CHILDREN

See back panel for additional Precautionary Statements

NET CONTENTS 32 FL. OZ.



3/6

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
If in eyes:	
-	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
-	Call a poison control center or doctor for treatment advice.
If on skin or clothing:	
-	Take off contaminated clothing.
-	Rinse skin immediately with plenty of water for 15-20 minutes.
-	Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment.	

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of rinsate.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WHAT IT IS: Safer® Brand Garden Fungicide Ready-To-Use is specially formulated with sulfur which inhibits the attack of fungal diseases by creating an environment which is not optimal for disease growth.

WHAT IT CONTROLS: Powdery mildew, black spot, leaf spot and rust.

WHERE TO USE IT: Roses, Flowers, Foliage Plants, Ornamentals, Beans, Peas, Broccoli, Cabbage, Cauliflower, Collards, Kale, Peppers, Strawberries, Apples, Pears, Potatoes, Cucumbers, Squash, Grapes, Blackberries, Blueberries, Raspberries, Loganberries, Currants, Melons and Tomatoes.

HOW TO USE IT: Spray at first sign of infection. Spray plants thoroughly to wet the upper and lower surfaces of foliage.



4/6

ROSES, FLOWERS, FOLIAGE PLANTS & ORNAMENTALS: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew, leaf spot, black spot, and rust. Be sure to treat new growth and buds.

AZALEA, BUTTONBUSH, CHRYSANTHEMUM, HYDRANGEA, MATRIMONY VINE, RHODODENDRON, AND SNAPDRAGON: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew, leaf spot, and rust. Be sure to treat new growth and buds.

CATALPA, DELPHINIUM, HONEYSUCKLE, HORSE CHESTNUT, LILAC, LINDEN, PHLOX, SWEETPEA, SYCAMORE, TRUMPET VINE, WITCH HAZEL, AND ZINNIA: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew and leaf spot. Be sure to treat new growth and buds.

CARNATION AND STAGHORN SUMAC: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control leaf spot and rust. Be sure to treat new growth and buds.

ALDER AND ASTER: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew and rust. Be sure to treat new growth and buds.

DAHLIA: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew. Be sure to treat new growth and buds.

BEANS, PEAS, BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE AND PEPPERS: Apply at first sign of disease infestation. Spray at 7 day intervals, following rain, or as needed to control powdery mildew. Can be used up to the day before harvest.

STRAWBERRIES: Apply at first sign of disease infestation. Repeat at 3 week intervals to control powdery mildew. Do not use on strawberry fruit that will be used for canning. Can be used up to the day before harvest.

APPLES & PEARS: Apply at pre-bloom stage or after petal fall as needed to control powdery mildew. NOTE: Some apple & pear varieties are sulfur sensitive; test for varietal sensitivity prior to full scale treatment.

POTATOES: Apply before disease infestation appears and repeat as necessary to control powdery mildew.

CUCUMBERS & SQUASH: Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants.

GRAPES: For control of powdery mildew, apply when new shoots are 6-10 inches. Repeat before blooms open and at 14 day intervals as needed. NOTE: Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord & other labrusca varieties.

BLACKBERRIES, BLUEBERRIES, RASPBERRIES, LOGANBERRIES AND CURRANTS: Apply each week starting at first bloom color to control powdery mildew.

5/6

Continue to spray at 10 to 14 day intervals until fruit is set, then repeat as needed or following rain.

MELONS: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals following rain, or as needed to control powdery mildew. NOTE: Melon foliage is extremely sensitive to sulfur burn. See paragraph on Use Cautions.

TOMATOES: Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants.

USE CAUTIONS: Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not use on Boston fern, spinach, apricots, filberts, walnuts and viburnum. Do not apply when temperature exceeds 85° F or in full sun. Do not apply for at least 4 weeks following application of oil spray. Avoid contact with plants until spray has dried.

STORAGE AND DISPOSAL

STORAGE: Store only in original container in a cool, dry area inaccessible to children and pets. Protect from freezing and heat. Keep container tightly closed to prevent evaporation.

DISPOSAL:

If Empty: Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled: Call your local solid waste agency or (800)-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Woodstream Corp. and its subsidiaries will not accept liability for damages or injury resulting from misuse. For information on this product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Distributed by

Safer®, Inc., a wholly owned subsidiary of Woodstream Corporation

69 North Locust Street

Lititz PA 17543

1-800-800-1812 for information

EPA Reg. No. 42697-17


H

D

EPA Est. No. 44616-MO-01

63456-MN-01

Circled or superscript letter corresponds to first letter of lot number on label, or on container side or bottom.



All ® Property of Woodstream Corporation

© Copyright Woodstream Corporation, 2002

6/6

[Optional Marketing Claims]

Effective
Proven effective through University testing
Water-based
Made with elemental sulfur
University Tested
Use as a preventative treatment.
Use up to day before harvest.
Always shake well before using.
Shake well
Made in USA
Printed in USA

Si no usted entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

The Right Solution™. Safer® Brand Garden Fungicide Ready-To-Use is used for prevention and control of fungal diseases, but is gentle on flowers, plants and produce. It treats foliage with the naturally occurring essential element sulfur, to create an environment where damaging fungi cannot live.

Its non-burning, easy to use spray formulation can be used on flowers, fruits and vegetables up until the day before harvest.

