

1 of 3
pma
42697-18

LABEL FRONT PANEL

ACCEPTED
SEP 19 1988
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 42697-18

SAFER™

GARDEN FUNGICIDE - MITICIDE
CONCENTRATE

FOR THE CONTROL OF
POWDERY MILDEW, BLACK SPOT, SCAB,
BROWN CANKER, LEAF SPOT AND RUST

PLUS
MITES, CHIGGERS AND THRIPS
ON


FRUITS, VEGETABLES, FLOWERS
FOLIAGE PLANTS, AND ORNAMENTALS
IN AND AROUND HOMES, NURSERIES,
AND GREENHOUSES

ACTIVE INGREDIENTS:	BY WEIGHT:
Sulfur	12.0%
INERT INGREDIENTS	88.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Side Panel for Additional Precautionary Statements

EPA Reg. No. 42697-18
EPA Est. No. 907-TN-01
Net Contents xx Fl. Oz.



DRAFT LABEL - BACK PANEL**PRECAUTIONARY STATEMENTS:**

Hazards to Humans and Domestic Animals:

CAUTION: Harmful if swallowed. Do not inhale spray mist. Avoid contact with eyes. Do not use for medical purposes or as a feed supplement.

Environmental Hazards: Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of waste.

USE CAUTIONS: Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not use on Boston Fern, spinach, apricots, filberts, walnuts, and viburnum. Do not apply when temperature exceeds 85° F. Do not apply for at least 4 weeks following application of an oil spray or within 4 days of other pesticides. Be sure to wash and rinse spray equipment thoroughly after each use. Do not allow mixed spray to stand in spray equipment.

Re-entry: Avoid contact with plants until spray has dried.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Do not re-use empty container. Wrap the empty container and put in trash for collection. Store in a cool place.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For control of blackspot, scab, brown canker, leaf spot, rust, mites and thrips. Mix one part concentrate to 30 parts water or 4 fl. oz. (8 tablespoons) per gallon of water. For powdery mildew mix 1 part concentrate to 60 parts water or 2 fl. oz. (4 tablespoons) per gallon of water. Place about half of the volume of water in sprayer, add concentrate, stir well, add rest of water and mix thoroughly. Run agitator on large tank sprayers. For coverage see individual applications.

Spray plants thoroughly to wet the upper and lower surfaces of foliage. Repeat as recommended. For control of mites, apply at seven day intervals.

FLOWERS, FOLIAGE PLANTS, AND ORNAMENTALS: To control powdery mildew, leaf spot, black spot, rust, and mites. Spray at 7 to 10 day intervals or as needed.

ROSES: To control powdery mildew, black spot, brown canker, rust, mites, aphids, and thrips. Spray at 7 to 10 day intervals or following rain. Be sure to treat new growth and buds. Avoid spraying when plants are in full sun or temperature exceeds 85° F. One gallon of spray covers approximately 350 sq.ft. (125 gal. per acre).

BEANS AND PEAS: To control powdery mildew and mites. Spray at first sign of disease or mite infestation and repeat at 7 day intervals as necessary. May be used up to day before harvest. Apply 1 gallon of spray to each 250 sq. ft. (175 gal. per acre).

DRAFT LABEL BACK PANEL (CONT.)

POTATOES: To control powdery mildew. Spray before disease appears and repeat as necessary. May be used to day before harvest. Apply one gallon of spray to each 350 sq. ft. (125 gal. per acre).

TOMATOES: To control russet mite. Spray when fruit begins to set and repeat as necessary. May be used up to day before harvest. Apply 1 gallon of spray to each 350 sq. ft. (125 gal. per acre).

CUCUMBERS AND SQUASH: To control powdery mildew. Spray when disease appears and repeat as necessary. Do not spray young plants. May be used up to day before harvest. Apply 1 gallon of spray to each 600 sq. ft. (75 gal. per acre).

STRAWBERRIES: To control powdery mildew. Spray when disease appears and repeat as necessary. Do not spray young plants. May be used up to day before harvest. Apply 1 gallon of spray to each 300 sq. ft. (150 gal. per acre). CAUTION: Do not apply to strawberry fruit that will be used for canning.

BLACKBERRIES, BRAMBLES, AND BLUEBERRIES: To control powdery mildew apply at the rate of 1 gal. to approximately 300 sq. ft. (150 gal. per acre) when new shoots are about 6 inches long or when disease is expected. Repeat at 10 day intervals.

APPLES AND PEARS: To control powdery mildew, scab, and mites apply to thoroughly wet foliage or at rate of 300 gal. per acre. Apply to pre-bloom stage or after petal fall as needed. CAUTION: Some apple and pear varieties are sulfur sensitive, test for varietal sensitivity prior to full scale treatment or consult local extension specialist.

CHERRIES AND PLUMS: To control powdery mildew and leaf spot apply to thoroughly wet foliage or at 300 gal. per acre rate. Apply from bloom on as needed.

PEACHES AND NECTARINES: To control powdery mildew, brown rot blossom blight, scab, and mites apply to thoroughly wet foliage or at 300 gal. per acre rate from pink stage on as needed. CAUTION: Do not use on apricots.

CITRUS (ALL VARIETIES): To control rust mite apply to thoroughly wet infested foliage or at rate of 125 - 150 gal. per acre as necessary. Do not apply within 4 weeks following use of an oil spray.

GRAPES: To control powdery mildew apply to thoroughly cover foliage or at a rate of 100 gal. per acre. Begin when new shoots are 6 to 10 inches; repeat before blooms open and at 14 day intervals as needed. CAUTION: Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord and other labrusca varieties.

GREENHOUSES: To control powdery mildew, leaf spot, rust, and mites on bedding plants, propagates, fruits, and vegetables. Mix 1 part concentrate to 30 parts water (8 tablespoons per gallon) and apply according to above recommendations by crop type. Do not use when greenhouse temperature is above 85° F. Test for phytotoxicity before full-scale application.

CHIGGERS: To control chiggers spray infested vegetation as necessary. Keep brush and grass trimmed to discourage re-infestation.

Safer Inc. will not accept liability for damage resulting from misuse or uses beyond those described on this label.