

Safer

FOR OUTDOORS

GARDEN FUNGICIDE

CONCENTRATE

CONTROLS POWDERY MILDEW, BLACK SPOT, SCAB, BROWN CANKER, LEAF SPOTS & RUST, ON FRUITS, VEGETABLES, FLOWERS & ORNAMENTALS.

ACTIVE INGREDIENTS BY WT
SULFUR 12.0% **KEEP OUT OF REACH OF CHILDREN**
INERT INGREDIENTS 88.0%
TOTAL 100.0%

CAUTION
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS 16 FL. OZ.

SAFER, INC. WILL NOT
ACCEPT LIABILITY FOR
DAMAGE OR INJURY
RESULTING FROM MISUSE.



DISTRIBUTED BY:
Safer, Inc.

189 WELLS, AVENUE
NEWTON, MA 02159
©1989, SAFER, INC.

SAFER® IS A REGISTERED
TRADEMARK OF
SAFER, INC.

69141409

USE CAUTIONS: Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not apply on Boston Fern, spinach, apricots, fibrous, walnuts and viburnum. Do not apply when temperature exceeds 85°F. Do not apply for at least 4 weeks following application of an oil spray or within 4 days of other pesticides. Be sure to wash & rinse spray equipment thoroughly after each use. Do not allow mixed spray to stand in spray equipment. Re-entry: Avoid contact with plants until spray has dried.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
For control of BLACKSPOT, SCAB, BROWN CANKER, LEAF SPOT and RUST, mix 1 part concentrate to 30 parts water or 4 fl. oz. (8 lbs.) per gal. For POWDERY MILDEW mix 2 fl. oz. (4 lbs.) per gal of water. Place about half of the volume of water in sprayer, add concentrate, stir well, add rest of water and mix thoroughly. Run agitator on large tank sprayers. For coverage see individual applications. Spray plants thoroughly to wet the upper and lower surfaces of foliage. Repeat as recommended.

To control powdery mildew, leaf spot, black spot and rust on FLOWERS, FOLIAGE PLANTS & ORNAMENTALS, spray at 7 to 10 day intervals or as needed. To control powdery mildew, black spot, brown canker and rust on ROSES, spray at 7 to 10 day intervals or following rain. Be sure to treat new growth & buds. Avoid spraying when plants are in full sun or temperature exceeds 85°F. Apply 1 gal. of mix to each 350 sq. ft.

To control powdery mildew on BEANS & PEAS, spray at first sign of disease, re-treatment & repeat at 7 day intervals as necessary. May be used up to day before harvest. Apply 1 gal. of mix to each 250 sq. ft. To control powdery mildew & scab on APPLES & PEARs, apply to thoroughly wet foliage. Apply at pre-bloom stage or after petal fall as needed. Some apple & pear varieties are sulfur sensitive, test for petal sensitivity prior to full scale treatment. Do not apply within 4 weeks following use of an oil spray.

To control powdery mildew on POTATOES, spray before disease appears & repeat as necessary. Apply 1 gal. of mix to each 350 sq. ft. On CLIMBERS & SQUASH, spray when disease appears & repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 600 sq. ft. On STRAWBERRIES, spray when disease appears & repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 300 sq. ft. Do not apply to strawberries that will be used for canning. Do not spray for day before harvest. On GRAPEs, tho' bushy cover foliage. Begin when new shoots are 6-10 inches long. Repeat before blooms open & at 14 day intervals as needed. Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord & other labrusca varieties.

GREENHOUSES: To control powdery mildew, leaf spot & rust on bedding plants, propagates, fruits & vegetables. Mix 1 part concentrate to 30 parts water (8 lbs. per gal.) & apply according to above recommendations by crop type. Test for phytotoxicity before full-scale application.

Product Code 2044

6044-B1189

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals
CAUTION: Harmful if swallowed. Do not inhale spray mist. Avoid contact with eyes. Do not use for medical purposes or as a feed supplement. Environmental Hazard: Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of waste.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feeding storage or disposal containers. Store in a cool place. Wrap the empty container and put in trash for collection. EPA REG NO 4269-18
⑧ EPA EST NO 4091 TN-03 61473 CN 002
MADE IN CANADA



LATE NOTICE 88-8
JUN 21 1988