

42697-37

06/08/92

CONDITIONAL

X

SAFER GARDEN FUNGICIDE

Mr. Eliot I. Harrison, Agent
Safer, Inc.
9959 Valley View Road
Minneapolis, MN 55344

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit the results of the 12-months storage stability test to the Agency within 15 months after granting this registration since the accelerated storage stability data for 30 days is only acceptable on an interim basis.
3. Make the labeling changes listed below before you release the product for shipment:
 - a. Under Environmental Hazards, change the second sentence to: "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark."

1.

- b. Under Disposal, Change the second sentence to: "Securely wrap the original container in several layers of newspaper and discard in trash."
- c. Add the EPA Registration No. 42697-37.
- 4. Submit five copies of your final printed labeling before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Susan T. Lewis
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure



Garden Fungicide Safer®

CONCENTRATE

Controls powdery mildew, black spot, leaf spot and rust on fruits, vegetables, flowers and foliage.



ACTIVE INGREDIENT:
Sulfur:
INERT INGREDIENTS
TOTAL:

BY WT.
..... 2.0%
..... 88.0%
..... 100.0%

Keep out of reach of children.

CAUTION
(See back panel for additional precautionary statements)

NET CONTENTS 16 FL. OZ.

5456-2731-RM 06058

No. 5456
Garden Fungicide
CONCENTRATE

See Directions for Use, Precautionary Statements and Storage and Disposal information inside.


Ringer Corp. and Safer®, Inc. will not accept liability for damage or injury resulting from misuse.

DISTRIBUTED BY:
Safer®, Inc., a wholly owned subsidiary of Ringer Corp., 9959 Valley View Rd., Eden Prairie, MN 55344
EPA. REG. NO. 42697-

EPA Est. No. 63456-MN-001 44616-MO-001
42705-MN-001

Circled letter corresponds to first lot no. on bottom of container.

©Copyright 1991 Ringer Corp.
5456-2731-RM06058



INTS

BEST AVAILABLE COPY

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

For control of BLACKSPOT, LEAF SPOT, and RUST, mix 1 part concentrate to 30 parts water or 4 fl. oz. (8 tbs) per gal. For POWDERY MILDEW mix 2 fl. oz. (4 tbs) per gal. of water. Place about half of the volume of water in sprayer, add concentrate, stir well, add rest of water and mix thoroughly. Run agitator on large tank sprayers. For coverage, see individual applications. Spray plants thoroughly to wet the upper and lower surfaces of foliage. Repeat as recommended. To control *powdery mildew*, *leaf spot*, *black spot*, and *rust* on FLOWERS, FOLIAGE, PLANTS & ORNAMENTALS, spray at 7 to 10 day intervals or as needed. To control *powdery mildew*, *black spot*, and *rust* on ROSES, spray at 7 to 10 day intervals or following rain. Be sure to treat new growth & buds. Avoid spraying when plants are in full sun or temperature exceeds 85°F. Apply 1 gal. of mix to each 350 sq. ft. To control *powdery mildew* on BEANS & PEAS, spray at first sign of disease & repeat at 7 day intervals as necessary. May be used up to day before harvest. Apply 1 gal. of mix to each 250 sq. ft. To control *powdery mildew* on APPLES & PEARS, apply to thoroughly wet foliage. Apply at pre-bloom stage or after petal fall as needed. CAUTION: Some apple & pear varieties are sulfur sensitive, test for varietal sensitivity prior to full scale treatment. CITRUS Do not apply within 4 weeks following use of an oil spray. To control *powdery mildew* on POTATOES, spray before disease appears & repeat as necessary. Apply 1 gal. of mix to each 350 sq. ft. On CUCUMBERS & SQUASH, spray when disease appears & repeat as necessary. Do not spray young plants

Apply 1 gal. of mix to each 600 sq. ft. On STRAWBERRIES, spray when appears & repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 300 sq. ft. CAUTION: Do not apply to strawberries that will be used for canning. May be used up to the day before harvest. On GRAPES, the cover foliage. Begin when new shoots are 6-10 inches. Repeat before open and at 14 day intervals as needed. CAUTION: Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord & other labrusca varieties. GREENHOUSES: To control *powdery mildew*, *leaf spot* & *rust* on bedding plants, propagates, fruits & vegetables: Mix 1 part concentrate to 30 parts water (per gal.) and apply according to above recommendations by crop type. Avoid phytotoxicity before full-scale application. USE CAUTIONS. Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not use on Boston fern, spinach, apricots, filberts, walnuts, and viburnum. Do not use when temperature exceeds 85°F. Do not apply for at least 4 weeks following application of an oil spray or within 4 days of other pesticides. Be sure to rinse spray equipment thoroughly after each use. Do not allow mixed chemicals to stand in spray equipment. RE-ENTRY: Avoid contact with plants until spray has dried.

INTEGRO™ is protected by the following Patents U.S.A. 4680080, 4592572, 4675062, Canada 422793, and others

Continued-Next Panel

BEST AVAILABLE COPY

as a feed supplement.

Environmental hazards: Do not apply directly to water or swamps, bogs, marshes, and potholes. Do not contaminate water when disposing of rinsate or equipment washwater.

Ringer Corp. and Safer[®], Inc. will not accept liability for damage or injury resulting from misuse.

STORAGE AND DISPOSAL:

Storage: Store in a cool place. Do not contaminate water, food, or feed by storage or disposal.

Disposal: Do not reuse empty container. Wrap the container and put in trash for collection.