

PM 21

42697-37

**Garden  
Fungicide  
CONCENTRATE**

**Safer®**

PM 1/3

10/8/98

**-Controls  
powdery mildew,  
black spot, leaf spot  
and rust on fruits,  
vegetables,  
flowers and  
foliage.**

**SHAKE WELL**

ACTIVE INGREDIENT	BY WT.	KEEP OUT OF REACH OF CHILDREN
Sulfur: .....	12.0%	CAUTION
INERT INGREDIENTS: .....	88.0%	(see back panel for additional
TOTAL: .....	100.0%	precautionary statements)

**NET CONTENTS 16 FL. OZ.**

**Garden Fungicide  
CONCENTRATE**

See Directions for Use, Precautionary Statements and Storage and Disposal information inside.

Ringer Corp. and Safer®, Inc. will not accept liability for damage or injury resulting from misuse. In case of emergency, call 1-800-228-5635 ext. 170.

DISTRIBUTED BY:  
**Safer®, Inc.** a wholly owned  
subsidiary of **Ringer Corp.**,  
9959 Valley View Rd.,  
Eden Prairie, MN 55344  
EPA REG. NO. 42697-37  
EPA EST. NO. 44616-MO-001

<b>ACCEPTED</b>
10/8/98
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 42697-37

Circled letter corresponds to first list no. on bottom of container.

©Copyright 1996 Ringer Corp.

Garden Fungicide  
CONCENTRATE

PRECAUTIONARY STATEMENTS  
Hazard to Humans and Domestic Animals:

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

**If in Eyes:** Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of rinsate or equipment washwater.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SHAKE WELL.** For control of **BLACK SPOT, LEAF SPOT, and RUST**, mix 1 part concentrate to 30 parts water or 4 fl. oz. (8 TBS) per gallon. For **POWDERY MILDEW** mix 2 fl. oz. (4 TBS) per gallon of water. Place about half of the volume of water in sprayer, add concentrate, stir well, add rest of water and mix thoroughly. Run agitator on large tank sprayers. For coverage, see individual applications. Spray plants thoroughly to wet the upper and lower surfaces of foliage. Repeat as recommended.

To control *powdery mildew, leaf spot,* and *rust* on **FLOWERS, FOLIAGE PLANTS & ORNAMENTALS** spray at 7 to 10 day intervals or as needed. To control *powdery mildew, black spot and rust* on **ROSES**, spray at 7 to 10 days intervals or following rain. Be sure to treat new growth & buds. Avoid spraying when plants are in full sun or temperature exceeds 85°F. Apply 1 gal of mix to each 350 sq. ft.

To control *powdery mildew* on **BEANS & PEAS**, spray at first sign of disease & repeat at 7 day intervals as necessary. May be used up to day before harvest. Apply 1 gal. of mix to each 250 sq. ft. To control *powdery mildew* on **APPLES & PEARS**, apply to thoroughly wet foliage. Apply at pre-bloom stage or after petal fall as needed. NOTE: Some apple and pear varieties are sulfur sensitive; test for varietal sensitivity prior to full-scale treatment.

To control *powdery mildew* on **POTATOES**, spray before disease appears & repeat as necessary. Apply 1 gal. of mix to each 350 sq. ft. On **CUCUMBERS and SQUASH**, spray when *powdery mildew* appears & repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 600 sq. ft.

To control *powdery mildew* on **STRAWBERRIES**, spray when disease appears and repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 300 sq. ft. NOTE: Do not apply to strawberries that will be used for canning. May be used up to the day before harvest.

To control powdery mildew on GRAPES, thoroughly cover foliage. Begin when new shoots are 6-10 inches. Repeat before blooms open and at 14 day intervals as needed. NOTE: Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord and other labrusca varieties.

HOME GREENHOUSES. To control *powdery mildew*, *leaf spot* and *rust* on bedding plants, propagates, fruits and vegetables: Mix 1 part concentrate to 30 parts water (8 TBS per gal.) and apply according to above recommendations by crop type. Test for phytotoxicity before full-scale application. NOTE: Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not apply on Boston fern, spinach, apricots, filberts, walnuts and viburnum. Do not apply when temperature exceeds 85°F. Do not apply for at least 4 weeks following application of an oil spray; or within 4 days of other pesticides. Be sure to wash & rinse spray equipment thoroughly after each use. Do not allow mixed spray to stand in spray equipment.

REENTRY: Avoid contact with plants until spray has dried.

**STORAGE AND DISPOSAL:**

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

**Storage:** Store in a cool place.

**Disposal:** Do not reuse empty container. Securely wrap the original container in several layers of newspaper and discard in trash. Rinse empty container well prior to disposal.