


42697-37

09/19/2002

1/4

EPA Reg. No. 42697-37
Draft Label Amendment
July 19, 2002
Page 1 of 4

MASTER LABEL-FRONT PANEL (Page #1 of 1)



SAFER® BRAND GARDEN FUNGICIDE CONCENTRATE

[Additional Brand Names]

Safer® Brand Garden Fungicide For Flowers, Fruit, and Vegetables

Controls powdery mildew, black spot, leaf spot and rust on fruits, vegetables, flowers and foliage.


ACTIVE INGREDIENT:	BY WT.
Sulfur.....	12.0%
OTHER INGREDIENTS:.....	88.0%
TOTAL:.....	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

(See back panel for Precautionary Statements and First Aid)

Net Contents: 16 fl. oz.



MASTER LABEL-BACK PANEL (Page #1 of 3)

PRECAUTIONARY STATEMENTS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of rinsate or equipment washwater.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL. For control of **BLACK SPOT**, **LEAF SPOT**, and **RUST**, mix 1 part concentrate to 30 parts water or 4 fl. oz. (8 TBS) per gallon. For **POWDERY MILDEW** mix 2 fl. oz. (4 TBS) per gallon of water. Place about half of the volume of water in sprayer, add concentrate, stir well, add rest of water and mix thoroughly. Run agitator on large tank sprayers. For coverage, see individual applications. Spray plants thoroughly to wet the upper and lower surfaces of foliage. Repeat as recommended.

To control powdery mildew, leaf spot, and rust on **FLOWERS**, **FOLIAGE PLANTS & ORNAMENTALS** spray at 7 to 10 day intervals or as needed. To control powdery mildew, black spot and rust on **ROSES**, spray at 7 to 10 day intervals or following rain. Be sure to treat new growth & buds. Avoid spraying when plants are in full sun or temperature exceeds 85°F. Apply 1 gal of mix to each 350 sq. ft.

To control powdery mildew on **BEANS & PEAS**, spray at first sign of disease & repeat at 7 day intervals as necessary. May be used up to day before harvest. Apply 1 gal. of mix to each 250 sq. ft. To control powdery mildew on **APPLES & PEARS**, apply to thoroughly wet foliage. Apply at pre-bloom stage or after petal fall as needed. **NOTE:** Some apple and pear varieties are sulfur sensitive; test for varietal sensitivity prior to full-scale treatment.

MASTER LABEL-BACK PANEL (Page #2 of 3)

To control powdery mildew on POTATOES, spray before disease appears & repeat as necessary. Apply 1 gal. of mix to each 350 sq. ft. On CUCUMBERS and SQUASH, spray when powdery mildew appears & repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 600 sq. ft.

To control *powdery mildew* on STRAWBERRIES, spray when disease appears and repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 300 sq. ft. **NOTE:** Do not apply to strawberries that will be used for canning. May be used up to the day before harvest.

To control *powdery mildew* on GRAPES, thoroughly cover foliage. Begin when new shoots are 6-10 inches. Repeat before blooms open and at 14-day intervals as needed. **NOTE:** Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord and other labrusca varieties.

HOME GREENHOUSES: To control *powdery mildew*, *leaf spot* and *rust* on bedding plants, propagates, fruits and vegetables: Mix 1 part concentrate to 30 parts water (8 TBS per gal.) and apply according to above recommendations by crop type. Test for phytotoxicity before full-scale application. **USE CAUTIONS:** Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not apply on Boston fern, spinach, apricots, filberts, walnuts, and viburnum. Do not apply when temperature exceeds 85°F. Do not apply for at least 4 weeks following application of an oil spray or within 4 days of other pesticides. Be sure to wash & rinse spray equipment thoroughly after each use. Do not allow mixed spray to stand in spray equipment.

REENTRY: Avoid contact with plants until spray has dried.

STORAGE AND DISPOSAL:

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

Storage: Store in cool place.

Disposal:

If Empty: Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled: Call your local solid waste agency or (800)-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Woodstream Corp. and its subsidiaries will not accept liability for damage or injury resulting from misuse. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents) call the National Pesticide Information Center at 1-800-858-7378.

EPA Reg. No. 42697-37
Draft Label Amendment
July 19, 2002
Page 4 of 4

MASTER LABEL-BACK PANEL (Page # 3 of 3)

Distributed by
Safer®, Inc., a wholly owned subsidiary of Woodstream Corporation
69 North Locust Street
Lititz PA 17543
1-800-800-1812 for information

EPA Reg. No. 42697-37

H D
EPA Est. No. 44616-MO-01 63416-MN-01
Circled letter corresponds to first letter on lot number on label, or container side or bottom.

All ® Property of Woodstream Corporation

© Copyright 2002 Woodstream Corporation

Made in USA
Printed in USA

