
 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 401 "M" St., S.W. Washington, D.C. 20460</p>	<p>EPA Reg. Number: 42697-57</p>	<p>Date of Issuance: SEP 15 2000</p>
	<p>Term of Issuance: Unconditional</p>	
<p>NOTICE OF PESTICIDE: <u> x </u> Registration <u> </u> Reregistration</p> <p>(under FIFRA, as amended)</p>		
<p>Name and Address of Registrant (include ZIP Code): Safer Brand, Inc. 213 SW Columbia Street Bend, OR 97702-1013</p>		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Make the following label change before you release the product for shipment: <ul style="list-style-type: none"> a) Revise the EPA Registration Number to read, "EPA Reg. No. 42697-57." 		
<p>Signature of Approving Official:  George T. LaRocca Insecticide Branch Registration Division (7505C)</p>	<p>Date: SEP 15 2000</p>	

b) Add to your First Aid Statement: Take off contaminated clothing. Rinse skin immediately with plenty of soap and water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

c) Following the pest claim termites on front panel, Add in parenthesis (localized control only).

d) The MRIDs you have cited for scorpions, wasps, and Smokey Brown Cockroaches were for studies that were found unacceptable by the Agency. Additionally, the MRIDs you cited for Deer Ticks, sowbugs and Fire Ants are unacceptable because the % active ingredient was at a higher level than your product. We could also not find any efficacy data in your matrix to support fleas, carpenter bees, and bedbugs. Please remove these pests from your label or submit or cite efficacy data to support these 10 pests. Please note, deer tick data and smokey brown cockroach data will not support general tick and cockroach claims. Additionally citations of other species will be necessary. We suggest you check with your supplier of the active ingredient in this regard.

e) Revise the second paragraph under "Termites" to read as follows: For subterranean termites the purpose of such applications is to kill workers or the winged forms which may be present in the treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement. For active subterranean termite infestations get a professional inspection.

3. Please submit (2) copies of your final printed label before you release your product for shipment. A stamped copy is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide Branch
Registration Division (7505C)

Enclose

SAFER BRAND'S®
DELTAMETHRIN INSECT CONTROL DUST

ACCEPTED
with COMMENTS
In EPA Letter Dated
SEP 15 2000

42697-57

- CONTROLS FIRE ANTS
- FOR INDOOR /OUTDOOR RESIDENTIAL USE ONLY
- FOR HOUSEHOLD USE
- HOUSEHOLD ORNAMENTAL GARDEN INSECT CONTROL
- CONTROLS FIRE ANTS, FLEAS, TICKS, TERMITES AND MORE
- KILLS INSECTS UP TO 8 MONTHS
- DOES NOT CLUMP OR ABSORB MOISTURE
- USE ON ROSES, FLOWERS AND ORNAMENTALS
- CONTROLS MANY DIFFERENT INSECT PESTS
- TO CONTROL IMPORTED FIRE ANTS ON LAWNS AND OTHER RESIDENTIAL AREAS
- NON-STAINING AND ODORLESS
- LONG-LASTING RESIDUAL ACTION
- LONG-LASTING
- 8 OZ. TREATS 15 FIRE ANT MOUNDS
- 1 LB. TREATS 30 FIRE ANT MOUNDS
- 2 LBS. TREATS 60 FIRE ANT MOUNDS
- FAST, EFFECTIVE KILL OF FIRE ANTS
- FAST, EFFECTIVE KILL OF STINGING ANTS
- EASY TO USE
- CONVENIENT
- KILLS FIRE ANTS ON CONTACTS

ACTIVE INGREDIENT:

Deltamethrin: [(s)-alpha-cyano-3-phenoxybenzyl-(1R,3R)-3-(2,2 dibromovinyl)-2,2-dimethyl

cyclopropanecarboxylate.....0.05%

INERT INGREDIENTS:..... 99.95%

Total.....100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Side Panel For Additional Precautionary Statements

Distributed By:

Safer Inc., A Wholly Owned Subsidiary of Verdant Brands, Inc.,
9555 James Ave. South, Suite 200,
Bloomington, MN 55431

EPA Reg. No. 42697-
EPA Est. No.

Net Contents:

For information on this product (including health concerns, medical emergencies or pesticide incidents) call the National Pesticide
Tele-communications Network at 1-800/858-7378.

PRECAUTIONARY STATEMENTS
Hazard To Humans And Domestic Animals

CAUTION: May be harmful if absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

Environmental Hazards

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water. Use with care when applying in areas adjacent to any body of water. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

STORAGE AND DISPOSAL

Storage: Store product in original container in a cool, dry, locked place out of reach of children.
Container Disposal: Do not reuse empty containers. Wrap in several layers of newspaper and discard in trash.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read the entire Directions for Use and Precautions before using this product. Deltamethrin Insect Control Dust is a ready-to-use insecticide dust which exhibits effective knockdown and long-lasting control against the pests listed when used according to label directions. Deltamethrin Insect Control Dust can be used for long-lasting pest control in and around homes and their immediate surrounding. Permitted areas of use include but are not limited to: animal quarters, apartments, household greenhouses, houses, mobile homes and recreational vehicles and ornaments.

This product is intended for applications with a hand duster or from the shaker can, with a paint brush or by other suitable means, to hiding and runway areas and other places where insects are found.

INDOOR USE: In living areas, make applications in such a manner as to avoid deposits on exposed surfaces or introducing the material into the air. To apply insecticide directly into cracks and crevices, use a bulbous duster or other suitable equipment. Apply lightly and uniformly to infested areas. Pay particular attention to cracks and crevices; service ducts; false floors and ceilings; wall voids; around electrical and telephone fittings and equipment; around water and sewer pipes; under and behind cabinets; refrigerators and sinks; around windows and door frames; along baseboards; in attics and crawl spaces. The amount to be applied will vary with the site but usually will be in the range of 2-3 grams of product per square yard (or 0.5 lbs. per 1000 sq. ft.). Repeat as necessary to maintain adequate control.

ANTS, BOXELDER BUGS, COCKROACHES (INCLUDING BOTH RESISTANT AND NON-RESISTANT STRAINS), CRICKETS, FLEAS, FIREBRATS AND SILVERFISH: Apply Deltamethrin Insect Control Dust thoroughly to all areas where these pests crawl and hide, especially in those areas described above. For ants, apply to ant trails, around doors and windows and other places where ants enter premises.

TERMITES, CARPENTER ANTS AND CARPENTER BEES: Apply to voids or channels in damaged wooden members of a structure; to cracks; spaces or bearing joints between wooden members of structures; or between such members and the foundation; and in locations vulnerable to termite attack. Drilling may be necessary prior to applying the dust in these locations, but it must be recognized that such drilling and

treating of timber in a particular location may not result in complete coverage of infested and vulnerable voids or galleries. This dust can be applied in attics and crawl spaces to prevent entry of drywood and formosan termites at a rate of 0.5 lb. per 1,000 sq. ft. of floor area. In these treatments, all vents and blowers should be closed. Inspect at 6 months and one year to ensure that colony has been killed. Retreat if necessary.

The purpose of such applications is to kill workers or winged forms which may be present in the treated channels at the time of treatment and to impart a temporary resistance at these places to attack which may aid in the prevention of immediate reinfestation. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement. This supplemental treatment can be particularly supportive of new construction.

CENTIPEDES, GROUND BEETLES, MILLIPEDES, SCORPIONS, SPIDERS AND SOWBUGS: Apply dust along and behind baseboards, to window and door frames, corner, pipes, storage locations, attics, crawl spaces, and other areas where these pests may enter and crawl.

WASPS AND BEES: It is generally advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Wear protective clothing if deemed necessary to avoid stings. Using hand or power duster (Actisol or similar equipment) or other suitable means, with extension tubes if necessary, thoroughly dust nest, entrance and surrounding areas where insects alight. Nests in wall voids can be located by listening with your ear against the wall. Drill a hole in the area, blow dust in and reseal. For ground wasps, dust entrance and surrounding areas. For best results check nests carefully one or two days after treatment to ensure complete kill, then remove and destroy nest to prevent emergence of newly-hatched insects. If removal is not feasible, retreat the nest, if necessary. Do not use this product to treat overhead nests.

WEBBING CLOTHES MOTHS: Use as a crack and crevice treatment in closets and other storage areas where these pests are found. This is an adjunct treatment and will not control moth larvae already on the clothes. DO NOT apply this product directly to clothing.

CARPET BEETLES (BLACK, FURNITURE, COMMON AND VARIED): Dust where these insects are found, for example: on and under the edges of floor coverings, around baseboards under rugs and furniture, and closets.

BEDBUGS: Apply dust thoroughly to all areas where these pests are typically found including: folds and tufts of mattresses, coils of springs, cracks and hollow posts of bedsteads, upholstery of chairs and sofas, picture frame moldings and all cracks and crevices in the room should be treated. Allow powder to remain in contact with surface for 4-6 hours. Thoroughly vacuum treated areas, dispose of vacuum bag, and place clean linens on bed.

STORE PRODUCT PESTS:

To aid in the control of exposed stages of:

Cadelles	Merchant grain beetles
Cheese mites	Red flour beetles
Cigarette beetles	Rice weevils
Confused flour beetles	Rusty grain beetles
Dark meal worms	Saw-toothed grain beetles
Drugstore beetles	Spider beetles
Grain mites	Tobacco moths
Granary weevils	Yellow meal worms
Lesser grain borers	

In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed. Clean up waste materials, dust, dirt and other debris. Thoroughly treat floors, walls and other surfaces in storage and handling cupboards, and other difficult to reach areas where insects may hide. DO NOT apply directly to food, utensils or on areas where food is stored.

ORNAMENTAL PLANT PESTS:

For control of :

Aphids	Leafrollers
Beet armyworms	Orange tortrix
Blister beetles	Periodical cicada
Boxelder bugs	Plant bugs
Certain leafminers	Rose chafer
(Boxwood, Holly	Rose midge
Oak, Spruce Needle)	Rose slugs
Elm leaf beetle	Tent caterpillars
Flea beetles	Thrips
Japanese beetle	Weevils
Leafhoppers	

SHRUBS AND ORNAMENTALS: Apply when insects or their damage first appear and repeat at weekly intervals or as necessary. Apply 8 ounces per 1,000 sq. ft.; or thoroughly apply to place a thin layer of dust on both upper and lower sides of stems, leaves and flowers.

ROSES, PETUNIAS, CHRYSANTHEMUMS, GERANIUMS, CARNATIONS AND OTHER RELATED FLOWERS: To control many leaf and flower chewing insects, apply as soon as the insect appears and repeat at 7 day intervals as long as insects continue to appear. Apply a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

OUTDOOR USE

PERIMETER TREATMENT: Use Deltamethrin Insect Control Dust as a residual treatment for control of ants, bees, centipedes, cockroaches, crickets, firebrats, fleas, ground beetles, millipedes, scorpions, silverfish, slugs, sowbugs, spiders, ticks and wasps. Make a residual treatment around windows and doors, porches, screens, eaves, patios, garages, under stairways, in crawl spaces, and other areas where pests hide. To help prevent the invasion of buildings and structures by these pests, apply thoroughly and uniformly to the foundation and crawl spaces where pests are active and may find entrance.

FLEAS AND TICKS IN RODENT BURROWS: To control fleas and ticks in and around rodent burrows around homes and other buildings, apply dust with a shaker can. Thoroughly apply dust to adequately cover mound and around the perimeter (18" from the outer edge). Be sure to treat abandoned rodent burrows which are suspected of containing fleas and ticks. Treat for a minimum of 5 days for light rodent populations, 10 days when populations are dense.

USE DIRECTIONS FOR FIRE ANT TREATMENT

IMPORTED FIRE ANT MOUND TREATMENT: Apply with a shaker can. Thoroughly apply dust to adequately cover the mound and around the perimeter (18" from the outer edge of the mound). A second treatment may be necessary if complete control is not obtained. **DO NOT DISTURB THE MOUND PRIOR TO APPLICATION.** If possible, treat all colonies in the vicinity. Also treat colonies which have not yet constructed a mound.

[Alternate directions: - Evenly sprinkle one tablespoon (approximately ½ oz.) of Deltamethrin Insect Control Dust over the top of each mound. Do not water in or disturb the mound. Apply when temperatures are between 65°F and 80°F. Allow 3 to 4 days for maximum control, then repeat as necessary. Food utensils such as a teaspoon or measuring cup should not be used for food after use with pesticides.]

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with label directions. If you are not completely satisfied with this product, or for consumer information, call (952) 703-3300 weekdays 9-5 Central Time to arrange a refund for the purchase or replacement of the product. Proof of purchase is required.