
42750-14





UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

MAY 3 1 2002

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES 117

Albaugh, Inc. c/o Michael Kellogg Pyxis Regulatory Consulting 11324 17th Ave. Ct. N.W. Gig Harbor, WA 98332

Gentlemen:

Subject: Delete Use on Rice Albaugh MCPA Amine 4 EPA Registration No. 42750-14 Your Submission Dated August 7, 2001

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. Use "Chemical Resistant Gioves Category A" instead of "Waterproof Gloves" wherever it appears on the labeling.

b. In the Environmental Hazards Section delete "For terrestrial uses,". Refer to PR Notice 93-8.

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records. The amended labeling supersedes all previously accepted ones. If you have any questions concerning this letter, please contact Mr. James Stone at 703-305-7391.

Sincerely yours,

Joanne I. Miller Product Manager (23) Herbicide Branch Registration Division (7505C)

Enclosure

## ALBAUGH MCPA AMINE 4

## HERBICIDE FOR SELECTIVE CONTROL OF CERTAIN WEEDS IN WHEAT, OATS, BARLEY, RYE, RICE, FLAX, ESTABLISHED GRASSLANDS AND NON-CROP AREAS

#### **ACTIVE INGREDIENT:**

Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid*	
OTHER INGREDIENTS:	
TOTAL	

\*This product contains 3.7 pounds 2-methyl-4-chlorophenoxyacetic acid per gallon or 39.9%. Isomer specific by AOAC Method No. 6.A-18-22 (13th Ed.).

## EPA RegACCEPTED<sup>4</sup> with COMMENTS In EPA Letter Dated:

EPA Est. No. 42750-MO-1

KEEP OUT OF REACH OF CHILDREN

MAY 3 1 2002 Under the Federal Insecticide,

# **DANGER - PELIGRO**

Si ust de not reprivate paratitiqueta, bi-sque a alguien para que se la explique a usted en detalis. (if you do material contratte de la detalis). (if you do material contratte de la detalis).

FIRST AID		
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes.	
	• Remove contact lenses, if present, after the first 5 minutes, then continue	
	rinsing eye.	
	Call a poison control center or doctor for treatment advice.	
If swallowed	Call a poison control center or doctor immediately for treatment advice.	
<ul> <li>Have person sip a glass of water if able to swallow.</li> </ul>		
	• Do not induce vomiting unless told to do so by the poison control center or	
	doctor.	
	Do not give anything by mouth to an unconscious person.	
If inhaled	Move person to fresh air.	
	• If person is not breathing, call 911 or an ambulance, then give artificial	
	respiration, preferably mouth-to-mouth, if possible.	
	Call a poison control center or doctor for further treatment advice.	
If on skin or	Take off containinated clothing.	
• Rinse skin immediately with plenty of water for 15-20 minutes.		
Call a poison control center or doctor for treatment advice.		
HOT LINE NUMBER		
Have the product container or label with you when calling a poison control center or doctor, or going		
for <b>treatm</b> ent.	You may also contact 1-800-424-9300 for emergency medical treatment	
information_NO4T-TO-PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric		
lavage.		
NOTE TO PHYSICIAN		
Probable mussis damage may contraindicate the use of gastric lavage.		

See additional Precautionary Statements on side panel.

Manufactured By: Albaugh, Inc.

NET CONTENTS \_\_\_\_\_ Gals. Ankeny, Iowa 50021

\_\_\_\_ Liters

#### FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE CALL CHEMTREC (800) 424-9300

#### PRECAUTIONARY STATEMENTS DANGER HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Harmful if swallowed, absorbed through skin or inhaled.

Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist.

## PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eyewear, and chemical-resistant headgear for overhead exposure.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS

## USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Drift or run-off may adversely affect nontarget plant. For terrestrial uses, to not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Avoid use of small-diameter spray nozzles. Do not apply when weather conditions favor drift from target area, as this product may injure cotton, beans, other vegetables, certain legumes and ornamentals.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Do not use, pour, spill or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

2

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks, protective eyewear, and chemical-resistant headgear for overhead exposure.

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter treated areas without protective clothing until sprays have dried.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

## STORAGE

Store in a dry location away from children, animals, foods, feeds, seeds, or other agricultural chemicals. Do not store below 40°F or expose to subfreezing temperatures. Handle in accordance with information given under "Precautionary Statements." In the event of spillage or leakage, soak up material with absorbent clay, sand, sawdust or other absorbent material. Scrape up and dispose of in accordance with information given under "Disposal." Repackage and relabel useable product in a sound container. In case of fire or other emergency, report at once by toll-free telephone to 800-424-9300.

## DISPOSAL

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

## **CONTAINER DISPOSAL**

**METAL CONTAINERS** - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**PLASTIC CONTAINERS -** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

4/1

3

#### USE PRECAUTIONS

Under no circumstances should MCPA Amine Herbicide be used in the vicinity of vegetables, flowers, grapes, tomatoes, cotton, beans and other legumes, etc., or other susceptible crops as severe damage may result. Extreme care must be exercised to prevent drifting of this material. Do not apply on windy days. Coarse sprays are less likely to drift. Do not apply this product through any type of irrigation system. Do not use around the home, recreation areas or similar sites. Equipment used to apply MCPA Amine should not be used to apply other agricultural chemicals to susceptible crops. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources.

#### AMOUNT OF SPRAY TO APPLY

Apply 5 to 15 gallons of total spray per acre when making applications with ground equipment and 3 to 5 gallons of total spray per acre when making applications by aircraft, unless directed otherwise under specific directions. Carefully read the Environmental Hazards section of the Precautionary Statements for further information on spray volume.

#### SUSCEPTIBLE (MOST AREAS):

Cock bur Dragonhead mint Field peppergrass Fanweed Goatsbeard Lambsquarters (goosefoot)

#### LESS SUSCEPTIBLE:

Buttercups Canada thistle Dandelions Hempnettle Kochia Pigweed Plantains Purslane Russian pigweed Sow thistle Stinging nettle Sunflower

#### **CONTROLS THESE WEEDS**

Marshelder Mustards (annual) Pennycress Puncturevine Ragweed Shepherdspurse Stinkweed Wild radish Yellow rocket or wintercress and many others

Vetch White top and hoary cress and many others

#### **DIRECTIONS FOR USE**

SELECTIVE SPRAYING: Wheat, Oats, Barley and Rye - Apply as a water mix spray by ground sprayer or airplane. Use 1/2 to 1 pint per acre for the more susceptible weeds after crop has reached the 3 to 4 leaf stage up to boot stage. Use up to 3 pints per acre for less susceptible weeds after crop has tillered and up to early boot stage. Do not spray from boot to dough stage. For small grains underseeded with legumes, see the following paragraph.

SMALL GRAINS: Underseeded with Alfalfa, Lespedeza, Red and White Clover: For emergency control of serious infestations of Mustard, Yellow Rocket and other susceptible broadleaf weeds, apply 1/4 to 1/2 pint in not more than 5 to 10 gallons of water per acre. Make application after cereal is well tillered (4 to 8 inches tall) and before reaching boot stage. The nurse crop and weeds should provide a protecting canopy which, together with the use of low gallonage applied at low pressure, will reduce the risk of damage to the legumes. Do not apply to small grains underseeded with Vetch or Sweet Clover, which are very susceptible. There is also some risk to other legumes if only thinly protected by a canopy. Do not forage or graze meat animals on treated areas within 7 days of slaughter.

**RED CLOVER AND ALFALFA (for use in the Pacific Northwest only):** For control of Yellow Rocket and other susceptible annual weeds such as Pennycress or Fanweed, use 1 pint per acre of MCPA Amine. Only use 1/2 pint per acre on new stands after Clover has two or more true leaves. Old stands of Red Clover may be retarded by application of MCPA. Apply in late fall following frosts when the legumes are dormant before active growth starts. The temperature at the time of spraying should be above 40° F.

۰.

**CANNING PEAS (for use in the Pacific Northwest only)**: Use MCPA Amine at 1/4 to 3/4 pint in sufficient water to cover 1 acre. Apply to peas after the 3 node stage and before the first pea flowering. Do not apply during bloom period of crop. To control Canada Thistle, use 1/2 to 3/4 pint of MCPA Amine per acre. Peas may be injured somewhat at the higher rate of application, but if thistle growth is heavy, control will more than compensate for injury to peas.

**RICE (for use in California only):** For early season broadleaf weed and sedge control in rice, apply 1 to 11/2 pints of MCPA Amine Herbicide per acre by air in 8 to 10 gallons of total spray volume when rice is in the 3 to 4 leaf stage, and with well established root systems, usually 18 to 24 days after planting. Applications made during hot weather or at the higher dosage range may cause temporary stunting of the rice. For maximum effectiveness the weeds should be exposed at time of treatment. If retreatment is necessary, usually 35 to 65 days after planting, the total amount per acre of MCPA Amine applied in the <sup>3</sup> two applications must not exceed 3 pints in one treatment. Do not grow crayfish or catfish in treated rice fields.

**FLAX**: Use 1/4 to 1/2 pint. Apply by ground sprayer or by airplane. Use sufficient water to give uniform and adequate coverage. Apply only when weeds are up and when flax is 4 to 8 inches high and before it comes into bud. Treatment after early bud stage may result in severe damage. If Canada Thistle is present, it may be necessary to go as high as 3/4 pint per acre to prevent seed head production. Some injury to the flax may result. Do not forage or graze meat animals on treated areas within 7 days of slaughter.

**ESTABLISHED GRASSLAND AND PASTURES**: Use 1 to 3 pints per acre in sufficient water (10 to 100 gallons) in airplane or ground sprayer application and give thorough coverage. Use higher rate for White Top, Canada Thistle and other less easy-to-kill weeds; spray perennials in early bud to full bloom stage and also regrowth in fall. Spray other weeds in spring or fall. Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not forage or graze dairy animals on treated areas within 7 days after treatment.

**GRASSES GROWN FOR SEED**: Use 1 to 2 pints per acre in sufficient water to give adequate coverage. Use higher rate where weed stands are heavy. In established grasses, apply in spring before head comes into boot and on seedling grass after grass has tillered.

**NOTE:** In weed control in grasses, repeat treatment may be needed for less susceptible weeds. White Clover and other legumes may be temporarily injured or killed. In some areas, Bent, Buffalo, Carpet and St. Augustine grasses may also be injured by the treatment.

**RANGE**: For control of Whitebrush - Use 1-1/4 quarts in a mixture of 1 gallon of diesel oil and sufficient water to make 8 gallons of solution per acre. Apply in spring or fall under good moisture conditions, full leaf, before blossoms begin to fall.

6/7

#### NON-CROP SPRAYING

60

<sub>6</sub> 7/7

CANADA THISTLE, WHITE TOP AND MEADOW BUTTERCUP (In non-crop areas such as Roadsides, Fence Rows, Rights-of-Way and similar places): For spot treatment, use 1/4 pint in 3 to 4 gallons of water, or 3/4 gallon per acre in sufficient water to give coverage for most extensive areas. Spray to wet weeds thoroughly when in bud to early bloom and again on fall regrowth. Do not forage or graze livestock or dairy animals on treated areas within 7 days of treatment. Local conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station Weed Specialists for specific recommendations for local weed problems and for information on possible lower dosages.

**Before buying or using this product**, read "Conditions of Sale and Warranty" below on this label. If terms are not acceptable, return **unopened** package at once to seller for full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under the Conditions of Sale and Warranty.

#### CONDITIONS OF SALE AND WARRANTY

The **DIRECTIONS FOR USE** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, INC. or the Seller. All such risks shall be assumed by the Buyer.

ALBAUGH, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. ALBAUGH, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. IN NO CASE SHALL ALBAUGH, INC. OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALBAUGH, INC. and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of ALBAUGH, INC. No employee or agent of ALBAUGH, INC. or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.