



Reg # 42750-19

PM-23

12/18

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

MAR 18 1994

ALICE WALKER
ALBAUGH INC.
1517 N. ANKENY BLVD.
ANKENY, IA. 50021

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 12/6/93 Response to PR Notice 93-7
EPA Reg. No. 42750-19
2,4-D AMINE NO. 4 HERBICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber

Page 2


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,


Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

3918

ALBAUGH, INC.

2,4-D AMINE NO. 4

HERBICIDE

ACCEPTED
with COMMENTS
in EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

42750-19

ACTIVE INGREDIENT:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid.....46.8%

INERT INGREDIENTS:.....53.2%

TOTAL100.0%

*Equivalent to 38.9% of 2,4-Dichlorophenoxyacetic acid or 3.8 lb./gal. Isomer specific by AOAC Method.

Before buying or using this product, read "Warranty Limitations and Disclaimer" elsewhere on this label. If terms are not acceptable, return unopened package at once to seller for full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under the Warranty Limitations and Disclaimer.

EPA Reg. No. 42750-19

EPA Est. No. 42750-MO-1

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

~~PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.~~

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN: Wash contaminated areas thoroughly with soap and water. ~~Do not re-use contaminated clothing until washed.~~ Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of clean water for 15 minutes. Get medical attention.

Manufactured By:
Albaugh, Inc.
1517 N. Ankeny Blvd.
Ankeny, Iowa 50021

NET CONTENTS
____ GALS.

**FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE
CALL CHEMTREC (800) 424-9300**

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS
DANGER
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Causes irreversible eye damage. Harmful if swallowed. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. In case of contact, wash skin with plenty of soap and water. ~~For eyes, flush with water for at least 15 minutes, then get medical attention. For skin, get medical attention if irritation persists. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.~~

~~When mixing, loading, or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical-resistant gloves (nitrile), long sleeved shirt, long pants, socks and shoes. For aerial applicators in enclosed cockpits and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required.~~

~~Wash hands, face and arms with soap and water as soon as possible after mixing, loading, or applying this product. Wash hands, face and arms with soap and water before eating, smoking, drinking, or using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.~~

~~If this container contains over 1 gallon and less than 5 gallons, persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.~~

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear, chemical-resistant headgear for overhead exposure, and chemical-resistant apron when cleaning equipment, mixing, or loading. ~~It is recommended that eye protection include front, brow and temple protection.~~

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

ENGINEERING CONTROLS STATEMENTS

If this container contains 5 gallons or more in capacity, a closed mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed

BEST AVAILABLE COPY

before removal. The mechanical system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)]. The handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use ~~closed systems~~, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Reclose all partially used containers by thoroughly tightening screw cap. Absorb any spill with a suitable clay absorbent and dispose of as indicated under "Pesticide Disposal."

Protect from freezing. If stored below freezing, the product must be warmed to at least 70°F and agitated before using. This does not affect the efficiency of the product.

For safety and prevention of unauthorized use, all pesticides should be stored in locked facilities. To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification.

Opened, partially used pesticides should be stored in original labeled containers when possible. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of the new container.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

METAL CONTAINERS: Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

PLASTIC CONTAINERS: Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear, and chemical-resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

~~Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your state Department of Agriculture for further information.~~

~~Re-Entry for Turf Sites (excluding sod farms): Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.~~

~~For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: eye protection (face shield or safety glasses), chemical-resistant gloves (nitrile), long-sleeved shirt, long pants, socks, and shoes.~~

~~Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. The front panel PRECAUTIONARY STATEMENTS should be read to workers as well as the instruction not to enter until spray has dried. When oral warnings are given, warnings shall be given in a language customarily understood by the workers. Written warnings must include the following information:~~

~~DANGER: Area treated with 2,4-D on (date of application). Do not enter~~

BEST AVAILABLE COPY

6 8 9 18

~~without appropriate protective clothing until sprays have dried. (Insert here Statements of Practical Treatment as on front panel).~~

GENERAL INFORMATION

Performance of this product may be affected by local conditions, crop varieties, and application method. User should consult local Extension Service, Agricultural Experiment, or University Weed Specialists, and state regulatory agencies for recommendations in your area.

Best results are obtained when product is applied to young succulent weeds that are actively growing. The lower recommended rates will be satisfactory on susceptible annual weeds. For perennial weeds and conditions such as the very dry areas of the western states, where control is difficult, the higher recommended rates should be used.

When product is used for weed control in crops, the growth state of the crop must be considered.

Some plants and weeds, especially woody varieties, are difficult to control and may require repeat applications.

Application rates should be 1 to 5 gallons of total spray by air or 5 to 25 gallons by ground equipment unless otherwise directed. In either case, use the same amount of 2,4-D recommended per acre. For crop uses, do not mix with oil, surfactants, or other adjuvants unless specifically recommended. To do so may reduce herbicide's selectivity and could result in crop damage.

Aerial applications should be used only when there is no danger of drift to susceptible crops. Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. This product contains Dimethylamine salt of 2,4-D, one of the least volatile forms of 2,4-D.

Because coarse sprays are less likely to drift than fine, do not use equipment (such as hollow cone small orifice nozzles) or conditions (such as high pressure) that produce such sprays.

Product should not be allowed to come into contact with desirable, susceptible plants such as beans, cotton, fruit trees, grapes, legumes, ornamentals, peas, tomatoes, and other vegetables. Product should not be used in greenhouses. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Users should note that herbicide treatment of public water requires a permit from appropriate state agencies in most states. Your state Conservation Department, or Game and Fish Commission will aid you in securing a permit in your state.

If stored below freezing, product should be warmed to 70°F and agitated before

using. This does not affect the efficiency of the product.

Spray equipment used to apply 2,4-D should not be used for any other purpose until thoroughly cleaned by a suitable chemical cleaner.

Spray Preparation: Add the recommended amount of product to approximately 1/2 the volume of water to be used for spraying. Agitate well, then add the remainder of the water. Continue agitation during application until spray tank is empty.

Use in Liquid Nitrogen Fertilizer: Product may be combined with liquid nitrogen fertilizer suitable for foliar application on corn, grass, pastures, or small grains in one operation. Use product according to directions on this label for those crops. Use liquid nitrogen fertilizer at rates recommended by supplier or Extension Service Specialist. Mix the product and fertilizer according to the following instructions:

Fill the spray tank approximately 1/2 full with the liquid nitrogen fertilizer. In a separate clean container, mix the amount of product to be used with an equal amount of water. Add the product mixture to the spray tank while agitating. Add the remainder of the fertilizer while continuing to agitate. Apply immediately, maintaining agitation during application until tank is empty. **DO NOT APPLY DURING COLD (NEAR FREEZING) WEATHER.** Spray mixture must be used immediately and may not be stored.

Note: Pre-mixing the product with an equal amount of water is important.

WHERE TO USE

This product is used to control broadleaf weeds in cereal crops, corn, and sorghum; weeds, and brush in rangelands, pastures, rights-of-way, and similar noncrop uses; tree injection and for aquatic weed control.

RECOMMENDED RATES OF PRODUCT PER ACRE**

Crop	Normal Rates (usually safe to crop)	Higher Rates for Special Situations* (more likely to injure crop)
<hr/>		
Small Grains:		
Spring Postemergent		
wheat, barley, rye	2/3 to 1-1/2 pts.	1 to 3 pints
oats	1/2 to 1 pint	1-1/2 to 2 pts.
Preharvest (dough stage)		
wheat, barley, oats	1 to 2 pints	2 to 3 pints
<hr/>		

Corn:

Preemergent	2 to 4 pints	
Emergent	1 pint	1-1/2 pints
Postemergent-up to 8 inches tall	1/2 to 1 pint	
8 inches to tasseling (use only directed spray)	1 pint	1-1/2 to 2-1/2 pts.
Preharvest	1 to 2 pints	

Sorghum:

Postemergent		
6 to 8 inches tall	2/3 to 1 pint	
8 to 15 inches tall (use only directed spray)	1 pint	1-1/2 to 2 pints

Rice:

1 to 2-1/2 pints	2 to 3 pints
------------------	--------------

Sugarcane:

Preemergent	4 pints
Postemergent	1-1/2 to 2 pints

NOTE: The higher rates as recommended above may be necessary to control difficult weed problems, such as dry conditions in the Western States. They should not be used, however, unless possible crop injury is acceptable. User should consult local Extension Service or Agriculture Experiment Station Weed Specialist for recommendations on special conditions.

*Arizona, Idaho, Montana, Oregon, Utah, Washington, Wyoming.

**If band treatment is used, base the dosage rate on the actual area sprayed.

WEEDS CONTROLLED

When used properly, product will kill or control the following, in addition to many other noxious plants susceptible to 2,4-D:

Alligatorweed	Dandelion	Pokeweed
Arrowhead	Dock	Povertyweed
Artichoke	Duckweed	Puncturevine
Bindweed	Elderberry	Purslane
(hedge, field and European)	Goldenrod	Rush
Bitter wintercress	Ground ivy	Russian thistle
Boxelder	Hemp	Sagebrush
Buckhorn	Hoary cress	Shepherdspurse
Bull thistle	Honeysuckle	Sowthistle
	Indigo	Stinkweed

Bulrush
Burdock
Bur ragweed
Buttercup
Catnip
Chickweed
Chickory
Cocklebur
Coffeebean
Creeping jenny
Curly indigo

Ironweed
Jimson weed
Lambsquarters
Locoweed
Mexican weed
Morningglory
Mustard
Parrotfeather
Pennywort
Pigweed
Plantain

Sumac
Sunflower
Virginia creeper
Waterhyacinth
Waterlily
Waterprimrose
Wild lettuce
Wild radish
Willow

LESS SUSCEPTIBLE WEEDS

Kochia
Pigweed (hybrid)

Poison ivy
Smartweed

Wild garlic
Wild onion

CROPS

Small grains (barley, oats, wheat, rye), not underseeded with a legume: See table for recommended use rates. Spray when weeds are small after grain begins tillering but before boot stage (usually 4 to 8 inches tall). Do not apply before the tiller stage nor from early boot through milk stage. To control large weeds that will interfere with harvest or to suppress perennial weeds, preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well.

Spring Planted Oats: Apply in sufficient water to give good coverage. Apply after the fully tillered stage, except during the boot to dough stage.

Fall Planted Oats: Apply after full tillering but before early boot stage. Some difficult weeds may require the higher rates of 1 to 1-1/2 pints per acre for maximum control but injury may result. Do not spray during or immediately following cold weather.

Note: Oats are less tolerant to 2,4-D than wheat or barley and more likely to be injured.

Note: Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 14 days after treatment. Do not feed treated straw to livestock.

For Emergency Weed Control in Wheat: Perennial broadleaf weeds apply 3 pints per acre when weeds are approaching bud stage. Do not spray grain in the boot to dough stage. The 3 pint per acre application can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury.

Wild Garlic in Grain Stubble: To prevent new growth of garlic following harvest, apply 2 to 3 quarts of product per acre to stubble. Do not forage for 14 days following application. Do not plant any crop for 3 months after treatment or until 2,4-D has disappeared from soil.

Corn: See table for recommended use rates.

Preemergent: Apply product from 3 to 5 days after planting but before corn emerges. Do not use on very light, sandy soils. Use the higher rates on heavy soils. Plant corn as deep as practical.

Post Emergent: Best results are usually obtained when weeds are small and corn is 5 to 18 inches tall. When corn is over 8 inches tall, use drop nozzles to keep spray off corn foliage as much as possible. Do not apply from tasseling to dough stage. If corn is growing rapidly and temperature and soil moisture is high, use 1/2 pint per acre rate to reduce possibility of crop damage. Delay cultivation for 8 to 10 days to prevent stalk breakage due to temporary brittleness caused by 2,4-D. Application rates of up to 1 pint per acre may be used to control some hard to control weeds. However, the possibility of injury to the corn is increased.

Do not use with atrazine, oil or other adjuvants. Since the tolerance to 2,4-D of individual hybrids varies, consult your seed supplier, local Extension Service, Agricultural Experiment Station, or University Weed Specialist for information.

Pre-Harvest: After the hard dough or denting stage, apply 1 to 2 pints of product per acre by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as Bindweed, Cocklebur, Dogbane, Jimsonweed, Ragweed, Sunflower, Velvetleaf, and vines that interfere with harvesting. Do not forage or feed corn fodder to livestock for 7 days following application.

Note: Hybrids vary in tolerance to 2,4-D. Some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for this information.

Rice: See table for recommended use rates. Apply in the later tillering stage of rice development at the time of first joint development (first to second green ring) usually 6 to 9 weeks after emergence. Do not apply after panicle initiation, after rice internodes exceed 1/2 inch, at early seedling, early panicle, boot, flowering, or early heading growth stages.

Note: Some rice varieties under certain conditions can be injured by 2,4-D. Therefore, before spraying consult local Extension Service or university specialists for appropriate rates and timing of 2,4-D sprays.

Sorghum (Milo): See table for recommended rate.

Apply to sorghum when crop is 4 to 12 inches high with secondary roots well established. Use drop nozzles when crop is over 10 inches high. Do not apply from flowering to dough stage. Rates of up to 1 pint per acre may be used to control some hard to control weeds. However, the chance of crop injury increased with the higher rates. Do not use with oil. Use lower rate if conditions of high temperature and high soil moisture exist.

Soybeans (Preplant Only) - For Use in Crop Residue Management Systems: Apply 1 pint **not less than 15 days** prior to planting soybeans or 2 pints **not less than 30 days** prior to planting. Apply to postemergent weeds when small, actively growing, and free of stress caused by extremes in climatic conditions, diseases, or insect damage. The response of individual weed species is variable. Consult your local county agent or state Agricultural Extension Service or crop consultant for advice. Use the higher rate on larger weeds and when perennials are present. (See WEEDS CONTROLLED below.)

Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of water per acre in ground equipment.

WEEDS CONTROLLED

alfalfa*	horseweed or maretail	ragweed, giant
bindweed*	ironweed	shepherdspurse
bullnettle	lambsquarters, common	smartweed, Pennsylvania*
bittercress, smallflowered	lettuce, prickly	sowthistle, annual
buttercup, smallflowered	morningglory, annual	speedwell
Carolina geranium	mousetail	thistle, Canada*
cinquefoil, common & rough	mustard, wild	thistle, bull
clover, red*	onion, wild*	velvetleaf
cocklebur, common	pennycress, field	vetch, hairy*
dandelion*	peppergrass*	Virginia copperleaf
dock, curly	plantains	
evening primrose, cutleaf	purslane, common	*Partially controlled
garlic, wild*	ragweed, common	

After applying, plant soybean seed as deep as practical or at least 1-1/2 to 2 inches deep. Adjust the planter press wheel, if necessary, to ensure that planted seed is completely covered.

If desired, this product may be applied preplant to soybeans in tank mixtures with other herbicides such as Poast, Poast Plus, Roundup, Roundup D-Pak, Honcho, Gramoxone Extra, Prowl DG, Prowl 3.3 EC, Pursuit Plus, Scepter 70 DG, Squadron and others that are registered for preplant soybean use.

Compatible crop oil concentrates, agricultural surfactant, and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of 2,4-D on certain weeds and may be added to the spray tank. Read and follow all directions and precautions on this label and on all labels of adjuvants or fertilizers mixed with

this product.

NOTE: Unacceptable injury to soybeans planted in treated fields may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.

Not registered for use in California.

RESTRICTIONS AND LIMITATIONS FOR USE IN SOYBEANS

Do not apply this product prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.

Do not use on low organic sandy soils (less than 1.0%).

Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.

Do not mow or cultivate weeds prior to treating with this product as poor control may result.

Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock. Do not feed or graze treated cover crops to livestock.

Only one application of this product may be made prior to planting soybeans per growing season, regardless of application rate used.

Do not replant fields treated with this product in the same growing season with crops other than those labeled for 2,4-D use.

Sugarcane: See table for recommended rate.

Apply as a pre- or post-emergent spray in the spring after canes emerge and through layby. Consult your local Agricultural Experiment or Extension Service Weed Specialists on specific use of this product to control broadleaved and grass weeds.

Ornamental Turf: Apply 2 to 4 pints on annual broadleaf weeds and 4 pints on biennial and perennial broadleaf weeds. Use enough water to give good coverage. Treat when weeds are young and actively growing. Perennial weeds should be near the bud stage but not flowering at application. Do not use on susceptible Southern grasses such as St. Augustine. Do not apply to newly seeded areas until grass is well established. Bentgrass, clover, legumes, and dichondra may be injured by this treatment. **NOTES FOR ALL TURF SITES (EXCLUDING SOD FARMS):** ~~Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried. The maximum number of broadcast applications per treatment site is 2 per year. When using this product,~~

~~wear long-sleeved shirt, long pants, socks, shoes, and chemical-resistant gloves. Wear eye protection (face shield or safety glasses). It is recommended that safety glasses include front, brow and temple protection. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.~~

Grass Seed Crops: Apply 1 to 4 pints of product per acre in the spring or fall to control broadleaf weeds in grass being grown for seed. Do not apply from early boot to milk stage. Spray seedling grass only after the five leaf stage, using 3/4 to 1 pint per acre to control small seedling weeds. After the grass is well established, higher rates of up to 4 pints per acre can be used to control hard to control annual or perennial weeds. For best results, apply when soil moisture is adequate for good growth. Do not use on Bent unless injury can be tolerated. Do not graze dairy cattle within 7 days of application. Do not apply this product within 30 days of cutting grass for hay. Remove meat animals from treated areas 3 days prior to slaughter.

Fallow Land: On established perennial species such as Canada thistle and Field bindweed, apply up to 3 quarts of product per acre. For annual broadleaf weeds, apply 1 to 2 quarts per acre. For annual broadleaf weeds, apply 1 to 2 quarts per acre. Do not plant any crop for 3 months after treatment or until 2,4-D has disappeared from soil.

Established Pastures and Rangelands: Use 1 to 4 pints of product in sufficient water to give good coverage to one acre depending on type of weeds and stage of growth. Use only on established stands of perennial grasses. Do not use on bentgrass, alfalfa, clover, or other legumes. Do not use from early boot to milk stage where grass seed production is desired. Do not graze dairy cattle within 7 days of application. Do not apply this product within 30 days of cutting grass for hay. Remove meat animals from treated areas 3 days prior to slaughter.

Control of Southern Wild Rose: On rangelands, roadsides, and fencerows, use 1 gallon of product plus 4 to 8 fluid ounces of an agricultural surfactant per 100 gallons of water and spray thoroughly as soon as foliage is well developed. Two or more treatments may be required. On rangeland, apply a maximum of 6 quarts of product per acre per application. Do not graze dairy animals on treated areas within 7 days after application.

General Weed Control: (Airfield, Roadsides, Vacant Lots, Drainage Ditch Banks, Fence Rows, Industrial Sites and similar areas):

Use 1 to 3 quarts of product per acre. Usually 2 quarts per acre will give adequate control. Do not use on herbaceous ground covers or creeping grass such as Bent. Legumes will usually be damaged or killed. Deep-rooted perennials may require repeat applications. Do not use on freshly seeded turf until grass is well established. Delay reseeding for 3 months or until 2,4-D has disappeared from soil.

Rights-of-Way: Apply up to 2 gallons of product per acre for the control of perennial broadleaf weeds and susceptible woody species. For less susceptible

BEST AVAILABLE COPY

perennial broadleaf weeds and difficult to control woody species, use a combination of 2 gallons of product plus 1 to 4 quarts of Garlon® 3A herbicide per acre. For ground application, apply in 20 to 400 gallons of water, depending on the height of the weeds and brush. Use the higher volumes of up to 400 gallons per acre for dense brush 6 feet tall or higher. For aerial application, use 10 to 30 gallons per acre total spray volume.

Woody Plant Control: To control woody plants susceptible to 2,4-D, such as Alder, Buckbrush, Elderberry, Sumac, and Willow on non-crop areas, use 2 to 3 quarts of product per acre in 100 gallons of water. Wet all parts of the plants thoroughly, including stem and foliage, to the point of runoff. Higher volumes of up to 400 gallons per acre are necessary where the brush is very dense and over 6 to 8 feet high. Applications are more effective when made on actively growing plants. Treatment should not be made during time of severe drought or in early fall when leaves lose their green color. Hard to control species may require re-treatment next season.

Tree Injection: For the control of unwanted hardwoods such as elm, oak, hickory, and sweet gum in forest and non-crop areas, apply undiluted product by injecting 1 ml through the bark, using one injection per inch of trunk diameter measured at breast height (4-1/2 feet). For harder to control species (ash, maple, dogwood), use 2 ml of undiluted product per injection. All injections should be as near the root collar as possible and should be evenly spaced around the trunk. Injections may be made at any time of the year but are most effective during the growing season. Maples should not be treated during the spring sap rise.

No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

For Dilute Injection: Mix 1 gallon of product in 19 gallons of water for dilute injections.

AQUATIC APPLICATIONS

Weeds and Brush on Irrigation Canal Ditchbanks: Seventeen Western States: Arizona, California, Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington, Wyoming.

For control of annual and perennial broadleaf weeds, apply 1 to 2 quarts of product per acre in approximately 20 to 100 gallons of total spray. Treat when weeds are young and actively growing before the bud or early bloom stage. For harder to control weeds, a repeat spray may be needed after 3 to 4 weeks for maximum results, using the same rates.

Apply no more than 2 treatments per season. For woody brush and patches of perennial broadleaf weeds, mix one gallon of product in 150 gallons of water. Wet foliage thoroughly, using approximately 1 gallon of spray solution per square rod.

Spraying Instructions: Low pressure (10 to 40 psi) power spray equipment should be used and mounted on a truck, tractor, or boat. Apply while traveling upstream to avoid accidental concentration of chemical into water. Spray when the air is calm, 5 mph or less. Do not use on small canals (less than 10 CFS) where water will be used for drinking purposes.

Boom spraying onto water surface must be held to a minimum and no cross-stream spraying to opposite banks should be permitted. When spraying shoreline weeds, allow no more than 2-foot overspray onto water with an average of less than 1-foot overspray to prevent introduction of greater than negligible amounts of chemical into the water.

Do not allow dairy animals to graze on treated areas for at least 7 days after spraying. Water within treated banks should not be fished.

For Aquatic Weeds in Lakes, Ponds, Drainage Ditches, and Marshes: Use 2-1/2 to 4-1/2 pints of product in 50 to 100 gallons of water per acre. Spray to wet foliage thoroughly. Application should be made when leaves are fully developed, above the water line, and plants are actively growing. Your State Conservation Department or Game and Fish Commission will assist you in determining the best time and rate for application under local conditions.

DO NOT APPLY to more than 1/3 to 1/2 of a lake or pond in any one month because excessive decaying vegetation may deplete oxygen content of water and kill fish.

Do not contaminate water intended for irrigation or domestic purposes except as indicated in directions for use on irrigation ditchbanks.

Perennial and other hard to control weeds may require a repeat application to give adequate control.

WARRANTY LIMITATIONS AND DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the DIRECTIONS FOR USE when used under normal conditions. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL. Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses or damages resulting from the use or handling of

this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify seller, in writing, of any claims to be eligible to receive either remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of seller will be one of the following, at the election of the seller:

1. Refund of purchase price paid by buyer or user for product bought or
2. Replacement of amount of product used.

The seller will not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations and Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations and Disclaimer in any manner.

Garlon® is a registered trademark of Dow Chemical.