

42750-20

11/16/2000

EPA 1/13

ALBAUGH

2,4-D LV 6

LOW VOLATILE HERBICIDE

ACTIVE INGREDIENT:

2-ethylhexyl ester of 2,4-dichlorophenoxyacetic acid\* .....86.5%

OTHER INGREDIENTS \*\*: .....13.5%

TOTAL .....100.0%

\* Equivalent to 57.4% of 2,4-dichlorophenoxyacetic acid or 5.5 lb./gal. Isomer specific by AOAC Method.

\*\*Contains petroleum distillates.

EPA Reg. No. 42750-20

EPA Est. No. 42750-MO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• <u>Immediately call a poison control center or doctor.</u></li> <li>• <u>Do not induce vomiting unless told to do so by a poison control center or doctor.</u></li> <li>• <u>Do not give any liquid to the person</u></li> <li>• <u>Do not give anything by mouth to an unconscious person.</u></li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• <u>Take off contaminated clothing.</u></li> <li>• <u>Rinse skin immediately with plenty of water for 15-20 minutes.</u></li> <li>• <u>Call a poison control center or doctor for treatment advice.</u></li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• <u>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</u></li> <li>• <u>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</u></li> <li>• <u>Call a poison control center or doctor for treatment advice.</u></li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN

May cause chemical pneumonitis if aspirated. If lavage is performed, suggest endotracheal and/or esophagosopic control.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Get medical attention. Do not give anything by mouth to an unconscious person. Avoid alcohol.

**ACCEPTED**

NOV 16 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 42750-20

2/13

~~IF ON SKIN: Wash contaminated areas thoroughly with soap and water. Get medical attention.~~

~~IF IN EYES: Flush with plenty of clean water for 10-15 minutes. Get medical attention.~~

~~NOTE TO ATTENDING PHYSICIAN: May cause chemical pneumonitis if aspirated. If lavage is performed, suggest endotracheal and/or esophagosopic control.~~

See page two for additional precautionary statements.

For chemical spill, leak, fire, or exposure, call CHEMTREC (800) 424-9300.

Manufactured By:  
Albaugh, Inc.  
Ankeny, Iowa 50021

NET CONTENTS  
\_\_\_\_ Gals.  
\_\_\_\_ Liters

Before buying or using this product, read "~~Warranty Limitations and Disclaimer~~" "Conditions of Sale and Warranty" elsewhere on this label. If terms are not acceptable, return **unopened** package at once to seller for full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under the ~~Warranty Limitations and Disclaimer~~ Conditions of Sale and Warranty.

**PRECAUTIONARY STATEMENTS**

**CAUTION**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing.

**PERSONAL PROTECTIVE EQUIPMENT**

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eyewear, and chemical-resistant apron when cleaning equipment, mixing, or loading.

If this container contains over 1 gallon and less than 5 gallons, mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

**ENGINEERING CONTROLS STATEMENTS**

If this container contains 5 gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

3/13

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

### PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

4/13

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

### STORAGE AND DISPOSAL

**STORAGE:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Do not store near heat or open flame. Reclose all partially used containers by thoroughly tightening screw cap. Absorb any spill with a suitable clay absorbent and dispose of as indicated under "Pesticide Disposal."

For safety and prevention of unauthorized use, all pesticides should be stored in locked facilities.

To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification.

Opened, partially used pesticides should be stored in original labeled containers when possible. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of the new container.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

### CONTAINER DISPOSAL

**METAL CONTAINERS:** Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**PLASTIC CONTAINERS:** Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**REFILLABLE CONTAINERS:** If this container has been designated by the supplier as refillable, return empty container to the place of purchase.

### GENERAL INFORMATION

Performance of this product may be affected by local conditions, crop varieties, and application method. User should consult local Extension Service, Agricultural Experiment or University Weed Specialists, and state regulatory agencies for recommendations in your area.

Best results are obtained when product is applied to young succulent weeds that are actively growing. The lower recommended rates will be satisfactory on susceptible annual weeds. For perennial weeds and conditions such as the very dry areas of the western states, where control is difficult, the higher recommended rates should be used.

When product is used for weed control in crops, the growth stage of the crop must be considered.

Some plants and weeds, especially woody varieties, are difficult to control and may require repeat applications.

Application rates should be 1 to 5 gallons of total spray by air or 5 to 25 gallons by ground equipment unless otherwise directed. In either case, use the same amount of 2,4-D recommended per acre. For crop uses, do not mix with oil, surfactants, or other adjuvants unless specifically recommended. To do so may reduce herbicide's selectivity and could result in crop damage.

Aerial applications should be used only when there is no danger of drift to susceptible crops. Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. Although this product is a low volatile formulation, at temperatures above 90°F vapors may damage susceptible crops growing nearby.

Because coarse sprays are less likely to drift than fine, do not use equipment (such as hollow cone small orifice nozzles) or conditions (such as high pressure) that produce such sprays.

Product should not be allowed to come into contact with desirable, susceptible plants such as beans, cotton, fruit trees, grapes, legumes, ornamentals, peas, tomatoes, and other vegetables. Product should not be used in greenhouses. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Users should note that herbicide treatment of public water requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit in your state.

If stored below freezing, efficacy is not affected if product is warmed to 40°F and agitated before using.

Spray equipment used to apply 2,4-D should not be used for any other purpose until thoroughly cleaned.

**Spray Preparation:** Add the recommended amount of product to approximately 1/2 the volume of water to be used for spraying. Agitate well, then add the remainder of the water. Continue agitation during application until spray tank is empty.

**Use in Liquid Nitrogen Fertilizer:** Product may be combined with liquid nitrogen fertilizer suitable for foliar application on corn, grass, pastures, or small grains in one operation. Use product according to directions on this label for those crops. Use liquid nitrogen fertilizer at rates recommended by supplier or Extension Service Specialist. Mix the product and fertilizer according to the following instructions:

Fill the spray tank approximately 1/2 full with the liquid nitrogen fertilizer. Add the product while agitating the tank. Add the remainder of the fertilizer while continuing to agitate. Apply immediately, maintaining agitation during application until tank is empty. DO NOT APPLY DURING COLD (NEAR FREEZING) WEATHER. Spray mixture must be used immediately and may not be stored.

**Note:** (1) If good, continuous agitation is not maintained, separation of the spray mixture and/or clogging of the nozzles is likely to occur.

**Note:** (2) If user's spray program includes frequent application of 2,4-D in liquid fertilizer, consideration should be given to using ~~SEE 2,4-D~~ SOLVE™ 2,4-D which is specially designed and formulated for such use.

### WHERE TO USE

This product is used to control broadleaf weeds in cereal crops, corn, and sorghum; weeds and brush in rangeland, pastures, rights-of-way, and similar noncrop uses.

### WEEDS CONTROLLED

When used properly, product will kill or control the following, in addition to many other noxious plants susceptible to 2,4-D:

Arrowhead	Ironweed
Artichoke	Jimsonweed
Bindweed (hedge)	Lambsquarters
(field and	Locoweed
European)	Mexicanweed
Bitter wintercress	Morningglory
Boxelder	Mustard
Buckhorn	Parrotfeather
Bull thistle	Pennywort
Burdock	Pigweed
Bur ragweed	Plantain
Buttercup	Pokeweed
Catnip	Povertyweed
Chickweed	Puncturevine
Chickory	Purslane
Cocklebur	Russian thistle
Coffebean	Sagebrush
Creeping jenny	Shepherdspurse
Curly indigo	Sowthistle
Dandelion	Stinkweed
Dock	Sumac
Elderberry	Sunflower
Goldenrod	Vetch
Ground ivy	Virginia creeper
Hemp	Wild lettuce
Hoary cress	Wild radish
Honeysuckle	Willow
Indigo	

### LESS SUSCEPTIBLE WEEDS

Kochia	Smartweed
Pigweed (hybrid)	Wild garlic
Poison ivy	Wild onion

### CROPS

**Small grains (barley, oats, wheat, rye), not underseeded with a legume:** See table for recommended use rates. Spray when weeds are small after grain begins tillering but before boot stage (usually 4 to 8 inches tall). Do not apply before the tiller stage nor from early boot through milk stage. To control large weeds that will interfere with harvest or to suppress perennial weeds, preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well.

**Spring Planted Oats:** Use 1/3 pint per acre in sufficient water to give good coverage. Apply after the

7/13

fully tillered stage, except during the boot to dough stage.

**Fall Planted Oats:** Apply 1/6 to 5/6 pints per acre after full tillering but before early boot stage. Some difficult weeds may require the higher rates of 1/2 to 5/6 pints per acre for maximum control, but injury may result. Do not spray during or immediately following cold weather.

**Note:** Oats are less tolerant to 2,4-D than wheat or barley and more likely to be injured. Do not forage or graze treated grain fields within 14 days after treatment with 2,4-D. Do not feed treated straw to livestock.

**Wheat and Barley:** Control of Wild Garlic and Wild Onion. For improved control of difficult weeds including Wild Garlic and Wild Onion, apply 1/3 to 1-1/3 pints of product per acre. Since these rates may injure the crop, do not use unless possible crop damage is acceptable. For the higher rates on spring wheat and barley, consult your local State Agricultural Experiment Station or Extension Service Weed Specialist for recommendations or suggestions to fit local conditions.

**Control of Wild Garlic in Stubble Grain Fields:** Following the harvest of small grains, Wild Garlic often produces new fall growth. This should be sprayed with 1-1/3 to 2 quarts of product per acre. This is a useful practice as one part of a Wild Garlic control program. Do not forage for 14 days following applications. Do not plant any crop for three months after treatment.

**Corn:** See table for recommended use rates.

**Preemergent:** Apply product to emerged weeds from 3 to 5 days after planting but before corn emerges. Do not use on very light, sandy soils. Use the higher rates on heavy soils. Plant corn as deep as practical. Product will not control weeds which have not emerged.

**Postemergent:** Best results are usually obtained when weeds are small and corn is 4 to 18 inches tall. When corn is over 8 inches tall, use drop nozzles to keep spray off corn foliage as much as possible. Do not apply from tasseling to dough stage. If corn is growing rapidly and temperature and soil moisture is high, use 1/3 pint per acre to reduce possibility of crop damage. Delay cultivation for 8 to 10 days to prevent stalk breakage due to temporary brittleness caused by 2,4-D. Application rates of up to 2/3 pint per acre may be used to control some hard to control weeds. However, the possibility of injury to the corn is increased.

Do not use with atrazine, oil or other adjuvants. Since the tolerance to 2,4-D of individual hybrids varies, consult your seed supplier, local Extension Service, Agricultural Experiment Station, or University Weed Specialist for information.

**Preharvest:** After the hard dough or denting stage, apply 2/3 to 1-1/3 pints of product per acre by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as Bindweed, Cocklebur, Dogbane, Jimsonweed, Ragweed, Sunflower, Velvetleaf, and vines that interfere with harvesting. Do not forage or feed corn fodder to livestock for 7 days following application.

**Postharvest:** Following the harvest of corn, Wild Garlic often produces new fall growth. This should be sprayed with 1-1/3 to 2 quarts of product per acre. This is a useful practice as one part of a Wild Garlic control program. Do not forage for 7 days following application. Do not plant any crop for three months after treatment.

**Sorghum (Milo):** See table for recommended rate. Apply to sorghum when crop is 4 to 12 inches high with secondary roots well established. Use drop nozzles when crop is over 10 inches high. Do not apply

8/13

from flowering to dough stage. Rates of up to 2/3 pint per acre may be used to control some hard-to-control weeds. However, the chance of crop injury is increased with the higher rates. Do not use with oil. Use lower rate if conditions of high temperature and high soil moisture exist.

**RECOMMENDED RATES OF PRODUCT PER ACRE\*\***

CROP (SEE DETAILED INSTRUCTIONS ABOVE)	RATE, AVERAGE CONDITIONS	RATE, DRY CONDITIONS AS IN WESTERN STATES*
<b>Small Grains (Wheat, Barley, Rye):</b>		
Annual Weeds	1/3 to 2/3 pt.	2/3 to 1-1/3 pts.
Perennial Weeds	2/3 pt.	5/6 to 1-1/3 pts.
Preharvest	2/3 to 1-1/3 pts.	
<b>Oats:</b>		
Spring	1/3 pt.	
Fall	1/3 to 1/2 pt.	
<b>Corn:</b>		
Preemergent	2/3 to 1-1/3 qts.	
Postemergent	1/3 pt.	1/3 to 1/2 pt.
Preharvest	2/3 to 1-1/3 pts.	
<b>Sorghum (Milo):</b>		
Postemergent	1/3 pt.	1/3 to 1/2 pt.

\*Arizona, Idaho, Montana, Oregon, Utah, Washington, Wyoming.

\*\*If band treatment is used, base the dosage rate on the actual area sprayed.

**Soybeans (Preplant Only) - For Use in Crop Residue Management Systems:** Apply 1/2 to 2/3 pint per acre **not less than 7 days** prior to planting soybeans or 2/3 to 1-1/3 pints per acre **not less than 30 days** prior to planting. Apply to postemergent weeds when small, actively growing, and free of stress caused by extremes in climatic conditions, diseases, or insect damage. The response of individual weed species is variable. Consult your local county agent or state Agricultural Extension Service or crop consultant for advice. Use the higher rate on larger weeds and when perennials are present. (See **WEEDS CONTROLLED** below.)

Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of water per acre in ground equipment.

**WEEDS CONTROLLED**

alfalfa*	horseweed or marestalk	ragweed, giant
bindweed*	ironweed	shepherdspurse
bullnettle	lambquarters, common	smartweed, Pennsylvania*
bittercress, smallflowered	lettuce, prickly	sowthistle, annual
buttercup, smallflowered	morningglory, annual	speedwell
Carolina geranium	mousetail	thistle, Canada*
cinquefoil, common & rough	mustard, wild	thistle, bull
clover, red*	onion, wild*	velvetleaf
cocklebur, common	pennycress, field	vetch, hairy*



9/13

dandelion*	peppergrass*	Virginia copperleaf
dock, curly	plantains	
evening primrose, cutleaf	purslane, common	
garlic, wild*	ragweed, common	*Partially controlled

After applying, plant soybean seed as deep as practical or at least 1-1/2 to 2 inches deep. Adjust the planter press wheel, if necessary, to ensure that planted seed is completely covered.

If desired, this product may be applied preplant to soybeans in tank mixtures with other herbicides such as Poast®, Poast® Plus, Roundup®, Roundup D-Pak®, Honcho®, Gramoxone® Extra, Prowl® DG, Prowl® 3.3 EC, Pursuit® Plus, Scepter® 70 DG, Squadron® and others that are registered for preplant soybean use.

Compatible crop oil concentrates, agricultural surfactant, and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of 2,4-D on certain weeds and may be added to the spray tank. Read and follow all directions and precautions on this label and on all labels of adjuvants or fertilizers mixed with this product.

**Note:** Unacceptable injury to soybeans planted in treated fields may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.

Not registered for use in California.

**Restrictions and Limitations for Use in Soybeans:**

Do not apply this product prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.

Do not use on low organic sandy soils (less than 1.0%).

Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.

Do not mow or cultivate weeds prior to treating with this product as poor control may result.

Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock. Do not feed or graze treated cover crops to livestock.

Only one application of this product may be made prior to planting soybeans per growing season.

Do not replant fields treated with this product in the same growing season with crops other than those labeled for 2,4-D use.

**Ornamental Turf:** Use 2/3 to 2 pints of product in enough water to give good coverage to one acre on established stands of perennial grasses, depending on type of weeds and stage of growth. Do not use on creeping grasses such as Bent except for spot spraying. Newly seeded turf should not be treated until after the second mowing and the lower dosage rate should be used. **Notes for all Turf Sites (excluding Sod Farms):** The maximum number of broadcast applications per treatment site is 2 per year.

**Grass Seed Crops:** Apply 2/3 to 2-2/3 pints of product per acre in the spring or fall to control broadleaf

10/13

weeds in grass being grown for seed. Do not apply from early boot to milk stage. Spray seedling grass only after the five leaf stage, using 1/2 to 2/3 pint per acre to control small seedling weeds. After the grass is well established, higher rates of up to 2-2/3 pints per acre can be used to control hard-to-control annual or perennial weeds. For best results, apply when soil moisture is adequate for good growth. Do not use on Bent unless injury can be tolerated. Do not graze dairy cattle within 7 days of application. Do not apply this product within 30 days of cutting grass for hay. Remove meat animals from treated areas 3 days prior to slaughter.

**Fallow Land:** On established perennial species such as Canada thistle and Field bindweed, apply up to 4 pints of product per acre. For annual broadleaf weeds, apply 1-1/3 to 2-2/3 pints per acre. Do not plant any crop for 3 months after treatment or until 2,4-D has disappeared from the soil.

**Established Pastures and Rangelands:** Use 2/3 to 2-2/3 pints of product in sufficient water to give good coverage to one acre depending on type of weeds and stage of growth. Use only on established stands of perennial grasses. Do not graze dairy cattle within 7 days of application. Do not apply this product within 30 days of cutting grass for hay. Remove meat animals from treated areas 3 days prior to slaughter.

**Wild Garlic and Wild Onion Control:** Apply 2-2/3 to 4 pints of product per acre making three applications, fall-spring-fall or spring-fall-spring, starting in the late fall or early spring. DO NOT graze dairy animals nor cut forage for hay within 7 days of application.

**General Weed Control: (Airfields, Roadsides, Vacant Lots, Fence Rows, Industrial Sites and similar areas):** Use 1-1/3 to 4 pints of product per acre. Usually 2-2/3 pints per acre will give adequate control. Do not use on herbaceous ground covers or creeping grass such as Bent. Legumes will usually be damaged or killed. Deep-rooted perennials may require repeat applications. Do not use on freshly seeded turf until grass is well established. Delay reseeding for 30 days.

**Woody Plant Control:** To control woody plants susceptible to 2,4-D, such as Alder, Buckbrush, Elderberry, Sumac, and Willow on non-crop areas, use 1-1/3 to 2 quarts of product per acre in 100 gallons of water. Wet all parts of the plants thoroughly, including stem and foliage, to the point of runoff. Higher volumes of up to 400 gallons per acre are necessary where the brush is very dense and over 6 to 8 feet high. Applications are more effective when made on actively growing plants. Treatment should not be made during time of severe drought or in early fall when leaves lose their green color. Hard-to-control species may require re-treatment next season.

#### **USES IN FOREST MANAGEMENT:**

**Conifer Release:** For control of alder, apply 2/3 to 2 quarts of product per acre in 8 to 25 gallons of water, and apply as a foliage spray. Treat when 3/4 of the brush foliage has attained full size leaves and before new conifer growth reaches 2" in length. This is usually between early May and mid-June. Adjust treatment date depending on stage of growth and brush species. This may cause leader deformation on exposed firs, but they should overcome this during the second year after spraying.

To control susceptible brush species such as Ceanothus spp., Chinquapin, Madrone, Manzanita, Oak and Tanoak and to release Douglas fir, Hemlock, Sitka spruce or Grand fir, apply up to 2 quarts per acre before new growth on Douglas fir is 2" long. To control Manzanita and Ceanothus in Ponderosa pine, apply up to 2 2/3 quarts per acre before pine growth begins in spring. To increase performance, add 2 to 4 quarts of diesel, fuel oil, kerosene, or a suitable approved agricultural surfactant at recommended label rate.

After Northern conifers, Jack pine, Red pine, Black spruce, and White spruce cease growth and "harden

off" in late summer, a spray of 1 to 2 quarts of product in 8 to 25 gallons of water per acre may be applied by air to control certain competing hardwood species such as Alder, Aspen, Birch, Hazel and Willow. Since this treatment may cause occasional conifer injury, do not use if such injury cannot be tolerated. Consult your regional or extension forester or state herbicide specialist for recommendations to fit local conditions.

**Tree Injections (Pine Release):** To control hardwoods, such as Oaks, Hickory, Maple, Pecan, Elm, Sumac, Sweetgum and Hawthorn in forest and other noncrop areas. apply undiluted product in a concentrate tree injector calibrated to apply .7 ml. per injection. Space injections 2" apart, edge to edge, completely around the tree and close to the base. The injector bit must penetrate the inner bark. On hard-to-kill species such as Hickory, Dogwood, Red maple, Blue beech and Ash, make injections 1 to 1-1/2 inches apart, edge to edge. Treatment may be made at any time of the year. For best results, injections should be made during growing season, May 15-October 15. For dilute injections, mix 2/3 gallon of product in 19 gallons of water.

**Dormant Application (other than pine):** For the control of susceptible deciduous brush species such as Alder, Cascara, Cherry, Poplar and Service berry, apply up to 2 quarts of product per acre in sufficient diesel, fuel oil or kerosene for good coverage. Application may be made by ground or air and should be made before conifer bud break.

**Pine Only:** Make application while pine buds are still dormant. Apply 1-1/3 quarts of product per acre in sufficient water for good coverage by air or ground equipment. Do not use this application unless some pine injury is acceptable. Use of diesel, kerosene, or other oil, or addition of surfactants to spray mix may cause unacceptable pine injury.

**Herbaceous Weed Control:** To control over-wintering susceptible weeds such as False dandelion, Klamath weed, Plantain, and Tansy ragwort, apply 2/3 to 2 quarts of product per acre in sufficient water for good coverage. Make application at rates and timing indicated above if pines are present. For control of hazel brush and similar species in the Lake States area, apply 1-1/3 quarts of product per acre in 8 to 25 gallons of water, when new shoot growth of Hazel is complete.

**Site Preparation: (As Dormant Spray)** - For control of Alder prior to planting seedlings, apply 1-1/3 to 2-2/3 quarts of product per acre in diesel, fuel oil, or similar oil before foliage is 1/4 full size. Application may be made by air or ground.

**(As Foliage Spray)** - For control of Alder prior to planting seedlings, apply 1-1/3 to 2-2/3 quarts of product per acre in 8 to 25 gallons of water, after most Alder leaves are full size. To increase penetration, 2 to 4 quarts per acre of diesel, fuel oil, kerosene, or a suitable approved agriculture surfactant at recommended label rates may be added to the spray mixture.

#### CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, INC. or the Seller. All such risks shall be assumed by the Buyer.

ALBAUGH, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. ALBAUGH, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. THIS WARRANTY DOES NOT EXTEND TO.

AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. IN NO CASE SHALL ALBAUGH, INC. OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALBAUGH, INC. and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of ALBAUGH, INC. No employee or agent of ALBAUGH, INC. or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

**WARRANTY LIMITATIONS AND DISCLAIMER**

~~Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the DIRECTIONS FOR USE when used under normal conditions. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL. Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller. When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify seller, in writing, of any claims to be eligible to receive either remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of seller will be one of the following, at the election of the seller:~~

- ~~1. Refund of purchase price paid by buyer or user for product bought or~~
- ~~2. Replacement of amount of product used.~~

~~The seller will not be liable for consequential or incidental damages or losses.~~

~~The terms of this Warranty Limitations and Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations and Disclaimer in any manner.~~

Roundup®, Roundup D-Pak®, and Honcho® are registered trademarks of Monsanto Chemical Company.

Gramoxone® is a registered trademark of Zeneca Agrochemicals.

Poast®, Prowl®, Squadron®, Scepter®, and Pursuit® are registered trademarks of BASF Corp.

Garlon® is a registered trademark of DowElanco.

SOLVE™ is a trademark of Albaugh, Inc.