



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

June 18, 2024

Carolyn Miter
Registration Specialist
Albaugh LLC
PO Box 2127
Valdosta, GA 31604

Subject: Label Amendment – Updated the Spray Drift language, updated the Weed Resistance language, revised directions for use, and fixed grammatical errors.
Product Name: ALBAUGH TRIFLURALIN 10G
EPA Registration Number: 42750-34
Application Date: 10/15/20
Case Number: 472306

Dear Carolyn Miter:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find

or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Eric Ingram at 202-566-0061 or at ingram.eric@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'Kable Bo Davis', enclosed within a hand-drawn oval.

Kable Bo Davis
Senior Regulatory Specialist
Office of Pesticide Programs
Registration Division, Immediate Office

Enclosure

ALBAUGH TRIFLURALIN 10G

A Selective Herbicide for Preemergence Control
of Annual Grasses and Broadleaf Weeds

ACTIVE INGREDIENT:

Trifluralin: a,a,a-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine10%

OTHER INGREDIENTS:90%

TOTAL:100%


KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	
See side panel for additional Precautionary Statements	

EPA Reg. No. 42750-34

EPA Est. No. xxxxxx-xx-xxx

NET CONTENTS: ____ lbs.



MANUFACTURED FOR:
Albaugh, LLC
1525 NE 36th Street
Ankeny, Iowa 50021

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Chemical resistant gloves Category A, such as butyl rubber > 14 mils, or natural rubber > 14 mils, or neoprene rubber > 14 mils or nitrile rubber > 14 mils,
- Shoes plus socks, and
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

This product is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. **DO NOT** apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This product must be soil-incorporated within 24 hours of application either by mechanical incorporation or at least 0.5 acre inch of water (rainfall, irrigation). If a second incorporation is needed it should be delayed 3 to 5 days after the first incorporation and be completed prior to planting.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of **12 hours**.

EXCEPTION: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

1. Coveralls,
2. Chemical resistant gloves Category A, such as butyl rubber > 14 mils, or natural rubber > 14 mils, or neoprene rubber > 14 mils or nitrile rubber > 14 mils,
3. Shoes plus socks, and
4. Protective eyewear.

Mandatory Spray Drift Management

Aerial Applications

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a coarser or coarser droplet size (ASABE S572.1).
- If the windspeed is 10 miles per hour or less, applicators must use $\frac{1}{2}$ swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use $\frac{3}{4}$ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site. If this windspeed is greater than 10 mph the boom length must be 65% or less of the wingspan for fixed wing aircraft at 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a coarse or coarser droplet size (ASABE S572.1)
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift. When applying aerially to crops, **DO NOT** release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

SHEILDED SPRAYERS

Shielding the boom of individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

PRODUCT INFORMATION

This product is a selective preemergence herbicide for control of many annual grasses and broadleaf weeds. The product must be mechanically incorporated within 24 hours after application to effectively control weeds. This product controls susceptible weeds as they germinate. It does not control established weeds.

GENERAL USE PRECAUTIONS

Applied according to directions and under normal growing conditions this product will not harm the treated crop. Overapplication may result in crop injury or rotational crop damage from soil residue. Uneven application or improper incorporation can result in erratic weed control or crop injury. Seedling disease, cold weather, deep planting, excessive moisture, high salt concentration or drought may weaken crop seedlings and increase the possibility of damage. Under these conditions, delayed crop development or reduced yields may result.

DO NOT apply this product to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.

DO NOT use this product on any crop grown in Pecos County or Reeves County, Texas.

In Montana, uses of this product are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

WEED RESISTANCE MANAGEMENT

For resistance management, please note that this product is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to this product and Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

Best Management Practices

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - (2) a spreading patch of non-controlled plants of a particular weed species;
 - (3) surviving plants mixed with controlled individuals of the same species.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Gowan Company representative.

ROTATION CROP RESTRICTIONS

Sugarbeets, Redbeets and Spinach:

In Arizona, Colorado, California, Idaho, Nevada, New Mexico, Oregon, Utah, Washington and Wyoming:

Sugarbeets, redbeets or spinach should not be planted for 12 months after a spring application or 14 months after a fall application. Plowing to a depth of 12 inches prior to planting these crops will reduce the possibility of crop injury. If land has not been irrigated, these crops should not be planted for 18 months after a spring application or 20 months after a fall application of this product.

In all other areas: Sugarbeets, redbeets, and spinach should not be planted for 12 months after a spring application or 14 months after a fall application. Before planting sugarbeets, moldboard plow to a depth of 12 inches to reduce the possibility of crop injury.

Proso Millet, Corn, Sorghum (Milo), Oats and Annual or Perennial Crops or Grass Mixtures:

In Arizona, Colorado, California, Idaho, Nevada, New Mexico, Oregon, Utah, Washington and Wyoming: Proso millet, corn, sorghum (milo), oats, and annual or perennial grass crops or grass mixtures should not be planted for 12 months after a spring application or 14 months after a fall application of this product to avoid the possibility of crop injury. If land has not been irrigated, these crops should not be planted for 18 months after a spring application or 20 months after a fall application. Moldboard plowing to a depth of 12 inches before planting these crops will reduce the possibility of crop injury.

In Minnesota, North Dakota and South Dakota: Proso millet, sorghum (milo), oats and annual or perennial grass crops or grass mixtures should not be planted for 18 months after a spring application or 21 months after a fall application of this product.

In those portions of Kansas, Nebraska, Oklahoma and Texas that receive less than 20 inches of rainfall and irrigation to produce a crop: **DO NOT** plant proso millet, sorghum (milo), oats and annual or perennial grass crops or grass mixtures for 18 months after an application of this product. In sorghum, cool wet weather conditions during early growth stages may increase the possibility of crop injury. In areas receiving more than 20 inches of rainfall and irrigation, these crops should not be planted for 12 months after a spring application or 14 months after a fall application of this product.

Other Crops:

Vegetable crops other than those listed on this label for use with preplant soil incorporated application of this product should not be planted within 5 months after an application of this product.

APPLICATION DIRECTIONS

Soil Preparation

This product may be applied to standing stubble or soil that has been pre-tilled. Existing weeds and crop residues should be reduced to a manageable level using tillage so that this product can be uniformly incorporated into the top 2 to 3 inches of the final seedbed. Soil surface conditions should allow this product to be thoroughly and uniformly mixed into the top 2 to 3 inches of soil. If this is not possible, the soil should be tilled prior to application.

General Soil Conditions

The soil surface should be smooth enough to allow for uniform application and efficient incorporation of this product. Apply when soil moisture is sufficient to allow breakup of large clods and uniform mixing during the incorporation process. Soil compaction and/or nonuniform incorporation may occur when soil is excessively moist.

Application

This product may be applied with ground or aerial broadcast application equipment. Apply only with equipment capable of accurate calibration and uniform application of herbicide granules. Apply at the recommended rate for soil texture class to be treated. Follow calibration directions provided by the equipment manufacturer. Avoid releasing granules in narrow bands on the soil surface as this may cause crop injury.

Freezing will not adversely affect this product. If product is frozen at the time of application, agitate or thaw to restore free-flowing properties to this product.

Soil Texture Guide for Application Rates

This product rate recommendations for incorporated treatments are based on "Soil Texture Class" (coarse, medium or fine) and soil organic matter content. A fine textured soil (e.g., clay loam) will require a higher application rate than a coarse textured soil (e.g., loamy sand). In the table below, find the "Soil Texture Class" (coarse, medium or fine) corresponding to the soil texture to be treated. Choose the proper rate for each application based on the "Soil Texture Class" and specific crop recommendations. **DO NOT** exceed recommended rates.

SOIL TEXTURE CLASS	SOIL TEXTURE TO BE TREATED
Coarse (Light) Soils	Sand, loamy sand, sandy loam
Medium Soils	Loam, silty clay loam†, silt loam, silt, sandy clay loam†
Fine (Heavy) Soils	Clay, clay loam, silty clay loam†, silty clay, sandy clay, sandy clay loam†

†Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine textured soils. If silty clay loam or sandy clay loam soils are predominantly sand or silt, they are usually classified as medium textured soils. If they are predominantly clay, they are usually classified as fine textured soils.

Application Rate Ranges

Where a rate range is shown, use the lower rate for coarser textured soils or soils with low organic matter content. Use the higher rate in the rate range for finer textured soils and on soils containing more than 5% organic matter. Where soil texture is variable within the same field, use the lower recommended this product rate.

APPLICATION TIMING

Spring Application:

Apply this product any time after January 1 when soil can be worked and is in condition suitable for good incorporation. See "Approved Crops" section for recommendations on specific crops.

Fall Application:

Application Timing: In California, Minnesota, North Dakota and South Dakota, apply this product any time between September 1 and December 31. In all other states, fall apply this product any time between October 15 and December 31.

Application Rates for Fall Application:

Refer to the "Approved Crops" section of this label for specific rate recommendations. Increased rates for fall application are recommended for certain crops grown in certain geographic areas. For crops for which there are no specific fall application instructions, and for which this product is recommended as a preplant incorporated treatment, use the rates listed for spring applications. In areas receiving greater than 20" total average annual rainfall and irrigation, use the higher rate in the recommended rate range. **DO NOT** apply this product prior to planting sugar beets, potatoes and direct seeded tomatoes the following spring.

Ground may be left flat or bedded-up over winter. On bedded ground, knock beds down to desired height before planting, moving some treated soil from beds into furrows. Where soil is left flat over winter, be careful not to turn up untreated soil during spring bedding operations. Destroy established weeds during seedbed preparation.

Prior to planting, destroy any weeds which have become established in furrows due to uncovering of untreated soil. **DO NOT** apply this product in the fall to soils that are wet or subject to prolonged periods of flooding, or where rice was grown the previous year.

Incorporation Directions:

This product requires two incorporation passes. The first should occur within 24 hours after application. For best weed control results, the second incorporation should be delayed 3 to 5 days after the first and be completed prior to planting. A minimum delay of 7 days after the first incorporation is recommended for certain this product uses in small grains. See specific use directions. The second incorporation pass should occur in a different direction than the first. Use incorporation equipment capable of thoroughly and uniformly mixing this product into the top 2 to 3 inches of the final seedbed. Erratic weed control may result if untreated soil is moved to the surface during the second incorporation pass. To avoid this problem, set equipment so that the second incorporation pass is not deeper than the first.

Incorporation to Established Crops:

This product may be applied and incorporated in certain established crops. Refer to the "Approved Crops" section of this label for specific incorporation directions.

Incorporation in Bedded Culture:

For optimum weed control, this product should be incorporated into the top 2 to 3 inches of the final seedbed.

Application Prior to Bedding:

Apply and make first incorporation with recommended equipment. The bedding operation serves as the second incorporation. **DO NOT** expose untreated soil during post-bedding operations†.

Application After Bedding:

Knock off beds to planting height before applying. Apply this product and incorporate with recommended equipment that will conform to the bed shape. **DO NOT** leave untreated soil exposed.†

†Avoid removal of treated soil from the seedbed before or during the planting operation. Exposure of untreated soil, will allow weeds to germinate in the drill row.

Cultivation After Planting:

Areas treated with this product may be shallowly cultivated without loss of weed control activity. Avoid deep cultivation that could bring untreated soil to the soil surface and result in loss of weed control.

RECOMMENDED INCORPORATION EQUIPMENT

Any recommended incorporation implement may be used alone or in combination with any other recommended implement. Two incorporation passes are necessary unless otherwise specified. The second incorporation should not be deeper than the first.

Tandem Disc:

Set to cut 4 to 6 inches deep and operate at 4 to 6 mph.

Field Cultivator:

Set equipment to cut 3 to 4 inches deep and operate at 5 or more mph. A field cultivator is defined as an implement with 3 to 4 rows of sweeps, spaced at intervals of 7 inches or less and staggered so that no soil is left unturned. Chisel points should not be used.

Chisel Plow: (For use in northern Great Plains)

The chisel plow may be used for the first incorporation pass only. Any other recommended incorporation implement may be used for the second pass for row crops. The chisel plow may be used for any tillage or incorporation pass in the Summer Fallow program. Operate chisel plow 4 to 5 inches deep at 4 to 6 mph. A chisel plow is defined as having 3 rows of 14 to 18-inch sweeps spaced no more than 12 inches apart. Sweeps should be staggered so that no soil is left unturned. Chisel points should not be used.

Combination Seedbed Conditioners:

These implements are defined as three or more tillage devices combined to operate as a single tillage unit. For example, 2 to 3 rows of field cultivator C- or S-shaped shanks with an effective sweep spacing of 6 to 9 inches (staggered so that no soil is left unturned), followed by a spike-tooth or flexline harrow, followed by a ground driven reel or basket. Combination implements should be set to cut 3 to 4 inches deep and operated at a minimum speed of 6 mph. This product can be incorporated with one pass when using a combination seedbed conditioner. Two incorporations are required under conditions which prevent optimum soil mixing such as excessive trash, roughness, high clay content or soil moisture.

Rolling Cultivator:

Set to cut 2 to 4 inches deep and operate at 6 to 8 mph. Rolling cultivators are adequate for use on coarse and medium textured soils only.

Mulch Treader (other similar disc-type implements):

Set to cut 3-4 inches deep and operate at 5 to 8 mph.

P.T.O. Driven Equipment (tillers, cultivators, hoes):

Adjust to incorporate this product into the top 2 to 3 inches of the seedbed with rotors spaced to provide a clean sweep of the soil. Only one incorporation is necessary. P.T.O. driven equipment should not be operated at a speed greater than 4 mph.

Other Equipment:

Other implements including a flexible tine-tooth harrow (Flextine or Melroe), sweep-type cultivator or rolling cultivator are recommended, but only for certain uses defined in the "Approved Crops" section of this label.

Weeds Controlled by this product

The following weeds are controlled by preplant incorporated treatments of this product.

GRASSES

Common Name

annual bluegrass
barnyardgrass (watergrass)
bromegrass (cheatgrass, downy brome)
crabgrass (large crabgrass, smooth crabgrass)
Foxtail (bottlegrass, bristleglass, giant foxtail, green
foxtail, foxtail millet, pigeongrass, robust foxtail, yellow foxtail)
johnsongrass (from seed)
junglerice
ryegrass, annual
sandbur (burgrass)
shattercane (wild cane)
signalgrass, broadleaf (brachiaria)
sprangletop
stinkgrass (lovegrass)
oat, wild†

Scientific Name

Poa annua
Echinochloa crus-galli
Bromus tectorum
Digitaria spp.
Setaria spp.

Sorghum halepense
Echinochloa colonum
Lolium multiflorum
Cenchrus incertus
Sorghum bicolor
Brachiaria platyphylla
Leptochloa filiformis
Eragrostis cilianensis
Avena fatua

BROADLEAF WEEDS

Common Name

carpetweed
chickweed, common
goosefoot
henbit
knotweed
lambsquarters, common
pigweed (carelessweed, prostrate pigweed,
redroot pigweed, rough pigweed, spiny
pigweed)
puncturevine (western U.S. only) (caltrop, goathead)
purslane, common
pusley, Florida (Florida purslane, Mexican clover, pusley)
stinging nettle (nettle)

Scientific Name

Mollugo verticillata
Stellaria media
Chenopodium hybridum
Lamium amplexicaule
Polygonum aviculare
Chenopodium album
Amaranthus spp.

Tribulus terrestris
Portulaca oleracea
Richardia scabra
Urtica dioica

†This product applied as a preplant incorporated (PPI) treatment will provide partial control of wild oats. This claim is for all PPI uses except for fall application for spring seeded cereals at foxtail (pigeongrass) control rates.

CROP SPECIFIC USE INSTRUCTIONS

ALFALFA, ESTABLISHED

Apply this product to established alfalfa prior to weed emergence with ground or aerial equipment. A single rainfall or overhead sprinkler irrigation of 0.5 inches or more, flood irrigation or furrow irrigation after application is required to activate this product. If activated using furrow irrigation, the surface of beds between furrows should be thoroughly wetted. If activation does not occur within 3 days after application this product should be activated using incorporation equipment that will ensure thorough soil mixing with minimum damage to the established alfalfa.

Weeds Controlled:

Barleygrass, bromegrass (cheatgrass, downy brome), canarygrass, cheat (chess), crabgrass, foxtail, junglerice, sandbur, and wild barley.

Application Rate per Acre:

Apply this product at a rate of 20 lbs./acre (2.0 lbs. a.i./acre) per application with a maximum of 4 lbs. a.i./acre per year for all soil textures.

Application Timing:

This product may be applied during dormancy or throughout the growing season immediately after a cutting. Because this product does not control established weeds, application must be made prior to the expected time of weed germination.

Fall Application:

Apply immediately after a cutting between August 1 and October 1. When fall applied, this product controls bromegrass and cheat in addition to other weeds listed above that germinate after application. Bromegrass and cheat begin to germinate in the fall with the onset of cooler weather.

Precautions:

In the season following a 20 lb./acre this product treatment, where established alfalfa is to be rotated to another crop, plant only those crops for which this product can be applied as a preplant incorporated treatment or crop injury may result. **DO NOT** apply more than 2.0 lbs. a.i. per application. **DO NOT** harvest alfalfa forage within 21 days of application and **DO NOT** harvest alfalfa hay within 20 days of application of this product.

ASPARAGUS, ESTABLISHED

Apply this product to established asparagus as a single or split application. This product will suppress volunteer seedling asparagus and field bindweed when recommendations for rates, application and timing are followed. Follow recommended soil preparation, application and incorporation procedures for this product.

Application Timing:

Apply in winter or early spring after mature ferns have been removed, but before new spears begin to emerge. Apply post-harvest applications immediately after harvest in late spring or early summer just before ferns are allowed to develop.

Broadcast Application Rates/Acre

Soil Texture	Split Application Before and After Harvest	Split Application Before and After Harvest
Coarse	5.0 + 5.0 lbs.	10 lbs.
Medium	7.5 + 7.5 lbs.	15 lbs.
Fine	10 + 10 lbs.	20 lbs.

• **DO NOT** apply more than 10 lbs. per acre on coarse soils, 15 lbs. per acre on medium soils, or 20 lbs. per acre on fine soils during the calendar year.

BEANS, DRY

Apply and incorporate this product in the spring before planting or in the fall. See instructions for fall application of this product under the heading "Application Timing" in the "Product Information" section of this label.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

BEANS, GUAR AND MUNGBEAN

Apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.
Fine	7.5 lbs.

- All soils with 2-5% organic matter--7.5 lbs.

BEANS, LIMA AND SNAP

Apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	5 lbs.
Fine	7.5 lbs.

- All soils with 2-5% organic matter--7.5 lbs.

CARROT

Apply and soil incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

CASTOR BEAN

Apply and soil incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.

- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

CELERY

Apply as a soil incorporated treatment. this product may be applied to direct seeded or transplant celery before planting, at planting or immediately after planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

CHICORY/ENDIVE

Apply and soil incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.
Fine	10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs. to 12.5 lbs.

COLE CROPS (BROCCOLI, BRUSSELS SPROUTS, CABBAGE, AND CAULIFLOWER)

Direct Seeded Cole Crops--Apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	5 lbs.
Fine	7.5 lbs.

- All soils with 2-5% organic matter--7.5 lbs.

Precautions:

Direct seeded cole crops exhibit marginal tolerance to higher than recommended rates of this product. Stunting or reduced stands may occur.

Transplanted Cole Crops--Apply and incorporate this product before transplanting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

CORN--FIELD CORN ONLY

This product may be applied as a postemergence treatment following the use of a preemergence herbicide. Uniformly apply to the soil surface when crop is well established (2 true leaf stage or taller), or immediately after a cultivation, up to a height of 30 inches. This product does not control established weeds.

Incorporation Directions:

In corn, this product should be incorporated within 24 hours after application. Incorporation may be accomplished with one pass of a sweep-type cultivator or properly adjusted rolling cultivator. The sweep-type cultivator should have 3 to 5 sweeps per row middle and be operated at a speed that will provide maximum soil movement. Set middle sweeps so as to avoid exposing untreated soil. Adjust incorporation equipment so as to avoid mechanical injury to the crop.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	3.75-5 lbs.†
Medium	5-7.5 lbs.
Fine	7.5-10 lbs.

†Apply 5 to 7.5 lbs. per acre on coarse soils in Alabama, Florida, Georgia, North Carolina, South Carolina, and Virginia to control fall panicum and Texas panicum.

- Apply lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Precautions

- **DO NOT** apply this product to sweet corn or corn grown for seed.
- **DO NOT** apply this product as a preplant or preemergence treatment or crop injury may occur.

COTTON

Apply and incorporate this product before or at planting, immediately after planting (before crop emergence), up to and at lay-by and in the fall. When incorporating this product after planting (postplant), be careful not to disturb the seed.

Broadcast Application Rates/Acre

Fall Application

Soil Texture	Spring† Application	Eastern U.S.††	Western U.S.†††
Coarse	5 lbs.	10 lbs.	7.5 lbs.
Medium	6.25-7.5 lbs.	10 lbs.	10 lbs.
Fine	7.5-10 lbs.	12.5 lbs.	12.5 lbs.

†Spring Application:

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

††Fall Application:

Use rates for eastern cotton producing areas including: Alabama, Arkansas, northern Florida, Georgia, Louisiana, Mississippi, southeastern Missouri (Bootheel), North Carolina, New Mexico, Oklahoma, South Carolina, Tennessee, and Texas.

†††Fall Application:

Use rates for western cotton producing areas including: Arizona, California and Nevada.

For cotton grown in areas other than those listed above, fall apply this product at broadcast rates recommended for areas receiving greater than 20 inches of annual rainfall and irrigation.

Restriction of Use:

DO NOT apply within 90 days of harvest. **DO NOT** apply more than 1.25 lb./A a.i. per application and **DO NOT** apply more than 2.0 lbs. a.i. per crop year (either fall application thru lay-by application or pre-plant plus post-plant thru lay-by).

Lay-by Treatment:

Apply and incorporate this product any time up to lay-by, but not less than 90 days before harvest. Apply so as to achieve a uniform distribution of granules on the soil surface beneath cotton plants. Use the same rates as for preplant incorporated treatments. Soil incorporate using one pass of a sweep-type cultivator or properly adjusted rolling cultivator. Operate cultivation equipment at speeds sufficient to provide vigorous soil mixing and exercise care to avoid mechanical injury to the crop.

Preemergence Overlay Applications Following this product:

Apply this product as a preplant incorporated treatment. Additional weeds tolerant to this product may be controlled using overlay applications of Cotoran, Karmex, Zorial or other products registered for use on cotton. Such applications may be made unless use following a this product application is specifically prohibited by the product label. Consult the overlay product label for additional weeds controlled, directions for use, cautions and limitations before use.

Postemergence Applications Following this product:

Apply this product as a preplant incorporated treatment. Additional weeds tolerant to this product may be controlled using postemergence applications of products registered for use on cotton. Such treatments may be made unless use following a this product application is specifically prohibited by the product label. Consult the postemergence product label for additional weeds controlled, directions for use, cautions and limitations before use.

COTTONWOOD TREES GROWN FOR PULP

New Plantings: Apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- All soils with 2-5% organic matter--7.5 to 10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Established Plantings:

Apply this product in established plantings as a ground application. A single rainfall or overhead sprinkler irrigation of 0.5 inches or more or flood irrigation after application is required to activate this product.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
All soils	10-20 lbs.

Application rate range may be adjusted according to weed pressure.

Johnsongrass Suppression:

Proper soil preparation before application is necessary for satisfactory results. Use incorporation methods not injurious to the crop.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
All soils	20 lbs.

Some johnsongrass plants will escape. Timely cultivation with tillage implements or spot spraying with effective postemergence herbicides will improve the level of johnsongrass control.

CROP STUBBLE OR CROP RESIDUE

Fall or Spring Application to Ground with Standing Crop Stubble or Crop Residue for Weed Control in Following Crop: This product may be applied in the fall or spring to ground with standing crop stubble or crop residue for weed control in the following crop. When applied in this manner, the first incorporation must occur within 72 hours after application. The second incorporation must be delayed a minimum of 5 days after the first and may occur any time up to planting. This application is limited to use on ground that has not had a primary tillage treatment following crop harvest. Following application, plant only crops on which This product may be applied as a preplant incorporated treatment.

Refer to the use directions for specific crops, including application rates and other use information. Apply at the recommended broadcast application rate for the crop, soil texture, and target weed species. When fall applied, use the highest rate in the rate range allowed for the soil texture to be treated.

Incorporation Directions: Incorporate once using minimum tillage equipment such as the rotary harrow, field cultivator, under cutter chisel plow or other suitable tillage implement. It is desirable to loosen and mix surface soil sufficiently to trap this product, but maintain a surface cover of stubble and crop residue sufficient for erosion protection. The second incorporation may occur any time prior to or in conjunction with tillage operations associated with planting.

CUCURBITS (CANTALOUPE, CUCUMBER AND WATERMELON)

Postemergence Application Only:

Apply and incorporate this product when plants have reached the 3 to 4 true leaf stage of growth. Set incorporation equipment to move treated soil around the base of plants during incorporation.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

FLAX AND FLAX SEED (Fall Application Only)

Apply and incorporate this product in the fall between September 1 to December 31 in California, Minnesota, North Dakota, and South Dakota, and between October 15 and December 31 in other states. Refer to instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.
Fine	10 lbs.

Special Instructions for Flax

1. Incorporation operations or other tillage performed in the spring prior to seeding should be relatively shallow so as to maintain a firm seedbed. The seedbed should be packed just prior to seeding.
2. Seeding should be done with a press drill or hoe drill. Seed into moist seedbed no more than 1.5 inches deep.
3. Flax should not be seeded until the seedbed has warmed up.

GRAIN SORGHUM (MILO)

This product may be applied as a postemergence treatment following the use of a preemergence herbicide. Uniformly apply to the soil surface when grain sorghum is 8 inches tall or taller. This product does not control established weeds.

Soil Preparation:

Cultivate before application to remove established weeds and cover the base of plants with soil. Set cultivation equipment to add approximately one inch of soil to the base of sorghum plants.

Activation and Incorporation Directions:

In grain sorghum, this product should be incorporated within 24 hours after application. Incorporation may be accomplished with one pass of a sweep-type cultivator or properly adjusted rolling cultivator. The sweep-type cultivator should have 3 to 5 sweeps per row middle and be operated at a speed sufficient to provide vigorous soil mixing. Set middle sweeps so as to avoid exposing untreated soil. Adjust incorporation equipment so as to avoid mechanical injury to the crop.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	4-5 lbs.
Medium	5-7.5 lbs.
Fine	7.5-10 lbs.

- Apply lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

GREENS (TURNIP GREENS GROWN FOR PROCESSING, COLLARD, KALE AND MUSTARD GREENS)

Apply to greens as a preplant incorporated treatment.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.
Fine	7.5 lbs.

HOPS

Apply and incorporate this product to established hops during dormancy. Incorporate once using incorporation equipment that will ensure thorough soil mixing with minimal damage to crop stand.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	6.25-7.5 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.

MUSTARD: GROWN FOR SEED OR PROCESSING FOR FOOD IN MINNESOTA, NORTH DAKOTA AND SOUTH DAKOTA

Apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.
Fine	7.5 lbs.

OKRA

Apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

ONION GROWN FOR DRY BULBS ONLY

A single rainfall or overhead sprinkler irrigation of 0.5 inches or more or flood irrigation after application is required to activate this product. **DO NOT** apply within 60 days of harvest.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	3.75-5 lbs.
Medium	5-6.25 lbs.

- Use the lower rates where light weed pressure is anticipated.

Soil incorporation may be accomplished by operating sweep-type or rolling cultivators 2 to 4 inches deep at 6 to 8 mph. Incorporation equipment must mix this product uniformly in the soil. Incorporate with two passes. The first pass must be within 24 hours of application or erratic weed control may result. Avoid covering exposed onion bulbs with treated soil during incorporation as injury to the crop may occur. Care should be taken to avoid injury to the roots during incorporation.

Precautions:

Applied according to these directions and under normal growing conditions, this product will not adversely affect onions. Diseases, improper incorporation depth, excessive moisture, high salt concentration or drought may weaken the crop and increase the possibility of damage from this product. Under these conditions, delayed crop development or reduced yields may result.

PEAS (DRY PEAS AND ENGLISH PEAS)

Apply and incorporate this product in the spring before planting or in the fall. Refer to instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre

Soil Texture	Spring Application	Fall† Application
Coarse	5 lbs.	5 lbs.
Medium	5 lbs.	6.25-7.5 lbs.
Fine	7.5 lbs.	7.5 lbs.

†This product may be fall applied to Dry and English Peas in the states of Idaho, Oregon, and Washington.

PEAS (SOUTHERN PEAS)

Apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.

Fine	7.5-10 lbs.
------	-------------

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

PEANUTS (SPANISH PEANUTS, FLORUNNER, AND FLORIGIANT VARIETIES)

For Use in Texas, Oklahoma, and New Mexico

Apply and incorporate this product before planting, at planting or immediately after planting. When incorporating after planting, take care not to disturb the seed.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.

PEPPER (Transplant Only)

Apply and incorporate this product prior to transplanting. **DO NOT** apply after transplanting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

POTATOES (Not for Use in Maine)

Apply and incorporate this product after planting prior to crop emergence, immediately following drag off, or after potato plants have fully emerged.

Incorporation:

Set incorporation equipment so that the bed and furrow are uniformly covered with a layer of treated soil. If the layer of treated soil is not uniform and the herbicide is concentrated over the bed, potato emergence may be retarded and stem brittleness can occur. When applying and incorporating this product after potato plants have fully emerged, do not completely cover the foliage with treated soil. Likewise, **DO NOT** completely cover foliage during subsequent cultivations. Be careful that incorporation equipment does not damage seed pieces or elongating sprouts.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

RADISH

Apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.
Fine	7.5 lbs.

RAPESEED (CANOLA) AND CRAMBE

Apply and incorporate this product in the spring or in late summer or in the early fall before planting. See instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.
Fine	10 lbs.

Precautions:

DO NOT apply to rapeseed (canola) in the state of Alaska. Where applications are made in late summer or fall, plant as rotation crops in the season following application only those crops to which This product may be applied as a preplant incorporated treatment or crop injury may occur. **DO NOT** graze or harvest treated crop for livestock forage.

Rapeseed (Canola), Safflower and Sunflower

For Distribution and Use Only in Montana

Apply and incorporate this product in the fall after September 1 or in the spring before planting. Make only one application of this product per crop cycle. Follow soil preparation, application and incorporation instructions in the this product product label.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.
Fine	10 lbs.

Use 7.5 to 10 pounds of this product per acre on coarse and medium soils with 2-5% organic matter.

Precautions:

Rotational Crop Planting Restriction: Plant only spring seeded barley (grown under irrigated conditions), rapeseed, safflower or sunflower as rotational crops in the crop year following the crop treated with this product. If one of these specified crops is not planted, the land should be left idle or fallow for the entire crop year following the crop treated with this product.

SAFFLOWER

Apply and incorporate this product in the spring before planting, or in the fall. See instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre

Soil Texture	Spring† Application	Fall†† Application
Coarse	5 lbs.	7.5 lbs.
Medium	6.25-7.5 lbs.	10 lbs.
Fine	7.5-10 lbs.	12.5 lbs.

†Spring Application:

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10-12.5 lbs.

- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

††This product may be fall applied to Safflower in Arizona, California, Idaho, Nevada, Oregon, Utah, Washington, and Wyoming.

SMALL GRAINS (BARLEY, DURUM AND WHEAT)

Special Precautions for Use of this product on Small Grains

Carefully follow directions for use of this product on small grains to minimize potential crop stress. Under certain conditions, delayed crop emergence and/or stand reduction may occur when this product is applied to barley, durum or wheat. The combined effect of certain cultural practices and unfavorable soil or environmental conditions may cause excessive crop seedling stress resulting in retarded crop growth, stand reduction and possibly reduced yield.

For best results, observe the following cultural practices or precautions:

Use tillage methods that provide a uniformly firm seedbed and time tillage operations to conserve moisture.

If irrigation is used, irrigate prior to planting or after germination and emergence. Moisture received between planting and emergence may cause crusting, especially on loose, friable seedbeds.

DO NOT exceed recommended application rates for this product. This is particularly important on coarse textured or low organic matter soils.

Carefully follow incorporation directions. When applying preplant incorporated treatments, operate equipment at recommended depth and speed to place this product into the upper 1 to 1.5 inches of soil. If applied after planting, set equipment so as to not disturb planted seed.

Set drills to place seed at a depth of 1 to 2 inches. A planting depth greater than 2.5 inches for spring wheat or durum will result in increased seedling stress and decreased emergence.

Use only high quality seed where this product is to be applied (avoid use of small seed with low starch reserves).

If seed treatments are used, apply at the correct rate and uniformly across all seeds. Misapplication may result in reduced germination and/or seedling vigor.

Avoid use of seed varieties known to have poor seedling (emergence) vigor.

DO NOT fall apply this product in combination with any other preplant incorporated herbicide.

Soil characteristics and environmental conditions which may contribute to crop seedling stress that may be accentuated by use of this product include:

- Soil related: High salinity, eroded knolls/hilltops, loose dry soils and compaction.
- Weather related: Cold and/or wet soils, excessively hot soils, excessive moisture, drought, and soil crusting from heavy rainfall.

Note:

DO NOT apply this product on small grains where a dinitroaniline herbicide such as this product was applied at a rate recommended for row crops (oil seeds) during the previous growing season.

Populations of green foxtail (pigeongrass) resistant to the dinitroaniline (DNA) class of herbicides have been identified in Bottineau, McHenry, Montrail, Ramsey, Renville, Rolette, Towner and Ward counties of North Dakota in fields which have a long history of dinitroaniline herbicide use. This product will not control green foxtail which has developed DNA resistance. Therefore, the grower assumes the risk of nonperformance due to DNA resistance if this product is used to control green foxtail in spring cereal grain production areas of North Dakota. Alternative green foxtail control

practices should be utilized in these fields. To prevent or delay the development or spread of DNA-resistant green foxtail in spring cereal production areas, utilize the following management practices:

1. Rotate herbicides so that the same product or same class of herbicide is not used repeatedly year after year. This product and/or other dinitroaniline herbicides should not be applied in consecutive years. Use other herbicides with different modes of action such as Tiller™, Hoelon™, Dakota™, or Stampede™ in alternate years.
2. Rotate crops and use alternative weed control methods, including tillage, fallow periods and/or other herbicides with different modes of action.
3. Thoroughly clean small grains harvested from fields with confirmed resistance before using as seed, or avoid using grain from DNA-resistant fields for seed.
4. Thoroughly clean all crop residues from tillage and harvesting equipment before moving out of fields with confirmed resistance.

Incorporation Directions and Equipment for Small Grains:

Use incorporation equipment in the manner described below unless otherwise specified in use directions. Any of the following implements listed below may be used for the first incorporation of This product. Use only the disc or field cultivator for the second incorporation pass and incorporate in a different direction. Poor weed control may result if untreated soil is moved to the surface during the second incorporation pass. To avoid this problem, the second incorporation should not be deeper than the first.

Chisel Plow:

May be used for the first incorporation pass only. Operate at 4 to 5 inches deep and 4 to 6 mph. A chisel plow is defined as having 3 rows of 14 to 18 inch sweeps on no greater than 12 inch centers. Stagger sweeps so that no soil is left unturned. Use of chisel points may result in inadequate soil mixing.

Tandem Disc:

Operate at 3 to 4 inches deep and at 4 to 6 mph.

Field Cultivator:

Operate at 3 to 4 inches deep and at 5 or more mph. A field cultivator is defined as having 3 to 4 rows of sweeps with "c" or "s" shaped shanks on no greater than 7 inch centers. Stagger sweeps so that no soil is left unturned.

SMALL GRAINS

Barley, Spring Seeded--Fall Application for General Weed Control During the Following Season (For Use in Minnesota, North Dakota, and South Dakota)

For general weed control (except for special rate or use programs) in Minnesota, North Dakota and South Dakota.

Apply in the fall for general weed control during the following growing season. Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and ensure uniform distribution of This product in treated soil. For best weed control results, the second incorporation should occur at least 7 days after the first.

Note:

See recommendations on incorporation and incorporation equipment at the beginning of the "Small Grains" section.

Planting Directions:

Set planting equipment to place seed approximately 2 inches deep.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.

Fine	7.5 lbs.
------	----------

Precautions:

While use of this practice may result in a stand reduction, slight stand reductions **DO NOT** normally affect yield.

Barley, Spring Seeded--Spring Application for Foxtail (Pigeongrass) Control (For Use in Minnesota, North Dakota, and South Dakota)

Apply in the spring as a preplant incorporated treatment for foxtail (pigeongrass) control in spring seeded barley.

Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and ensure uniform distribution of this product in treated soil. For best weed control results, the second incorporation should occur at least 7 days after the first.

Note:

See recommendations on incorporation and incorporation equipment at the beginning of the "Small Grains" section.

Planting Directions:

Set planting equipment to place seed approximately 2 inches deep.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	5 lbs.
Fine	5 lbs.

Precautions:

While use of this practice may result in a stand reduction, slight stand reductions **DO NOT** normally affect yield.

Barley, Spring Seeded--Spring Application for Foxtail (Pigeongrass) Control in Barley Used as a Cover Crop or in the Acreage Conservation Reserve Program (For Use in Minnesota, North Dakota and South Dakota)

Apply in the spring as a preplant incorporated treatment for foxtail (pigeongrass) control in spring seeded barley used as a cover crop or to acreage in the conservation reserve programs.

Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and ensure uniform distribution of this product in treated soil. For best weed control results, the second incorporation should occur at least 7 days after the first.

Note:

See recommendations on incorporation and incorporation equipment at the beginning of the "Small Grains" section.

Planting Directions:

Set planting equipment to place seed approximately 2 inches deep.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	7.5 lbs.
Fine	7.5 lbs.

Precautions:

Use of this practice may result in a slight stand reduction. Follow the most severe grazing restrictions imposed by either the label for This product or the USDA Acreage Conservation Reserve Program, whichever is longest. Consult the local ASCS office or other state agency to determine the period of the USDA grazing restriction.

Spring Seeded Wheat or Durum--Spring, Preplant Incorporated Application for Suppression of Foxtail (Pigeongrass) (For Use in North Dakota West of the Red River Valley and in South Dakota)

Apply this product in the spring as a preplant, incorporated treatment for suppression of foxtail (pigeongrass) in spring seeded wheat and durum. Broadcast 4 lbs. per acre for all soil types.

Incorporation Directions:

The first incorporation should occur within 24 hours after application. Incorporate a second time before planting to destroy existing weeds and ensure uniform distribution of product in treated soil. The second incorporation should be delayed at least 14 days after the first. Operate equipment to place this product into the upper 1.5 to 2 inches of soil.

Note:

See recommendations on incorporation and incorporation equipment at the beginning of the "Small Grains" section. Use a chisel plow (first incorporation only) or a field cultivator.

Incorporation Equipment:

See instructions for chisel plow and field cultivator in the "Small Grains" section.

Planting Directions:

Set equipment to place seed 1.5 to 2 inches deep.

Precautions:

Carefully follow directions under the "Product Information" section for "Small Grains" except place this product into the upper 1.5 to 2 inches of soil.

Spring Seeded Wheat or Durum--Fall Applied, Preplant Incorporated for Suppression of Foxtail (Pigeongrass) (For Use in North Dakota West of the Red River Valley and in South Dakota)

Apply this product in the fall as a preplant incorporated treatment for suppression of green and yellow foxtail (pigeongrass) in spring seeded wheat and durum.

Incorporation Directions:

Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and ensure uniform distribution of product in treated soil. The second incorporation should be 3 to 5 days after the first. Operate equipment to place this product into the upper 1.5 to 2 inches of soil.

Note:

See recommendations on incorporation and incorporation equipment at the beginning of the "Small Grains" section.

Incorporation Equipment:

See instructions for chisel plow or field cultivator in the "Small Grains" section.

Planting Directions:

Set equipment to place seed 1.5 to 2 inches deep.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	3.5-4 lbs.
Medium	3.5-4 lbs.
Fine	5 lbs.

Use the lower rate where light weed pressure is anticipated.

Precautions:

Carefully read and follow "Special Precautions for Use of this product in Small Grains" before application.

Spring Seeded Wheat or Durum--Fall Application for Foxtail (Pigeongrass) Control During the Following Growing Season

Apply in the fall for foxtail (pigeongrass) control during the following growing season. Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and ensure a uniform distribution of this product in treated soil.

Note:

See recommendations on incorporation and incorporation equipment at the beginning of the "Small Grains" section.

Planting Directions:

Set planting equipment to place seed approximately 2 inches deep.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	5 lbs.
Fine	5 lbs.

Precautions:

While use of this practice may result in a stand reduction, slight stand reductions do not normally affect yield.

Winter Wheat--Trifluralin 10G Preplant Incorporated for Control of Cheatgrass and Other Annual Grasses (For Use in Colorado, Idaho, Kansas, Nebraska, Oregon, and Washington)

Apply this product as a preplant incorporated treatment up to 3 weeks before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	7.5 lbs.
Medium	7.5 lbs.
Fine	10 lbs.

Incorporation Directions:

Incorporate this product with a flexible tine-tooth harrow (Flextine or Melroe), set to cut 1 to 2 inches deep and operated at 3 to 6 mph. Thorough incorporation requires 2 incorporation passes over the field in different directions. Incorporate 1 time within 24 hours after application. A second incorporation pass prior to planting should occur at least 5 days after the first. **DO NOT** till the soil with a disc after this product has been incorporated with a flexible tine harrow.

Planting Directions:

Use only a deep furrow or semi-deep furrow drill that will place the seed below the zone into which this product has been incorporated.

Precautions:

DO NOT plant wheat directly into the zone of soil treated with this product as crop injury (delayed emergence or stand reduction) may occur. Delayed emergence or slight stand reduction does not normally affect yield.

Winter Wheat--Fallow Soil Application of Trifluralin 10G for Annual Grass Control Prior to Planting (For Use in Colorado, Idaho, Kansas, Nebraska, Oregon, and Washington)

Apply and shallowly incorporate this product up to 4 months before planting to control cheatgrass and certain other annual grasses and broadleaf weeds during the fallow period and during the following growing season. Apply any time from May to September prior to fall planting of winter wheat.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	7.5 lbs.

Medium	7.5 lbs.
Fine	10 lbs.

Incorporation Directions:

Incorporate this product with a flexible tine-tooth harrow (Flextine or Melroe), set to cut 1 to 2 inches deep and operated at 3 to 6 mph. Thorough incorporation requires 2 incorporation passes over the field in different directions. Incorporate 1 time within 24 hours after application. A second incorporation pass prior to planting should occur at least 5 days after the first. **DO NOT** till the soil with a disc after this product has been incorporated with a flexible tine harrow.

Planting Directions:

Use only a deep furrow or semi-deep furrow drill that will place the seed below the zone into which this product has been incorporated.

Precautions:

DO NOT plant wheat directly into the zone of soil treated with this product as crop injury (delayed emergence or stand reduction) may occur. Delayed emergence or slight stand reduction does not normally affect yield.

Summer Fallow Weed Control with this product Followed by Spring Seeded Wheat, Durum or Barley

This product may be applied to coarse medium and fine textured soils for control of labeled weeds in the summer fallow period and for pigeweed (foxtail) control in wheat, durum and barley seeded the following spring.

This product may be applied to standing stubble or land which has been fallowed or pretilled. Existing weeds, surface trash or crop litter should be reduced by tillage if present in quantities that will prevent uniform soil incorporation. The first incorporation is required within 24 hours after application. The second incorporation may occur in conjunction with tillage to destroy resistant weed growth during the remainder of the fallow year. During the fallow year, susceptible weeds may not be controlled until after the second incorporation.

Broadcast Application Rates/Acre

Application Date	Areas with less than 10 inches annual rainfall	All other areas
Apr. 15-Apr. 30	8.75 lbs.	10 lbs.
May 1-May 31	7.5-8.75 lbs.	8.75-10 lbs.
June 1-June 30	6.25-7.5 lbs.	7.5-8.75 lbs.
July 1-July 31	5-6.25 lbs.	6.25-7.5 lbs.
Aug. 1-Aug.31	5 lbs.	5-6.25 lbs.

Where a rate range is shown, use the higher rate per acre during the early part of an application period and the lower rate per acre during the latter part of an application period.

Incorporation:

See recommendations on incorporation and incorporation equipment at the beginning of the "Small Grains" section.

Seeding Directions:

Wheat, durum or barley should be seeded approximately 2 inches deep.

Precautions:

While use of this practice may result in a stand reduction, slight stand reductions do not normally affect yield.

SOYBEANS

Apply and incorporate this product in the spring prior to planting or in the fall. See instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre

Soil Texture	Spring† Application	Fall†† Application
--------------	---------------------	--------------------

Coarse	5 lbs.	10 lbs.
Medium	7.5 lbs.	10 lbs.
Fine	10 lbs.	12.5 lbs.

†Spring Application:

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10-12.5 lbs.

††Fall application use rates for soybeans grown in Alabama, Arkansas, northern Florida, Georgia, Louisiana, Mississippi, southwestern Missouri (Bootheel), North Carolina, Oklahoma, South Carolina, Tennessee, and Texas. For soybeans grown in areas other than those listed above, fall apply this product at broadcast rates recommended for areas receiving greater than 20 inches of annual rainfall and irrigation.

Precautions:

DO NOT fall apply this product in the fall to soils which are wet or subject to prolonged periods of flooding, or where rice was grown the previous year.

Preemergence Overlay Applications Following Trifluralin 10G (Not for Use in California)

Apply This product as a preplant incorporated treatment. Additional weeds tolerant to This product may be controlled using preemergence overlay applications of Canopy, Dual, Gemini, Lasso, Lexone, Lorox, Lorox Plus, Preview, Pursuit†, Scepter††, Sencor, or other products registered for use on soybeans. Such treatments may be made, unless use following a this product application is specifically prohibited by the product label. Consult the overlay product label for additional weeds controlled, directions for use, cautions and limitations before use.

†The use of Pursuit is limited to certain states. Use Pursuit as an overlay treatment following this product only in states specified on the Pursuit label.

††Use of Scepter is limited to certain states. **DO NOT** use Scepter as a preemergence overlay application following a this product preplant incorporated treatment in the "Northern Use Area" as defined by the Scepter label.

Postemergence Treatments Following Trifluralin 10G (Not for Use in California)

Apply this product as a preplant incorporated treatment. Additional weeds tolerant to this product may be controlled using postemergence applications of Basagran, Blazer, Classic, Cobra, Galaxy, Pinnacle, Pursuit†, Reflex, Scepter††, Storm, Tackle, or other products registered for use on soybeans. Such treatments may be made, unless use following a this product application is specifically prohibited by the product label. Consult the overlay or postemergence product label for additional weeds controlled, directions for use, cautions, and limitations before use.

†The use of Pursuit is limited to certain states. Use Pursuit as a post-emergence treatment following This product only in states specified on the Pursuit label.

††Use of Scepter is limited to certain states. **DO NOT** use Scepter as a postemergence application following a This product preplant incorporated treatment in the "Northern Use Area" as defined by the Scepter label.

SUGAR BEETS

Apply and incorporate this product when sugar beets are 2-6 inches tall.

Precautions:

Exposed sugar beet roots should be covered with soil before applying this product to reduce the possibility of girdling. When incorporating, set equipment to move treated soil into the row. Set incorporation equipment carefully so as to prevent damage to sugar beet tap root.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.

Medium	6.25-7.5 lbs.
Fine	6.25-7.5 lbs.

Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

SUGARCANE

Apply and incorporate this product twice a year. Make the first application in the fall on firmly packed beds immediately after the seed pieces are planted. Make a second application in the spring before or shortly after the cane emerges. Loosen rain-packed beds 2-3 inches deep before spring application. Take care that incorporation equipment does not damage the seed pieces or emerging shoots.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
All soil textures	10-20 lbs.

Application rate range may be adjusted according to weed pressure.

Applications up to Lay-by for Plant Cane or Ratoon Cane (For Use in Louisiana and Texas)

Apply and incorporate this product in the spring from shortly before or after cane emergence until lay-by. Apply after beds have been shaved or false shaved. Loosen rain-packed beds 2-3 inches deep before application. Avoid incorporation equipment damage to seed pieces or emerging shoots. Incorporate with a rolling cultivator or bed chopper for all soil textures. Set rolling cultivator to cut 2 to 4 inches deep and operate at 6 to 8 mph. Set bed chopper to cut 3 to 4 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
All soil textures	10-20 lbs.

Application rate range may be adjusted according to weed pressure.

Itchgrass (Raoulgrass) Control (For Use in Louisiana)

Apply and incorporate this product on plant or ratoon cane. Follow use directions in preceding section for lay-by application.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
All soil textures	10-20 lbs.

Application rate range may be adjusted according to weed pressure.

SUNFLOWER

Apply and incorporate this product in the spring or in the fall between September 15 and December 31 in California, Minnesota, North Dakota, and South Dakota, and between October 15 and December 31 in other states.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5-10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

TOMATO

For direct seeded tomato, apply this product at blocking or thinning to the soil between rows and beneath plants and incorporate. For transplant tomato, apply and incorporate prior to transplanting only. **DO NOT** apply this product after transplanting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5-10 lbs.

- Coarse and medium soils with 2-5% organic matter--7.5 lbs.
- Fine soils with 2-5% organic matter--10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

TREE AND VINE CROPS--CITRUS, FRUIT AND NUT CROPS AND VINEYARDS Application to New Plantings of Citrus, Fruit and Nut Crops

For new plantings of almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, plum, prune, tangelo, tangerine and walnut trees, apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5 lbs.
Medium	6.25-7.5 lbs.
Fine	7.5 lbs.

- All soils with 2-5% organic matter--7.5-10 lbs.
- All soils with 5-10% organic matter--10 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Application to New Plantings of Vineyards

For new plantings of vineyards, apply and incorporate this product before planting.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
Coarse	5-7.5 lbs.
Medium	7.5-15 lbs.
Fine	15-20 lbs.

- All soils with 2-10% organic matter--15-20 lbs.
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Note:

DO NOT use more than 10 lbs. per acre on heat treated grape rootings.

Application to Established Non-Bearing and Bearing Citrus, Fruit and Nut Crops and Vineyards

This product may be applied in established non-bearing and bearing vineyards and plantings of almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, plum, prune, tangelo, tangerine and walnut trees. In established plantings, apply to the soil surface and use incorporation methods not injurious to the crop. **DO NOT** apply to vineyards within 60 days of harvest.

Broadcast Application Rates/Acre

Soil Texture	Trifluralin 10G
All soil textures	10-20 lbs.

Use the higher rate in the rate range for longer term weed control.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container only. In case of spill, contain material and dispose as waste.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling, if available. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in sanitary landfill or by incineration, in accordance with applicable regulations, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, LLC, its Supplemental Distributors, or the Seller. All such risks shall be assumed by the Buyer.

ALBAUGH, LLC, its Supplemental Distributors and the Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. NEITHER ALBAUGH, LLC NOR ITS SUPPLEMENTAL DISTRIBUTORS MAKE ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

BUYER'S EXCLUSIVE REMEDY AND THE EXCLUSIVE LIABILITY OF ALBAUGH, LLC, ITS SUPPLEMENTAL DISTRIBUTORS AND THE SELLER FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. IN NO CASE SHALL ALBAUGH, LLC, ITS SUPPLEMENTAL DISTRIBUTORS, OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALBAUGH, LLC, its Supplemental Distributors, and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of ALBAUGH, LLC

No employee or agent of ALBAUGH, LLC, its Supplemental Distributor, or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

All trademarks are property of their respective owners.

101520

LABEL HISTORY

Not Part of Final printed Label

File Name	Version Mark	Comment
042750-00034.20201015.DRAFT	101520	Label Amendment to EPA