SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR D-638 BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the lab supplement.

Albaugh D-638

EPA Reg. No. 42750-36

MAY 2 2 2001

Under the Pederal Insecticides, Fungicides, and Rodenticide Act. as amended, for the proficide registered under EPA Rep. No. A 2750-36

COTTON STUBBLE

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of application.

For use on cotton stubble:

Weeds	Rate Per Acre	Directions
Annual broadleaf weeds	1-1/3 to 2-2/3 pts.	Use the lower rate when weeds are small (2 to 3 inches tall) and growing actively. Use the higher rate on older and drought-stressed plants, except as directed in the SPECIAL WEED PROBLEMS section of the label.
Biennial broadleaf weeds	2-2/3 to 5-1/3 pts.	Spray while musk thistles or other biennial species are in the seedling to rosette stage, and before flower stalks become apparent. The lower rate can be used in spring during rosette stage. In fall or after flower stalks have developed, use the highest rate, except as directed in the SPECIAL WEED PROBLEMS section of the label.
Perennial broadleaf weeds (including cotton regrowth)	1-1/2 pts. to 4 qts.	Spray to actively growing weeds, including cotton stubble regrowth, following harvest. Do not disturb treated area for at least 2 weeks after treatment, or until weed tops are dead, except as directed in the SPECIAL WEED PROBLEMS section of the label.

A minimum spray volume of 5 gallons per acre by air or 10-15 gallons per acre by ground is recommended. Do not graze diary animals on treated areas within 7 days after application. Do not graze meat animals on treated areas within 3 days before slaughter. Do not cut treated grass for hay within 30 days of application.

Within 29 days following an application of this product, plant only those crops named as use sites on the D-638 label or other registered 2,4-D labels. Follow more specific limitations, if any, provided in the directions for individual crops. Labeled crops may be at risk for crop injury or loss when planted soon after application, especially in the first 14 days. All other crops may be planted 30 or more days following an application without concern for illegal residues in the planted crop. However, under certain conditions, there may be risk of injury to susceptible crops.

See the "GENERAL INFORMATION" and "MIXING INSTRUCTIONS" sections of the label for D-638 for essential product information.

Read the "WARRANTY LIMITATIONS AND DISCLAIMER" in the label for D-638 before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc. 121 NE 18" Street Ankeny, IA 50021