

42750-59

02/25/2009

1/3



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460**

**PESTICIDES AND**

**OFFICE OF  
PREVENTION,  
TOXIC SUBSTANCES**

FEB 25 2009

Mr. Morris Gaskins  
Albaugh, Inc  
P. O. Box 2127  
Valdosta, GA 31604

Dear Mr. Gaskins:

Subject: Glyphosate 53.8% (Add Use on Roundup Ready Sugarbeets)  
EPA Registration No. 42750-59  
Application Dated January 20, 2009

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable. A stamped copy of labeling is enclosed for your records.

Supplemental labeling must be incorporated into master labeling and copies of master labeling submitted to the Agency for our files at your next printing or within two years from the date of acceptance of supplemental labeling, whichever comes first. The Agency will consider a convincing argument as to why a supplemental label should not appear on a master label on a case by case basis.

Sincerely,

*Vukie Swatter for*  
James A. Tompkins  
Product Manger 25  
Herbicide Branch  
Registration Division (7505P)

### SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 53.8% HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL

When using Glyphosate 53.8% herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

### GLYPHOSATE 53.8%

EPA Reg. No. 42750-59

### FOR WEED CONTROL APPLICATIONS IN SUGAR BEETS WITH THE ROUNDUP READY® GENE

**ACCEPTED**


FEB 25 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. 42750-59

KEEP OUT OF REACH OF CHILDREN.

CAUTION

In case of an emergency involving this product, Call CHEMTREC toll free at 1-800-424-9300



### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, ESPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIREABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENREAL INFORMATION" and "MIXING" sections of the label booklet for GLYPHOSATE 53.8% herbicide for essential product performance information.

ALBAUGH RECOMMENDS USE OF THIS PRODUCT FOR POSTEMERGENCE APPLICATION ONLY ON SUGAR BEET VARIETIES DESIGNATED AS CONTAINING A ROUNDUP READY GENE.

The Roundup Ready designation indicates that the sugar beet contains a patented gene, which provides tolerance to this product. Information on Roundup Ready sugarbeet may be obtained from your seed supplier or Albaugh representative. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.

See the "ROUNDUP READY CROPS" section of the Glyphosate 53.8% herbicide label booklet for general precautionary instructions for use in Roundup Ready crops. Do NOT combine these instructions with other recommendations made for crop varieties that do not contain a Roundup Ready gene listed in the "ANNUAL AND PERENNIAL CROPS (Alphabetical)" sections of the Glyphosate 53.8% herbicide label booklet.

3  
3

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop)

APPLICATION INSTRUCTIONS

MAXIMUM ALLOWABLE APPLICATION RATES

Combined total per year for all application	192 fluid ounces per acre
Preplant, Preemergence applications	120 fluid ounces per acre
Emergence to 8 leaf stage	60 fluid ounces per acre
Between 8 leaf stage and canopy closure	48 fluid ounces per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. Tank mixtures of this product with herbicides, insecticides or fungicides may result in crop injury or reduced weed control.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting of Roundup Ready sugar beets.

PRECAUTIONS, RESTRICTIONS: Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 120 fluid ounces per acre per season.

Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied over the top of Roundup Ready sugar beets for control of annual grasses and broadleaf weeds from emergence to 30 days prior to harvest. To maximize yield potential, spray sugar beets early to eliminate competing weeds. Up to 4 sequential applications of this product may be made with at least 10 days between applications. This product will control or suppress most perennial weeds. For some perennial weeds, repeat applications may be required to eliminate crop competition throughout the growing season.

RESTRICTIONS: Follow all general precautionary instructions for use in Roundup Ready crops.

- The combined total application from crop emergence through harvest must not exceed 108 fluid ounces per acre.
- The maximum rate for any single application between emergence to the 8 leaf stage is 36 fluid ounces per acre.
- The maximum rate for any single application between the 8 leaf stage and canopy closure is 24 fluid ounces per acre.
- Allow a minimum of 30 days between last application and sugar beet harvest.
- For any crop NOT listed in the "CROPS" section of this label booklet, applications must be at least 30 days prior to planting.

Manufactured By:  
Albaugh, Inc.  
Ankeny, IA 50021