

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% AND ARSENAL™ 2 WSL BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label for Glyphosate 41% and the label instructions for Arsenal 2 WSL and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



**GLYPHOSATE 41% PLUS ARSENAL 2 WSL FOR NONCROP AREAS SUCH AS
NONIRRIGATION DITCHBANKS, DRY DITCHES, DRY CANALS, FENCEROWS,
ROADSIDES, INDUSTRIAL SITES, PIPELINES, UTILITY RIGHTS-OF-WAY AND
RAILROADS.**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

When applied as directed, this tank mixture will control or partially control labeled woody brush, trees and herbaceous weeds in noncrop areas. In addition to the weeds labeled for Glyphosate 41% herbicide, this tank mixture will control arrowweed, Saltcedar and yaupon.

HAND-HELD AND HIGH-VOLUME APPLICATIONS

Use 4 to 8 quarts of Glyphosate 41% plus ½ to 4 pints of Arsenal 2 WSL per 100 gallons of spray solution. Nonionic surfactants which are labeled for use with herbicides may be used. If used, add 2 quarts of nonionic surfactant per 100 gallons of spray solution. Apply to foliage of actively growing vegetation. Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

BROADCAST APPLICATIONS WITH GROUND EQUIPMENT

Use 2 to 5 quarts of Glyphosate 41% plus ½ to 4 pints of Arsenal in sufficient water to apply in a total spray volume of 10 to 40 gallons per acre. Apply to foliage of actively growing vegetation.

AERIAL APPLICATIONS

Use 2 to 5 quarts of Glyphosate 41% plus ½ to 4 pints of Arsenal in sufficient water to apply in a total spray volume of 10 to 20 gallons per acre. Apply to foliage of actively growing vegetation.

2/2

Apply to woody brush and trees after full leaf expansion until initiation of fall color.

Avoid direct applications to any body of water. Do not apply on ditches used to transport irrigation water.

Read and carefully observe the label directions, cautionary statements and all information on all the labels of both products used in this tank mixture before proceeding with this recommendation. Additional precautionary statements are made in these labels. Use according to the most restrictive label directions for each product in the mixture.

When used in combination as recommended by Albaugh, Inc., the liability of Albaugh, Inc., shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Albaugh product in such combination use.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

4/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



FOR DISTRIBUTION AND USE ONLY IN WEST TEXAS – IN THE AREA BORDERED BY HWY. 86 ON THE NORTH, HWY. 287 ON THE NORTHEAST, HWY. 283 ON THE EAST, HWY. 190 ON THE SOUTH, HWY. 285 ON THE SOUTHWEST, AND THE TEXAS-NEW MEXICO BORDER ON THE WEST.

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION", "APPLICATION EQUIPMENT AND TECHNIQUES" and "CROPPING SYSTEMS" sections of the label booklet for Glyphosate 41% for essential product performance information.

Read the entire label booklet for Glyphosate 41% herbicide as well as labels for Def 6, Foilex and Prep, if tank mixing, and follow all use directions and observe all precautions contained in these labels.

RECOMMENDATIONS

When applied as directed under the conditions described, this product controls annual and perennial weeds listed in the label booklet for Glyphosate 41% prior to the harvest of cotton.

Broadcast Applications – This product may be applied using either aerial or ground spray equipment. For ground applications with broadcast equipment, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 15 gallons of water per acre.

2/2

FOR AERIAL APPLICATIONS, REFER TO THE "APPLICATION EQUIPMENT AND TECHNIQUES" AND "AERIAL EQUIPMENT" SECTIONS OF THE LABEL BOOKLET FOR GLYPHOSATE 41%.

DO NOT EXCEED A MAXIMUM RATE OF 2 QUARTS PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR.

Weed Control – For specific rates of application and instructions for control of various annual and perennial weeds for this product used alone or in the following tank mixtures, see the "Weeds Controlled" section of the label booklet for Glyphosate 41%.

To control johnsongrass using multiple-directed or broadcast over-the-top spray equipment, apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. Ensure complete coverage.

For partial control of field bindweed, apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 20 gallons of water per acre. Apply when bindweed is actively growing and 12 inches or greater in length. Reduced performance may result if bindweed is under drought stress.

To control silverleaf nightshade (whiteweed), apply 2 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 15 gallons of water per acre. Applications should be made when at least 60 percent of the plants have berries. Treatments must be applied before a killing frost. Do not treat when the plants are under drought stress since good soil moisture is needed for active growth.

TANK MIXTURES

Glyphosate 41% plus Def™ 6
Glyphosate 41% plus Folex™ 6
Glyphosate 41% plus Prep™ 6
Glyphosate 41% plus Prep plus Def 6 or Folex

When applied as recommended under the conditions described, these tank mixtures control annual and perennial weeds listed on the label booklet for Glyphosate 41% prior to the harvest of cotton. For application guidelines, precautions and use-rates, refer to the Def, Folex and Prep labels.

This product when tank-mixed with Def 6 or Folex defoliant may provide enhancement of cotton leaf drop and regrowth inhibition.

Timing of Application – Apply this product or these tank mixtures for preharvest weed control after 60 percent of the cotton bolls have opened.

NOTE: DO NOT APPLY TO CROPS GROWN FOR SEED. Allow a minimum of 7 days between application and harvest. Do not feed or graze treated cotton forage or hay following preharvest applications.

™Def is a trademark of Mobay Chemical Company.

™Folex and Prep are trademarks of Rhone-Poulenc, Inc.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/1


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41%, OUST™ AND 2,4-D AMINE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41%, the label instructions for Oust and 2,4-D amine and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



**GLYPHOSATE 41% HERBICIDE PLUS OUST AND 2,4-D AMINE FOR NONCROP AREAS,
INCLUDING RAILROAD RIGHTS-OF-WAY, UTILITY SUBSTATIONS, AIRPORTS,
INDUSTRIAL PLANTS, PIPELINES, ROADSIDES, STORAGE AREAS, AND SIMILAR SITES.**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

When applied as directed, this tank mixture will control or partially control labeled annual and perennial weeds in noncrop areas.

Apply the recommended rate of Glyphosate 41% plus 1 to 2 pints of 2,4-D amine and 2 to 4 ounces of Oust in 25 to 40 gallons of total spray solution per acre.

Use the higher rates of these mixtures when weed growth is heavy or dense.

Do not apply this tank mixture, drain or flush equipment on or near desirable trees or other plants, on areas where their roots may extend, or in locations where Oust or 2,4-D may be washed or moved into contact with their roots.

Read and carefully observe the label directions, cautionary statements and all information on the labels of each product used in this tank mixture.

Additional precautionary statements are made in these labels. Use according to the most restrictive label directions for each product in the mixture.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% AND FOR GARLON™ 4 BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



GLYPHOSATE 41% PLUS GARLON 4 TANK MIXTURE FOR BURNDOWN AND PARTIAL CONTROL OR SUPPRESSION OF WOODY BRUSH AND WEEDS IN INDUSTRIAL SITES.

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

NONCROP USES

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information and the following NONCROP SECTIONS for specific recommended uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OR DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds or regrowth from areas not adequately contacted by the spray mixture.

Glyphosate 41% does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

INDUSTRIAL SITES

When applied as directed for "Noncrop Uses" under the conditions described, this product plus Garlon 4 provides burndown and partial control or suppression of woody brush and vegetation labeled for this product. This tank mixture is recommended for use on rights-of-way (utility,

railroad, highway, pipeline), fencerows, roadsides, nonirrigation ditchbanks, wasteland and similar noncrop or industrial sites.

Hand-Held and High-Volume Applications:

Use 2 to 4 quarts of Glyphosate 41% herbicide plus 1 to 2 quarts of Garlon 4 per 100 gallons of spray solution and apply to foliage of actively growing woody brush and weeds. Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff.

Broadcast Applications with Ground Equipment:

Use 2 to 4 quarts of Glyphosate 41% plus ½ to 2 quarts of Garlon 4 in sufficient water to make 20 to 100 gallons of total spray per acre.

Aerial Applications (Helicopter Only):

Use 2 to 4 quarts of Glyphosate 41% plus 1 to 2 quarts of Garlon 4 and apply in a total spray volume of 10 to 20 gallons per acre. Aerial sprays should be applied using suitable drift control.

Apply when plants are actively growing and after full leaf expansion of woody brush. Use the higher rates of these products where vegetation is heavy or dense, or where hard-to-control brush species are prevalent. Repeat applications may be necessary to maintain control or suppress areas where canopying of vegetation prevents good spray coverage and penetration.

Nonionic surfactants which are labeled for use with herbicides may be used. Use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 50 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 50 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Read and carefully observe the label claims, cautionary statements and all information on the labels of both products used in this tank mixture. Use according to the most restrictive label directions for each product in the mixture.

When used in combination as recommended by Albaugh, Inc., the liability of Albaugh, Inc., shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Albaugh product in such combination use.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41%, DIURON AND ATRAZINE HERBICIDES BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



**GLYPHOSATE 41% PLUS DIURON PLUS ATRAZINE HERBICIDE TANK MIXTURE FOR
CONTROL OR PARTIAL CONTROL OF HERBACEOUS VEGETATION ONLY ON
RAILROAD RIGHTS-OF-WAYS.**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

NONCROP USES

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information and the following NONCROP SECTIONS for specific recommended uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OR DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Apply when plants are actively growing. Use the higher recommended rates of these products where vegetation is heavy or dense, or where hard-to-control species are prevalent. Repeat applications may be necessary to maintain control where dense vegetation prevents good spray coverage. Applications should be made when weeds are less than 12 inches tall for best results.

Nonionic surfactants which are labeled for use with herbicides may be used. Use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 50 percent active ingredient, or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 50 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Read and carefully observe the label claims, cautionary statements and all information on the labels of both products used in this tank mixture. Use according to the most restrictive label directions for each product in the mixture.

When used in combination as recommended by Albaugh, Inc., the liability of Albaugh, Inc., shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Albaugh product in such combination use.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



**TANK MIXTURES OF GLYPHOSATE 41% PLUS ARSENAL APPLICATORS
CONCENTRATE FOR FORESTRY SITE PREPARATION PRIOR TO PLANTING DOUGLAS
FIR IN WASHINGTON AND OREGON**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

FORESTRY SITE PREPARATION

Use as a site preparation treatment for forestry sites being planted to Douglas fir species in Washington and Oregon.

Apply 2 to 4 quarts of this product with 4 fluid ounces to 8 fluid ounces of Arsenal Applicators Concentrate in 5 to 15 gallons of spray solution per acre as a broadcast spray to control big leaf maple resprouts. Where big leaf maple resprouts are not a primary concern, addition of 1 fluid ounce to 4 fluid ounces per acre of Arsenal Applicators Concentrate to the recommended rate of this product will improve control of most other woody brush species, such as willow, pin cherry, dogwood, and vine maple.

Nonionic surfactants which are labeled for use with herbicides may be used. If used, add 2 quarts of nonionic surfactant per 100 gallons of spray solution.

The tank mixtures may be applied by air (helicopter only).

APPLICATION TIMING

Big leaf maple resprouts should have vigorous growth prior to the application of these tank mixtures. Fall applications will provide best results.

Read and carefully observe the label directions, cautionary statements and all information on the labels of both products used in this tank mixture. Additional precautionary statements are made in these labels. Use according to the most restrictive label directions for each product in the mixture.

2/2

When used in combination as recommended by Albaugh, Inc., the liability of Albaugh, Inc., shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Albaugh product in such combination use.

™ Arsenal is a trademark of American Cyanamid Company.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

4/1


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



LOW-VOLUME BACKPACK APPLICATIONS OF GLYPHOSATE 41% FOR TREES AND BRUSH

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

When applied as directed, this product will control or partially control labeled woody brush and trees in noncrop areas. This application technique is recommended for spot treatment of trees and brush. It is most effective in areas where there is a low density of undesirable trees or brush.

Use a 5 to 10 percent solution of this product in water. Apply the spray solution with a nozzle which produces a straight stream. Begin at the top of the targeted tree and spray the leaves with a lateral zig-zag motion. The spray pattern should go from top to bottom and from side to side of the part of the tree that is covered with leaves. The spray should contact approximately 50 percent of the tree's leaves.

In small, open branched trees, applications need only be made from one side. If the foliage is thick or there are multiple root sprouts, applications must be made from several sides to ensure adequate spray coverage.

Use the higher rates for difficult-to-control species or under stressful environmental conditions.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



FOR USE IN WIPER APPLICATORS

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product may be applied through a wiper applicator in any noncrop site and only when specifically recommended in cropping systems as stated in the label booklet for Glyphosate 41%.

AVOID CONTACT WITH DESIRABLE VEGETATION.

Equipment must be designed, maintained, and operated to prevent the herbicide solution from contacting desirable vegetation.

In severe infestations, reduce equipment ground speed to ensure that adequate amounts of this product are wiped on the weeds. A second treatment in the opposite direction may be beneficial.

APPLICATION RATES – Use solutions consisting of 33 to 100 percent of this product. Limit solutions to 33 percent when using a standard wick-type applicator. Use the more concentrated solutions only with equipment that can handle a thick solution such as the porous plastic-type applicators.

2/2

Add 1 quart of nonionic surfactant for every 4 gallons of solution.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.


Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx

FOR NONSELECTIVE WEED CONTROL IN CHERIMOYA, COCOA BEANS, GENIP AND POMEGRANATES

Keep out of reach of children

CAUTION!



FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product is recommended for weed control in established groves or orchards, or for site preparation prior to transplanting cherimoya, cocoa beans, genip or pomegranates. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, or orchard guns, or with wiper applicator equipment. See the "Application Equipment and Techniques" section of the label booklet for Glyphosate 41% for specific information on the use of equipment.

When applying this product, refer to the "Weeds Controlled" and the "Tree and Vine Crops" sections of the label booklet for rates to be used.

NOTE: Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. This product does not provide residual weed control. For subsequent weed control, use repeated applications of this product. Do not apply more than 10.6 quarts of this product per acre per year.

2/2

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

AVOID PAINTING CUT STUMPS WITH THIS PRODUCT AS INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES.

Allow a minimum of 14 days between the last application and harvest.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% AND ARSENAL™ APPLICATORS CONCENTRATE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Arsenal Applicators Concentrate and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



**TANK MIXTURE OF GLYPHOSATE 41% HERBICIDE AND ARSENAL APPLICATORS
CONCENTRATE FOR FORESTRY CONIFER RELEASE IN THE STATES OF MAINE, NEW
HAMPSHIRE AND VERMONT**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

™Arsenal is a trademark of American Cyanamid Company

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

Apply a mixture of 2 quarts of this product and 1 to 2.5 fluid ounces of Arsenal Applicators Concentrate per acre as release treatment for balsam fir and red spruce.

This mixture is recommended for controlling woody brush, deciduous trees and herbaceous weeds on sites regenerated with balsam fir and red spruce. Make applications only after formation of final resting buds on these conifers.

Use the higher recommended rate for sites with dense, tough-to-control woody brush and deciduous trees.

When using ground application equipment, use 10 to 60 gallons of spray solution per acre. For aerial application (helicopter only), use 5 to 15 gallons of spray solution per acre.

Injury may occur to conifers treated for release, especially where spray patterns overlap. Injury can be accentuated if applications are made when conifers are actively growing or are under stress.

Read and carefully observe the label claims, cautionary statements, and all information on the label for all products used.

2/2

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/3


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% label for tank mix products and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



GLYPHOSATE 41% AND TANK MIXTURES FOR NONCROP AREAS, INCLUDING RAILROAD, RIGHTS-OF-WAY, UTILITY SUBSTATIONS, AIRPORTS, INDUSTRIAL PLANTS, ROADSIDES, STORAGE AREAS AND SIMILAR SITES.

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See the "GENERAL INFORMATION", "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" and "NONCROP USES" sections of the label booklet for Glyphosate 41% for essential product information.

Do not allow spray mixtures of this herbicide to mist, drip, drift or splash onto desirable vegetation since injury or destruction may occur. Do not apply when wind or other conditions favor drift.

See the "WEEDS CONTROLLED" section of the Glyphosate 41% label booklet for rate recommendations. For difficult to control species, where dense stands occur, or where conditions for control are not ideal, 5 to 10 quarts per acre of this product may be used for improved results.

TANK MIXTURES

This product provides control of the emerged weeds listed in the label booklet. When applied as a tank mixture, the following herbicides will provide preemergence and/or postemergence control of the weeds listed in the individual product labels.

The following list of products may be tank mixed with this product. Any recommended rate of this product may be used in a tank mixture with these products.

<u>Tank Mix Product</u>	<u>Rate per Acre</u>
Arsenal™	0.5 to 4 pints
Banvel	1 to 4 pints
2,4-D	0.5 to 1 pound
Garlon™ 3A	1 to 6 pints
Garlon 4	1 to 6 pints
Diuron	4 to 8 pounds
Diuron + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Diuron + Garlon 3A	4 to 10 pounds + 1 to 2 pints
Diuron + Garlon 4	4 to 10 pounds + 1 to 2 pints
Hyvar™ X	4 to 8 pounds
Hyvar X + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Hyvar X + Garlon 3A	4 to 8 pounds + 1 to 2 pints
Hyvar X + Garlon 4	4 to 8 pounds + 1 to 2 pints
Krovar™ I DF	4 to 6 pounds
Krovar I DF + 2,4-D	4 to 6 pounds + 0.5 to 1 pound
Krovar I DF + Garlon 3A	4 to 6 pounds + 1 to 2 pints
Krovar I DF + Garlon 4	4 to 6 pounds + 1 to 2 pints
Oust™	2 to 6 ounces
Oust + 2,4-D	2 to 6 ounces + 0.5 to 1 pound
Oust + Garlon 3A	2 to 6 ounces + 1 to 2 pints
Oust + Garlon 4	2 to 6 ounces + 1 to 2 pints
Spike™ 80W	2 to 5 pounds
Spike 80W + 2,4-D	2 to 5 pounds + 0.5 to 1 pound
Spike 80W + Garlon 3A	2 to 5 pounds + 1 to 2 pints
Spike 80W + Garlon 4	2 to 5 pounds + 1 to 2 pints

*Arsenal is not approved for use in the state of California.

Refer to the individual product labels for specific noncrop sites, rates, carrier volumes and precautionary statements.

Read and carefully observe the label claims, cautionary statements, recommended use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Maintain good agitation at all times during the mixing process. Ensure that the tank mix products are well mixed with the spray solution before adding this product.

Mix only the quantity of spray solution which can be used during the same day. Tank mixtures allowed to stand overnight may result in reduced weed control.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Nonionic surfactants which are labeled for use with herbicides may be used. Use a 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution). Use surfactants that contain at least 50 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

When used in combination as recommended by Albaugh, Inc., the liability of Albaugh, Inc., shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Albaugh product in such combination use.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

™Arsenal is a trademark of American Cyanamid Company.

™Banvel is a trademark of Sandoz Crop Protection Corporation.

™Garlon and Spike are trademarks of DowElanco Products Company.

™Hyvar, Krovar and Oust are trademarks of E.I. duPont de Nemours and Company.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/1


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



GLYPHOSATE 41% FOR USE IN COCONUTS

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product may be applied to control annual and perennial weeds listed on this label in established plantings of coconut and for site preparation prior to transplanting coconut trees.

Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns, or with wiper equipment.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT OR OTHER PARTS OF TREES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

AVOID PAINTING OUT STUMPS WITH THIS PRODUCT AS INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41%, container labels and this supplement.

WARNING: USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



USE ONLY FOR SEED PRODUCTION OF SOYBEANS WITH THE ROUNDUP READY® GENE.

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300
Roundup Ready® is a registered trademark of the Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATIONS INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT SOYBEANS IN PRODUCTION FIELDS OF SOYBEANS CONTAINING THE ROUNDUP READY® GENE. SEVERE INJURY OR DEATH WILL RESULT IF SOYBEAN VARIETIES WHICH DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT. AVOID CONTACT OF THE SPRAY WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS OR ANY DESIRABLE PLANTS AND TREES, OTHER THAN SOYBEANS WITH THE ROUNDUP READY GENE. SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

2/2

ROUNDUP READY® SOYBEAN VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE SOYBEAN VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT. IT IS UNLAWFUL TO SELL OR PLANT SAVED SEED.

SOYBEANS WITH THE ROUNDUP READY® GENE MAY ONLY BE USED FOR PLANTING A COMMERCIAL CROP IN A SINGLE SEASON. SEED MAY NOT BE SAVED FOR REPLANTING AND SAVED SEED MAY NOT BE SUPPLIED TO OTHERS FOR REPLANTING. ALBAUGH, INC. DOES NOT WARRANT THE SAFETY OR PERFORMANCE OF THIS PRODUCT WHEN USED ON "BROWN BAG" OR FARMER-MADE SEED.

USE RECOMMENDATIONS:

When applied as directed under the conditions described, this product will control non-glyphosate tolerant soybeans in seed production fields of soybeans containing the Roundup Ready gene. This product may be applied using ground spray equipment only. Apply 1 quart of this product plus 0.5 to 1.0% nonionic surfactant by total spray volume in 5 to 20 gallons of spray solution per acre as a broadcast spray.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT.

Application Timing – This product can be applied to Roundup Ready soybeans from the cracking stage through the flowering stage.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

11


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



SHIELDED APPLICATOR USE IN GRASSES GROWN FOR SEED IN IDAHO, OREGON AND WASHINGTON ONLY.

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

When applied using shielded applicator equipment designed to prevent direct contact, movement of spray droplets, or mist onto desirable grasses grown for seed production, this product may be used to control labeled weeds. Use of low spray pressure through low-pressure nozzles will minimize the potential of spray drift.

Apply 1 to 3 quarts of this product per acre as a broadcast spray in 10 to 20 gallons of total spray volume per acre.

Uniform planting in straight rows aid in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by or through the protective shields.

GROWER ASSUMES ALL RESPONSIBILITY FOR CROP LOSSES FROM MISAPPLICATION.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



GLYPHOSATE 41% FOR USE IN HYBRID POPLAR (*Populus spp.*) PRODUCTION.

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

PREPLANT

This product is recommended for use prior to planting *Populus spp.* This includes, but is not limited to hybrid poplars and hybrid cottonwoods.

See the Glyphosate 41% label booklet for specific rates for the weeds being controlled.

DIRECTED SPRAYS

Use a 2 percent spray solution as a spray-to-wet application for the control of undesirable woody brush and trees. To control herbaceous weeds, use a 1 to 2 percent solution. Avoid contact of spray, drift, or mist with foliage, green bark or non-woody surface roots of the *Populus spp.*

Mix 2 to 6 quarts of a nonionic surfactant per 100 gallons of spray solution (0.5 to 1.5 percent spray volume). Use a surfactant with greater than 70 percent active ingredient

WIPERS

This product may be used through wick or other suitable wiper applicators for control or partial control of grass and broadleaf weeds listed on the label.

For wick applicators, mix 1 gallon of this product with 2 gallons water to make a 33% solution. For wiper systems that can handle thicker solutions, such as force fed systems, a 33 to 100% Glyphosate 41% solution may be used.

For best results ensure that the herbicide solution is allowed to contact the maximum amount of leaf surface. As weed densities increase, decrease equipment speed to allow sufficient herbicide flow to wet all weed surfaces contacted. Weeds not contacted will be unaffected.

AVOID HERBICIDE CONTACT WITH DESIRABLE VEGETATION. Desirable vegetation contacted by the herbicide solution may be injured or controlled. This includes foliage, fruit, or green stems.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 13th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



RED RICE CONTROL IN TEXAS ONLY

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product may be used to control red rice, prior to planting rice. Apply 1-½ quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Use surfactants with a minimum of 70 percent active ingredient. Add 17 pounds of ammonium sulfate per 100 gallons of water.

Flush fields prior to applications to obtain uniform germination and stand of red rice. Make applications when the majority of the red rice plants are in the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may only be partially controlled. Do not treat rice fields or levees when the fields contain floodwater.

For aerial applications drift control additives may be used.

Avoid spraying during low humidity conditions, as reduced control may result.

Do not re-flood treated fields for 8 days following application.

2/2

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

Label as used in this supplemental labeling refers to the label for Glyphosate 41% and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



**ROUNDUP HERBICIDE FOR CONTROL OR SUPPRESSION OF *BROMUS* SPECIES AND
MEDUSAHEAD RYE FOUND IN INDUSTRIAL, RANGELAND AND PASTURE SITES IN
COLORADO, IDAHO, IOWA, KANSAS, MONTANA, NEBRASKA, NORTH DAKOTA,
OREGON, SOUTH DAKOTA, UTAH, WASHINGTON AND WYOMING**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

When applied as directed in this label, there are no grazing restrictions.

APPLICATION EQUIPMENT AND TECHNIQUES

Applications may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre.

Mix 0.5 to 1.0 percent nonionic surfactant (2 to 4 quarts per 100 gallons of spray solution) with the spray solution. The surfactant should contain at least 80 percent active ingredient.

BROMUS SPECIES

This product may be used to treat downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*) and cheatgrass (*Bromus setchuensis*) found on industrial rangeland and pasture sites. Apply 8 to 16 fluid ounces of product per acre on a broadcast basis.

2/2

For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses are able to become reestablished on the site.

MEDUSAHEAD RYE

To treat medusahead rye, apply 16 fluid ounces of this product per acre as soon as plants are actively growing, and prior to seedhead emergence. Applications may be made in the fall or spring.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

4/6

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41%, container labels and this supplement.

WARNING: USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx

ACCEPTED

APR 25 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 42750-60

FOR SOYBEANS WITH THE ROUNDUP READY® GENE
OVER-THE-TOP APPLICATIONS

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300
Roundup Ready® is a registered trademark of the Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

General Information

NOTE: THIS PRODUCT MAY BE USED ONLY OVER THE TOP OF IMPROVED SOYBEAN VARIETIES THAT ARE DESIGNATED AS SOYBEANS WITH THE ROUNDUP READY® GENE. SEVERE INJURY OR DEATH OF SOYBEANS WILL RESULT IF ANY SOYBEAN VARIETIES NOT DESIGNATED AS HAVING THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT. AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF

2/6

CROPS, OR ANY DESIRABLE PLANTS AND TREES, OTHER THAN SOYBEANS WITH THE ROUNDUP READY GENE, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

ROUNDUP READY® SOYBEAN VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE SOYBEAN VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT. IT IS UNLAWFUL TO SELL OR PLANT SAVED SEED.

SOYBEANS WITH THE ROUNDUP READY® GENE MAY ONLY BE USED FOR PLANTING A COMMERCIAL CROP IN A SINGLE SEASON. SEED MAY NOT BE SAVED FOR REPLANTING AND SAVED SEED MAY NOT BE SUPPLIED TO OTHERS FOR REPLANTING. ALBAUGH, INC. DOES NOT WARRANT THE SAFETY OR PERFORMANCE OF THIS PRODUCT WHEN USED ON "BROWN BAG" OR FARMER-MADE SEED.

Application Instructions

This product may be applied postemergence to soybeans with the Roundup Ready Gene from the cracking stage through the flowering stage. Single and repeat in-crop applications of this product are not to exceed 2 quarts per acre per growing season. Preharvest applications are not to exceed 1 quart per acre. For preharvest applications in soybeans, refer to the "PREHARVEST APPLICATIONS" section of the Glyphosate 41% label booklet. Total Glyphosate 41% use should not exceed 8 quarts per acre per year. Applications may be made following pre-plant burndown applications of products containing Glyphosate 41%, or on conventionally tilled seedbeds.

When applied as directed under the conditions described, this product will control labeled annual grasses and broadleaf weeds in soybeans with the Roundup Ready gene. Many perennial grass and broadleaf weeds will be controlled or suppressed with one or more applications of this product.

Applications should be made to actively growing weeds before they reach the maximum size listed for each recommended rate in the "WEEDS CONTROLLED" section of this label. Refer to the "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" of the label booklet for proper use instructions.

There are no rotational crop restrictions following applications of this product.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS OTHER THAN SOYBEANS WITH THE ROUNDUP READY® GENE. Do not allow the herbicide solution to mist, drip, drift or splash onto other desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of plant injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions which allow spray drift to occur such as combinations of pressure and nozzle type that will result in fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR SPRAY PRESSURE.**

For ground applications: Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. See the "WEEDS CONTROLLED" section of this label for specific recommended rates. Carefully select proper nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

3/6

For aerial applications: Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart per acre. See the "WEEDS CONTROLLED" section on this label. AVOID DRIFT – DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

WEEDS CONTROLLED

APPLICATION RATES – ANNUAL WEEDS:

Rate per acre: 12 fluid ounces

WEED	WEED HEIGHT (INCHES)
Foxtail	Up to 18

Rate per acre: 16 fluid ounces

<u>WEED</u>	<u>WEED HEIGHT (INCHES)</u>
Barley	Up to 18
Rye	Up to 18
Wheat	Up to 18
Chickweed	Up to 12
Cocklebur	Up to 12
Corn, volunteer	Up to 12
Crabgrass	Up to 12
Mustard*	Up to 12
Pigweed sp.	Up to 12
Seedling Johnsongrass	Up to 12
Shattercane	Up to 12
Sunflower	Up to 12
Witchgrass	Up to 8
Field Pennycress	Up to 6
Groundcherry	Up to 6
Lambsquarters	Up to 6
Panicum, fall	Up to 6
Panicum, Texas	Up to 6
Shepherdspurse	Up to 6
Sprangletop	Up to 6
Woolly cupgrass	Up to 6
Kochia**	3 to 6
Barnyardgrass	Up to 3
Broadleaf signalgrass	Up to 3

4/6

Fleabane, Rough	Up to 3
Goosegrass	Up to 3
Horseweed/marestail	Up to 3
Smartweed, Pennsylvania	Up to 3
Velvetleaf	Up to 3
Junglerice	Up to 3
Sicklepod	Up to 2

*Do not treat mustard after flowering.

**Do not treat Kochia in button stage (1-2" tall)

Rate per acre: 24 fluid ounces

<u>WEED</u>	<u>WEED HEIGHT (INCHES)</u>
Chickweed	12 to 18
Cocklebur	12 to 18
Mustard	12 to 18
Pigweed	12 to 18
Shattercane	12 to 18
Sunflower	12 to 18
Crabgrass	12 to 18
Witchgrass	9 to 12
Field Pennycress	6 to 12
Groundcherry	6 to 12
Kochia	6 to 12
Lambsquarters	6 to 12
Panicum, fall	6 to 12
Shepherdspurse	6 to 12
Sprangletop	6 to 12
Wooly cupgrass	6 to 12
Goosegrass	3 to 6
Barnyardgrass	3 to 6
Broadleaf signalgrass	3 to 6
Fleabane, Rough	3 to 6
Horseweed/marestail	3 to 6
Junglerice	3 to 6
Pennsylvania Smartweed	3 to 6
Velvetleaf	3 to 6
Sicklepod	3 to 4
Black Nightshade	Up to 6
Ragweed, common	Up to 6
Ragweed, giant	Up to 6
Red rice	Up to 3
Hemp sesbania	Up to 2
Morningglory sp.	Up to 2
Prickly sida, Teaweed	Up to 2

5/6

Rate per acre: 32 fluid ounces

<u>WEED</u>	<u>WEED HEIGHT (INCHES)</u>
Groundcherry	12 to 18
Horseweed/marestail	12 to 18
Lambsquarters	12 to 18
Panicum, fall	12 to 18
Scraggletoe	12 to 18
Goosegrass	8 to 12
Black nightshade	6 to 12
Fleabane, Rough	6 to 12
Ragweed, common	6 to 12
Ragweed, giant	6 to 12
Velvetleaf	6 to 12
Pennsylvania smartweed	6 to 9
Broadleaf signalgrass	6 to 8
Junglerice	6 to 8
Henbit	Up to 6
Sicklepod	4 to 6
Red rice	4 to 5
Hemp sesbania	3 to 4
Morningglory sp.	3 to 4
Prickly sida, Teaweed	3 to 4

Rate per acre: 48 fluid ounces

<u>WEED</u>	<u>WEED HEIGHT (INCHES)</u>
Hemp sesbania	5 to 9
Morningglory sp.	5 to 9
Prickly sida, Teaweed	4 to 6

NOTE:

Non-ionic Surfactant: Use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient.

Ammonium Sulfate: This additive may improve performance of Glyphosate 41% under environmental stress conditions. Ammonium sulfate is most effective under cool conditions or under mild moisture stress conditions when weeds are not growing as rapidly. Refer to the "ADDITIVES" section of the label booklet for use instructions for ammonium sulfate.

Sequential Applications: Some weeds with multiple germination times or suppressed (stunted) weeds may require a second application of this product. The second application should be made after some regrowth has occurred.

APPLICATION RATES – PERENNIAL WEEDS:

Glyphosate 41% at 16 to 64 oz per acre (single or sequential applications) will control or burndown the following perennial weeds: Canada thistle, common milkweed, field bindweed,

6/6

hemp dogbane, nutsedge, quackgrass, rhizome johnsongrass, tall fescue, trumpetcreeper, swamp smartweed, and wirestem muhly. For some perennial species, repeat applications may be required to eliminate crop competition throughout the growing season.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

4/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement

**Albaugh
GLYPHOSATE 41%**

EPA Reg No. 42750-xxx



HOODED SPRAYERS IN GRAIN SORGHUM (MILO)

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES. BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product may be used through hooded sprayers for weed control between the rows of grain sorghum (milo). Only hooded sprayers that completely enclose the spray pattern may be used.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way.

If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

When applying to milo that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 1 quart of this product per acre per application.
- Milo must be at least 12 inches tall, measured without extending leaves. Treat before milo sends tillers between the drill rows. If such tillers are contacted with the spray solution, the main plant may be killed.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 mph.
- Maximum wind speed: 10 mph.
- Use low-drift nozzles.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

For specific rates of application and instructions for control of various annual and perennial weeds, see the Weed Control tables in the label booklet for Glyphosate 41%.

Do not graze or feed milo forage or fodder following applications of this product through hooded sprayers.

Do not apply more than 3 quarts of this product per acre per year for hooded sprayer applications.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

Label as used in this supplemental labeling refers to the label book for Glyphosate 41% and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



HOODED SPRAYERS IN CORN

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way.

If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 1 quart of this product per acre per application.
- Corn must be at least 12 inches tall, measured without extending leaves.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 mph.
- Maximum wind speed: 10 mph.
- Use low-drift nozzles.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

For specific rates of application and instructions for control of various annual and perennial weeds, see the Weed Control tables in the label booklet for Glyphosate 41%.

Do not graze or feed corn forage or fodder following applications of this product through hooded sprayers.

Do not apply more than 3 quarts of this product per acre per year for hooded sprayer applications.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



HOODED SPRAYERS IN SUGARCANE

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product may be used through hooded sprayers for weed control between the rows of sugarcane.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of the label booklet for Glyphosate 41%.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding desirable vegetation from the spray solution. When applying to sugarcane that is grown on raised beds, ensure that the hood is designed to completely enclose the spray. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Hooded sprayers which apply the herbicide solution as a spray band should be calibrated on a broadcast equivalent rate and volume basis. Use the following equation to convert from a broadcast rate per acre to a band rate per acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Herbicide Broadcast RATE per acre} = \text{Herbicide Band RATE per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME of solution per acre} = \text{Band VOLUME of solution per acre}$$

Use nozzles that provide uniform coverage within the treated area. The herbicide solution should contact the entire weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation.

Do not allow treated weeds to come into contact with desirable vegetation.

Droplets, mist, foam, or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Minimize the potential for spray particles to escape from under the hood by operating the sprayer at appropriate ground speeds, nozzle pressures, and wind speeds. Operation on rough or sloping ground may result in spray particles escaping from the hood.

Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

11


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



FOR CONTROL OF VOLUNTEER GRAIN SORGHUM (MILO)

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product may be used to control volunteer grain sorghum (milo). Apply the following rates in 3 to 10 gallons of water per acre with ground equipment and 3 to 5 gallons of water per acre with aerial equipment:

<u>Rates</u>	<u>Heights</u>
12 fl oz/A	Up to 6 inches
16 fl oz/A	Up to 12 inches
24 fl oz/A	Up to 18 inches

Allow at least 3 days after application before tillage.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



**FOR DISTRIBUTION AND USE ONLY WITHIN TEXAS AND CHAVES, DONA ANA, EDDY
AND LUNA COUNTIES IN NEW MEXICO.
FOR PREHARVEST APPLICATIONS TO COTTON**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

When applied as directed under the conditions described, this product controls annual and perennial weeds listed in the label booklet for Glyphosate 41% prior to the harvest of cotton.

This product may be applied using either aerial or ground spray equipment. For aerial applications, refer to the "APPLICATION EQUIPMENT AND TECHNIQUES" and "AERIAL EQUIPMENT" sections of the label booklet.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT IN EAST AND CENTRAL TEXAS AND IN NEW MEXICO OR 2 QUARTS PER ACRE IN WEST TEXAS* WHEN MAKING APPLICATIONS BY AIR.

NOTE DO NOT APPLY TO CROPS GROWN FOR SEED.

Allow a minimum of 7 days between application and harvest. Do not feed or graze treated cotton forage or hay following preharvest applications.

WEED CONTROL

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of the label booklet.

TANK MIXTURES

This product may be tank-mixed with the following harvest-aid products or combination of harvest-aid products:

DEF™ 6	Folex™
Prep™	Dropp™

For application guidelines, precautions and use rates refer to the DEF, Folex, Prep, and Dropp labels.

HARVEST AID

At rates of 24 ounces to 2 quarts per acre and when tank-mixed with Def, Folex or Dropp, this product may provide enhancement of cotton leaf drop and regrowth inhibition. The degree of leaf drop or regrowth inhibition achieved is strongly affected by plant and weather factors.

TIMING OF APPLICATIONS

Apply this product after 60 percent of the cotton bolls have opened, or when there are sufficient mature bolls to produce the desired yields. Care must be taken not to apply this product before a sufficient number of bolls have matured since further boll maturation and development will most likely NOT occur after application.

A boll is matured when it is too hard to be dented when squeezed between thumb and finger, too hard to be sliced with a sharp knife, and when the seed coat becomes light brown in color.

In the Texas counties listed below, apply this product at any time after October 1 since the boll maturation process has essentially stopped. Prior to October 1, follow the application instructions listed above. For optimum control of whiteweed, make applications two weeks prior to frost.

Armstrong, Bailey, Borden, Briscoe, Castro, Childress, Cochran, Collingsworth, Crosby, Dawson, Deaf Smith, Dickens, Donley, Floyd, Gaines, Garza, Hale, Hall, Hockley, Kent, Lamb, Lubbock, Lynn, Motley, Parmer, Randall, Scurry, Swisher, Terry and Yoakum.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

*West Texas as used in this label is defined as the area bordered by Hwy. 86 on the north, Hwy. 287 on the northwest, Hwy. 283 on the east, Hwy. 190 on the south, Hwy. 285 on the southwest, and the Texas-New Mexico border on the west.

™Def is a trademark of Mobay Chemical Corporation.

™Folex and Prep are trademarks of Rhone-Poulenc, Inc.

™Dropp is a trademark of NOR-AM Chemical Company.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/3


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



**FOR DISTRIBUTION AND USE WITHIN DELAWARE, MARYLAND, NEW JERSEY,
PENNSYLVANIA AND VIRGINIA.**

**APPLICATIONS WITH FLUID FERTILIZERS FOR THE CONTROL OF ANNUAL AND
PERENNIAL WEEDS PRIOR TO THE EMERGENCE OF CORN, COTTON, SORGHUM,
SOYBEANS AND WHEAT.**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This supplemental label and the label booklet for Glyphosate 41% must be in the possession of the user at the time of application.

**AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE
PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.**

See the label booklet for Glyphosate 41% for essential product performance information.

**THIS PRODUCT MAY BE APPLIED IN WATER, CLEAR LIQUID NITROGEN FERTILIZER
AND CLEAR LIQUID COMPLETE-ANALYSIS FERTILIZER CARRIERS.**

USE WITH LIQUID FERTILIZERS

Glyphosate 41% may be applied in clear liquid nitrogen fertilizers and clear liquid complete-analysis fertilizers. The Glyphosate 41% tank mixtures listed in the "CONSERVATION TILLAGE, MINIMUM TILLAGE AND NO-TILL SYSTEMS" of the Glyphosate 41% label booklet may also be applied in these fertilizer carriers.

Do not use Glyphosate 41% with suspension-type liquid fertilizers.

When used with clear liquid fertilizers, this product will provide control of annual weeds and suppression of perennial weeds listed in the label booklet for Glyphosate 41% and in this supplemental labeling. Depending on the weed species, fertilizer composition, application

method and climatic conditions, rates of Glyphosate 41% at the higher end of the rate range may result in faster burndown and control of perennial weeds.

Add a nonionic surfactant at 0.5 to 1 percent by volume of spray solution.

The addition of 2 percent ammonium sulfate by weight or 17 pounds per 100 gallons of spray solution may increase the performance of this product.

The maximum rate for perennial weeds is 5 quarts or 160 fluid ounces per acre.

The maximum rate for annual weeds is 1.5 quarts or 48 fluid ounces per acre.

The combined total of all treatments must not exceed 8 quarts of this product per acre per year.

Minimum rates of Glyphosate 41% to use when liquid fertilizer carriers for all labeled weeds are shown in the tables below.

For control of spurred anoda, Virginia copperleaf, cotton, eveningprimrose, Carolina geranium, morningglory, hemp sesbania, Sicklepod, Teaweed or velvetleaf, make applications when the weeds are less than 6 inches tall.

USE RATES WITH CLEAR LIQUID NITROGEN FERTILIZERS

FERTILIZER VOLUME (Gallons per acre)	MAXIMUM WEED HEIGHT/LENGTH (Inches)		
	0 to 5"	6 to 10"	11 to 15"
	(Glyphosate 41% Quarts per Acre)		
25	0.5 to 0.75	0.75 to 1	1 to 1.5
50	1 to 1.5	1.5 to 2	2 to 3
75	1.5 to 2.25	2.25 to 3	3 to 4.5
100	2 to 3	3 to 4	*

*Not recommended

USE RATES WITH CLEAR LIQUID COMPLETE-ANALYSIS FERTILIZERS

FERTILIZER VOLUME (Gallons per acre)	FERTILIZER TYPE	MAXIMUM WEED HEIGHT/LENGTH (Inches)		
	N-P-K	0 to 5"	6 to 10"	11 to 15"
(Glyphosate 41% Quarts per Acre)				
25	N-0-0	0.5 to 0.75	0.75 to 1	1 to 1.5
	N-5-5	0.75 to 1	1 to 1.25	1.25 to 2
	N-10-10	0.75 to 1.25	1.25 to 1.5	1.5 to 2.5
	N-15-15	1 to 1.5	1.5 to 1.75	1.75 to 2.75
50	N-0-0	1 to 1.5	1.5 to 2	2 to 3
	N-5-5	1.25 to 2	2 to 2.5	2.5 to 3.75
	N-10-10	1.5 to 2.25	2.25 to 3	3 to 4.5
	N-15-15	1.75 to 2.75	2.75 to 3.5	3.5 to 4.75
75	N-0-0	1.5 to 2.25	2.25 to 3	3 to 4.5
	N-5-5	2 to 3	3 to 3.75	*
	N-10-10	2.25 to 3.5	3.5 to 4.5	*
	N-15-15	2.75 to 4	4 to 4.75	*
100	N-0-0	2 to 3	3 to 4	*
	N-5-5	2.5 to 3.75	*	*
	N-10-10	3 to 4.5	*	*
	N-15-15	*	*	*

*Not recommended.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



FOR CONTROL OF EMERGED WEEDS PRIOR TO THE PLANTING OR EMERGENCE OF CORN, COTTON, RICE, SORGHUM AND SOYBEANS; PRIOR TO THE HARVEST OF COTTON AND SOYBEANS; AND FOLLOWING THE HARVEST OF ANY CROP* IN THE FALL VIA AERIAL APPLICATIONS

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

*Do not plant subsequent crops other than those on the label for 30 days following application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS OR TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product controls annual and perennial weeds listed in the label booklet for Glyphosate 41% prior to the planting or emergence of corn, cotton, rice, sorghum and soybeans; prior to the harvest of cotton and soybeans; and following the harvest of any crop in the fall via aerial applications. See the "WEEDS CONTROLLED" and the "PREHARVEST APPLICATIONS" sections of the Glyphosate 41% label booklet for specific application rates and timing instructions.

Do not plant subsequent crops other than those on the label for 30 days following application.

Aerial applications of this product may be made in fallow systems and in conventional, reduced and zero tillage systems.

2/2

For applications via aerial equipment, use the recommended rates of this product in 3 to 10 gallons of water per acre. **Do not exceed a rate of 3 quarts per acre.**

AVOID DRIFT – DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser velocities, will allow spray drift to occur.

Read the "AERIAL EQUIPMENT" section of the Glyphosate 41% label booklet for additional precautions and instructions.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

4/7


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



GLYPHOSATE 41% FOR FORESTRY AND UTILITY RIGHTS-OF-WAY.

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of application.

APPLICATION EQUIPMENT AND TECHNIQUES

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

AERIAL EQUIPMENT

This product is recommended for aerial application in forestry sites by helicopter only. Use the recommended rates of this product in 5 to 30 gallons of spray solution per acre as a broadcast spray. (This product plus oil and surfactant mixtures may not be applied by air in California.)

AVOID DRIFT: DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT; DRIFT WILL CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine droplets.

Drift control additives may be used for forestry site preparation and utility rights-of-way applications. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label. The use of a drift control

2/17

agent for conifer and herbaceous release applications may result in conifer injury and is not recommended.

GROUND BROADCAST EQUIPMENT

This product is recommended for broadcast applications using suitable ground equipment in forestry sites, utility sites and utility rights-of-way. Use the recommended rates of this product plus surfactant in 10 to 60 gallons of clean water per acre as a broadcast spray. Check for even spray distribution throughout the spray pattern.

BACKPACK, HANDGUN OR MISTBLOWER EQUIPMENT

This product is recommended for application through backpack, handgun or hand-held mistblower equipment.

For spray-to-wet applications, coverage should be uniform and complete, but not to the point of runoff.

) This product may be used for low volume directed sprays for spot treatment of trees and brush. Coverage should be uniform with at least 50 percent of the foliage contacted. Coverage in the top one-half of the plant is important for best results. It is most effective in areas where there is a low density of undesirable trees or brush. If a straight stream nozzle is used, start the application at the top of the targeted vegetation and spray from top to bottom in a lateral zig-zag motion. For flat fan and cone nozzles and with mist blowers, mist the application over the foliage of the targeted vegetation. Small, open branched trees need only be treated from one side. If the foliage is thick or there are multiple root sprouts, applications must be made from several sides to ensure adequate spray coverage.

FORESTRY SITE PREPARATION AND UTILITY RIGHTS-OF-WAY

This product is recommended for the control or partial control of woody brush, trees and herbaceous weeds. This product is labeled for use in forestry and utility sites, including utility rights-of-way. This product is also recommended for use in maintaining logging roads.

) In forestry, this product is recommended for use in site preparation prior to planting any tree species, including Christmas trees and silvicultural nursery sites.

In utility rights-of-way, this product is recommended for use along power, pipeline, telephone and in other utility sites such as substations, etc.

APPLICATION RATES AND TIMING

<u>APPLICATION</u>	<u>GLYPHOSATE 41%</u>	<u>SPRAY VOLUME (GAL/A)</u>
<u>BROADCAST</u>		
Aerial	2 to 5 qts/a	5 to 30
Ground	2 to 5 qts/a	10 to 60
<u>SPRAY-TO-WET</u>		
Handgun, Backpack, Mistblower	¾ to 2% by volume	Spray-to-wet
<u>LOW VOLUME DIRECTED SPRAY</u>		
Handgun, Backpack, Mistblower	5% to 10% by volume	Partial coverage*

*See the BACKPACK, HANDGUN OR MISTBLOWER EQUIPMENT section of this label for application instructions.

In forestry site preparation and utility rights-of-way applications, this product requires use with a nonionic surfactant. Use a nonionic surfactant with greater than 50 percent active ingredient and labeled for use with herbicides. Use of this product without surfactant will result in reduced performance. See the "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" section of the label booklet for this product for more information.

Mix 2 or more quarts of the nonionic surfactant per 100 gallons of spray solution (0.5 percent or more by spray volume). Use of surfactant concentrations greater than 1.5 percent by spray volume with handgun applications or 2.5 percent by spray volume with broadcast applications is not recommended.

Use higher rates of this product within the recommended range for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the recommended range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product within the recommended range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

This product has no herbicidal or residual activity in the soil. Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

TANK MIXTURES

Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of both products used. Use according to the most restrictive label directions for each product in the mixture. Any recommended rate of this product may be used in a tank mix.

NOTE: For forestry site preparation, make sure the tank mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

4/7

<u>PRODUCT</u>	<u>BROADCAST RATE</u>	<u>USE SITE</u>
Arsenal™ Applicators Concentrate	2 to 16 fluid ounces per acre	Forestry site preparation
Oust™	1 to 4 ounces per acre	Forestry site preparation
Garlon™ 3A*, Garlon 4	1 to 4 quarts per acre	Forestry site preparation, Utility sites
Arsenal 2 WSL	2 to 32 fluid ounces per acre	Utility sites

SPRAY-TO-WET RATES

Arsenal™ Applicators Concentrate	1/32 to 1/2 % by volume	Forestry site preparation
Arsenal 2 WSL	1/32 to 1/2 % by volume	Utility sites

LOW VOLUME DIRECTED

SPRAY RATES

Arsenal™ Applicators Concentrate	1/8 to 1/2 % by volume	Forestry site preparation
Arsenal 2 WSL	1/8 to 1/2 % by volume	Utility sites

*Insure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray compatibility problems.

™Arsenal is a trademark of American Cyanamid Company.

™Oust is a trademark of E.I. du Pont de Nemours and Company.

™Garlon is a trademark of DowElanco Products Company.

FORESTRY CONIFER AND HARDWOOD RELEASE

DIRECTED SPRAY

In forestry conifer and hardwood sites, including Christmas tree plantations and silvicultural nurseries, use a 2 percent spray solution as a spray-to-wet application for the control of undesirable woody brush and trees. To control herbaceous weeds, use a 1 to 2 percent solution. Avoid contact of spray, drift or mist with foliage, green bark or non-woody surface roots of desirable species.

Mix 2 to 6 quarts of a nonionic surfactant per 100 gallons of spray solution (0.5 to 1.5 percent by spray volume). Use a surfactant with greater than 50 percent active ingredient.

BROADCAST SPRAY

Except where specifically recommended below, use only where conifers have been established for more than one year.

APPLICATION MUST BE MADE AFTER FORMATION OF FINAL CONIFER RESTING BUDS IN THE FALL OR PRIOR TO INITIATION OF BUD SWELLING IN THE SPRING.

Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied. Damage can be accentuated if applications are made when conifers are actively growing, or are under stress from drought, floodwater, insects or diseases.

This product may require use with a surfactant. Unless otherwise recommended in this section of this label, use Entry™ II surfactant at 5 to 15 fluid ounces per acre.

For release of the following conifer species, outside the southeastern United States:

Douglas fir

Pseudotsuga menziesii

Fir

Abies spp.

Hemlock**

Tsuga spp.

Pines*

Pinus spp.

Redwood, California**

Sequoia spp.

Spruce

Picea spp.

*Includes all species except loblolly pine, longleaf pine, shortleaf pine or slash pine.

**Use of additional surfactant is not recommended for release of hemlock species or California redwood. In mixed conifer stands, injury to these species may result if a surfactant is used.

Apply 1 to 2 quarts of this product per acre as a broadcast spray.

NOTE: For release of Douglas fir growing in the coastal range in Washington and Oregon at elevations above 1500 feet, Entry II surfactant may be used at rates up to 10 fluid ounces per acre. At elevations below 1500 feet in this area, additional surfactant is not recommended.

In Maine, up to 3 quarts per acre of this product may be used for the control of difficult species.

OUST TANK MIXTURES – To release jack pine, red pine, white pine, and white spruce, apply 1 to 2 quarts of this product with 1 to 3 ounces of Oust per acre. Make applications to actively growing weeds as a broadcast spray over the top of established conifers. Applications at these rates should be made after formation of conifer resting buds in the late summer or fall.

ARSENAL APPLICATORS CONCENTRATE TANK MIXTURES – This product may be tank mixed with Arsenal Applicators Concentrate for release of Douglas fir. Use 1 to 1-½ quarts of this product tank mixed with 2 to 6 fluid ounces of Arsenal per acre.

For release of the following conifer species in the southeastern United States:

Loblolly pine

Pinus taeda

Eastern white pine

Pinus strobes

Shortleaf pine

Pinus echinate

Slash pine

Pinus elliotii

Virginia pine

Pinus virginiana

Apply 1.5 to 2.5 quarts of this product per acre as a broadcast spray during late summer or early fall after the conifers have hardened off.

ARSENAL APPLICATORS CONCENTRATE TANK MIXTURES – Apply 1 to 2 quarts of this product with 2 to 16 fluid ounces of Arsenal Applicators Concentrate per acre as a broadcast spray for conifer release. Use only on conifer species that are labeled for over-the-top sprays

6/7

for both products. Use the higher recommended rates for dense, tough-to-control woody brush and trees.

Read and carefully observe the label claims, cautionary statements and all information on the labels of each product used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

HERBACEOUS RELEASE

When applied as directed, this product plus listed residual herbicides provides postemergence control of the annual weeds and control or suppression of the perennial weeds listed in this label, and residual control of the weeds listed in the residual herbicide label. Make applications to actively growing weeds as a broadcast spray over the top of labeled conifers. The use of additional surfactant is not recommended for these applications.

OUST TANK MIXTURES – To release loblolly pines, apply 16 to 24 fluid ounces of this product, plus 2 to 4 ounces of Oust per acre.

To release slash pines, apply 12 to 16 fluid ounces of this product, plus 2 to 4 ounces of Oust per acre.

Applications can be made over newly planted pines after the emergence of herbaceous weeds in the spring or early summer. Best results are obtained from applications made in May and June.

Weed control may be reduced if water volumes exceed 25 gallons per acre for these treatments.

ATRAZINE TANK MIXTURES – To release Douglas fir, apply 1 quart of this product, plus 4 pounds a.i. of atrazine per acre. Apply only over Douglas fir that has been established for at least one full growing season. Apply in the early spring, usually mid-March through early April. Injury will occur if applications are made after bud swell in the Spring.

Always read and follow the manufacturers label recommendations for all herbicides used.

INJECTION AND CUT STUMP APPLICATIONS

Woody brush and trees may be controlled using injection or cut stump applications of this product.

INJECTION APPLICATIONS

Apply the equivalent of 1 mL of this product per each 2 inches of trunk diameter. This is best achieved by applying a 25 to 100 percent concentration of this material either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In these species, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, application should be made during periods of active growth and after full leaf expansion.

CUT STUMP TREATMENTS

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in

7/7

reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, this product will CONTROL, PARTIALLY CONTROL or SUPPRESS most woody brush and tree species, some of which are listed below:

Alder

Alnus spp.

Coyotebrush

Baccharis consanguinea

Dogwood

Cornus spp.

Eucalyptus

Eucalyptus spp.

Hickory

Carya spp.

Madrone

Arbutus menziesii

Maple

Acer spp.

Oak

Quercus spp.

Poplar

Populus spp.

Reed, giant

Arundo donax

Saltcedar

Tamarisk spp.

Sweetgum

Liquidambar styraciflua

Sycamore

Platanus occidentalis

Tan Oak

Lithocarpus densiflorus

Willow

Salix spp.

Read and carefully observe the label directions, cautionary statements and all information on the labels of each product used. Additional precautionary statements are made in these labels. Use according to the most restrictive label directions for each product in the mixture.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 5002

NEXT

LABEL

42750-60

4-25-2001

1/1


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



PREHARVEST APPLICATIONS IN GRAIN SORGHUM (MILO)

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product may be used prior to harvest of grain sorghum (milo) to control annual and perennial weeds listed on the label booklet for Glyphosate 41% herbicide.

Apply up to 2 quarts of this product per acre.

Make applications at 30% grain moisture or less and at least 7 days prior to harvest.

This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for instructions.

Note: It is not recommended that grain sorghum grown for seed be treated because a reduction in germination or vigor may occur.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% container labels and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



**BROADCAST APPLICATIONS FOR WEED CONTROL IN CHRISTMAS TREE
PLANTATIONS IN THE STATES OF WASHINGTON AND OREGON ONLY**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

NOTE: IF IMPROPERLY APPLIED, THIS PRODUCT HAS THE POTENTIAL TO CAUSE SEVERE CHRISTMAS TREE INJURY. FOLLOW ALL LABELED DIRECTIONS.

This product may be applied as a broadcast spray over established Christmas trees. Ensure that adequate buffers are maintained to prevent drift onto nearby desirable crops or vegetation. Read the entire "APPLICATION EQUIPMENT AND TECHNIQUES" section of the Glyphosate 41% label booklet for additional application precautions.

This application is approved for the following Christmas tree species:

Douglas fir	(<i>Pseudotsuga menziesii</i>)
Fir species	(<i>Abies spp.</i>)
Spruce species	(<i>Picea spp.</i>)

Applications may be made only after trees have completed at least a full growing season since planting or transplanting. Applications should not be made within 1 full year prior to tree harvest.

Applications may only be made in the fall after the formation of final conifer resting buds. Final resting buds must be fully hardened and in the dormant stage. Applications made at any other time may result in unacceptable Christmas tree injury.

2/2

Avoid spray pattern overlap, as injury may occur.

Apply 1 quart of this product per acre in 5 to 30 gallons of water per acre.

NOTE: DO NOT ADD SURFACTANTS, ADDITIVES CONTAINING SURFACTANTS, OR ANY OTHER ADDITIVES TO THIS PRODUCT OR SEVERE CHRISTMAS TREE INJURY MAY RESULT.

This product may be used at rates from 1 to 2 quarts per acre in some areas. Consult your local Albaugh, Inc. representative for specific recommendations if you require rates greater than 1 quart per acre.

Drift control additives may increase Christmas tree injury and their use is not recommended.

The use of other herbicides tank mixed with Glyphosate 41% is not recommended since severe Christmas tree injury may result.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement:

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



FOR AERIAL APPLICATION IN ARKANSAS ONLY

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING" sections of the Glyphosate 41% label booklet for essential product performance information.

USE DIRECTIONS

AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the recommended rate of this product in 3 to 15 gallons of water per acre. ✓

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are recommended.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety

The distance of the outermost nozzles on the boom must not exceed 75% of the length of the wingspan rotor. In many cases reducing this distance to 65% of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing airflow on rotary winged aircraft. Avoid the use of nozzles with wide-angle discharge.

Do not apply this product when wind speeds are in excess of 10 miles per hour.

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/5

SUPPLEMENTAL LABELING


READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% container labels and this supplement

WARNING. USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-~~60~~ ^{AN}

Roundup Ready® is a registered trademark of the Monsanto Company.



**FOR POSTEMERGENCE APPLICATIONS TO CORN
WITH THE ROUNDUP READY® GENE**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

GENERAL INFORMATION

ROUNDUP READY® CORN VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE CORN VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT. IT IS UNLAWFUL TO SELL OR PLANT SAVED SEED.

CORN WITH THE ROUNDUP READY® GENE MAY ONLY BE USED FOR PLANTING A COMMERCIAL CROP IN A SINGLE SEASON. SEED MAY NOT BE SAVED FOR REPLANTING AND SAVED SEED MAY NOT BE SUPPLIED TO OTHERS FOR REPLANTING. ALBAUGH, INC. DOES NOT WARRANT THE SAFETY OR PERFORMANCE OF THIS PRODUCT WHEN USED ON "BROWN BAG" OR FARMER-MADE SEED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

2/3

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, OTHER THAN CORN WITH THE ROUNDUP READY GENE, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% for essential product performance information.

GENERAL INFORMATION

ALBAUGH, INC. RECOMMENDS USE OF THIS PRODUCT ONLY ON CORN HYBRIDS DESIGNATED AS CONTAINING THE ROUNDUP READY® GENE.

- Applying this product to corn hybrids which are not designated as Roundup Ready® will result in severe crop injury and yield loss.
- The Roundup Ready designation indicates that the corn contains a patented gene which provides tolerance to this herbicide. Information on Roundup Ready corn may be obtained from your seed supplier.

Application Instructions

This product may be applied postemergence to Roundup Ready corn from emergence through the V8 (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first. Single in-crop applications of this product are not to exceed 1 quart per acre.

Sequential in-crop applications of this product from emergence through the V8 stage or 30 inches must not exceed 2 quarts per acre per growing season.

Maximum Yearly Rates Allowed

Preplant: Maximum amount of this product which can be applied prior to crop emergence is 5 quarts per acre.

In-crop: Maximum combined total of multiple in-crop applications from emergence through the V8 stage or 30 inches is 2 quarts per acre.

Preharvest: Maximum amount of this product that can be applied after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest is 1 quart per acre.

Cropping Season: Combined total per year for all applications may not exceed 8 quarts per acre.

When applied as directed, this product controls labeled annual grass and broadleaf weeds in Roundup Ready corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. Applications should be made to actively growing weeds before they reach the maximum size listed in the "WEEDS CONTROLLED" section of the label booklet for Glyphosate 41%. Refer to the "MIXING" section of the label booklet for proper use instructions.

3/5

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product under hard water conditions, drought conditions or when tank mixed with Bullet®, Micro-Tech® or Partner® Herbicides. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. The addition of other additives, including fertilizers and micronutrients are not recommended with this product since this may result in increased potential for crop injury.

Allow a minimum of 50 days between application of this product and harvest of corn forage and 7 days between application and harvest of corn grain. Allow a minimum of 10 days between in-crop applications of this product. **Do not graze, harvest or feed corn forage or silage following sequential in-crop applications of this product on Roundup Ready corn.** There are no rotational crop restrictions following applications of this product.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY® GENE.

THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

For ground applications: Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select correct nozzles and spray pressure to avoid spraying a fine mist. Check for even distribution of spray droplets.

For aerial applications: Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart per acre. See "WEEDS CONTROLLED" section on this label. AVOID DRIFT – DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Weeds Control Recommendations

Apply 24 to 32 ounces of Glyphosate 41% per acre for control of labeled grasses and broadleaf weeds in conventional and no-till corn production systems. Refer to the label booklet for Glyphosate 41% for rate recommendations for specific annual weeds. Glyphosate 41% applied at up to 1 quart per acre will control or suppress the growth of perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, Horsenettle, nutsedge, quackgrass, rhizome johnsongrass, Redvine, Trumpetcreeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the "PERENNIAL WEED RATE TABLE" of the label booklet for Glyphosate 41%.

Preemergence followed by Postemergence Weed Control Program

This product may be applied postemergence in-crop following any labeled preemergence herbicide applications. The post application of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop. A single in-crop application of this product at the recommended rate will provide control of emerged weeds listed on the label. This product may be applied postemergence to Roundup Ready corn from

emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing), whichever comes first.

Postemergence Only Weed Control Program

This product may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on the label. The postemergence application of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 24 to 32 fluid ounces per acre will control the labeled grasses and broadleaf weeds. This product may be applied postemergence to Roundup Ready corn from emergence through the V8 stage or until corn height reaches 30 inches (free standing), whichever comes first.

This product may be applied in tank mixing with a labeled rate of Harness®, Harness® Xtra, Harness® Xtra 5.6L, Micro-Tech®, Bullet®, Partner®, Permit® or Atrazine. Refer to the specific product label and observe all precautions and limitations on the label for all products used in tank mixtures, including application timing restrictions, soil restrictions, minimum recropping interval and rotational guidelines – the more restrictive requirements apply. Tank mixtures with other products may result in increased potential for crop injury and/or weed antagonism. Refer to the table below for height limitation for tank mix partner.

Tank Mix Partner	Maximum Height Of Corn For Application
Harness Harness Xtra Harness Xtra 5.6	11 inches
Bullet* Micro-Tech* Partner*	5 inches
Permit	24 inches
Atrazine	12 inches

*Bullet, Micro-Tech and Partner are not registered for use as a postemergence application in Texas.

NOTE: Non-ionic Surfactant: Non-ionic surfactants which are labeled for use with postemergence herbicides may be used. When using additional surfactant, use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient.

The addition of certain surfactants to this product may result in some crop response including leaf necrosis, leaf chlorosis or leaf speckling due to the surfactant added to the spray mixture. Read and carefully observe cautionary statements and other information appearing on the surfactant label.

*Bullet, Harness, Micro-Tech and Partner are registered trademarks of Monsanto Company.

*Permit is a trademark of, and used under license from Nissan Chemical Industries, Ltd.

5/5

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

4/10


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% container labels and this supplement.

WARNING: USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



FOR POSTEMERGENCE APPLICATIONS TO SOYBEAN VARIETIES WITH THE ROUNDUP
READY® GENE

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

Roundup Ready® Gene and Roundup Ready® are registered trademarks of the Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

GENERAL INFORMATION

FOR POSTEMERGENCE APPLICATION ONLY ON SOYBEAN VARIETIES WHICH HAVE THE ROUNDUP READY® GENE

- 2/6
- Applying this product to soybean varieties which are not designated as Roundup Ready® will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Roundup Ready gene, since severe injury or destruction will result.
 - Roundup Ready varieties must be purchased from an authorized seed supplier. Crop safety and weed control performance are not warranted by Albaugh, Inc. when this product is used in conjunction with "brown bag" or "bin run" soybean seed saved from previous year's production and replanted.
 - The Roundup Ready designation indicates that the soybean contains a patented gene which provides tolerance to Albaugh's Glyphosate brand herbicides. Information on Roundup Ready soybeans may be obtained from your seed supplier.

APPLICATION INSTRUCTIONS

This product may be applied postemergence to Roundup Ready soybeans from the cracking stage through the full flowering stage.

Allow a minimum of 14 days between application and harvest of soybeans.

MAXIMUM ALLOWABLE YEARLY RATES

Preplant: Maximum amount of this product which can be applied prior to crop emergence is 5 quarts/A.

In-crop: Maximum combined total of single or multiple in-crop applications from cracking to flowering is 2 quarts/A.

Preharvest: Maximum amount of this product that can be applied after loss of green color in soybean pods until 14 days before harvest is 1 quart/A.

Cropping Season: Combined total per year for all applications may not exceed 8 quarts/A.

When applied as directed, this product will control labeled annual grasses and broadleaf weeds in Roundup Ready soybeans. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product.

There are no rotational crop restrictions following applications of this product.

For ground applications: Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use nozzles which provide a flat fan pattern. Check for even distribution of spray droplets.

For aerial applications: Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart of this product per acre. DO NOT APPLY DURING LOW LEVEL INVERSION CONDITIONS. WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

AERIAL APPLICATIONS ON ROUNDUP READY SOYBEANS MAY BE MADE ONLY IN THE FOLLOWING STATES: ALABAMA, ARKANSAS, FLORIDA, GEORGIA, KANSAS,

3/14

LOUISIANA, MISSISSIPPI, MISSOURI (BOOT-HEEL ONLY), NORTH CAROLINA,
OKLAHOMA, SOUTH CAROLINA. TENNESSEE AND TEXAS.

ANNUAL WEED RATE TABLES

The following rate recommendations will provide control of labeled grasses and broadleaf weeds in conventional and no-till soybean production systems. Refer to the label booklet for rate recommendations for specific annual weeds.

Tank mixtures with other herbicides are not recommended due to potential for crop injury and/or weed antagonism and to rotational crop restrictions of the tank mix partner.

This product may be used up to 64 fluid ounces per acre in any single application for control of annual weeds, where heavy weed densities exist.

NOTE: In no-till and stale seedbed systems, always burn down existing weeds before soybeans emerge. Apply a preplant burn-down treatment of 16-64 fluid ounces per acre of this product.

MIDWEST RECOMMENDATIONS

Narrow row or drilled soybeans: A single in-crop application of this product will provide effective control of labeled weeds.

Weed Height (inches)	Rate (fluid oz/A)
3-6	24
6-12	32
12-18	48

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this product at 16-32 fluid ounces per acre may be necessary to control late flushes of weeds.

Wide row soybeans: A single in-crop application of this product will provide effective control of initial stand of labeled weeds. If new flushes of weeds occur, they can be controlled by sequential applications of this product.

Initial Treatment

Weed Height (inches)	Rate (fluid oz/A)
3-6	24
6-12	32
12-18	48

Sequential Application*

Weed Height (inches)	Rate (fluid oz/A)
1-3	16
3-6	24
6-12	32

*Combined total application in-crop not to exceed 64 fluid ounces per acre.

4/16

For morningglory, black nightshade, groundcherry, and Pennsylvania smartweed apply:

Weed Height (inches)	Rate (fluid oz/A)
1-3	24
3-6	32
6-12	48

Giant ragweed. Apply 32 fluid oz/A when the weed is 8-12" tall to avoid the need for sequential application.

Some weeds, such as black nightshade, woolly cupgrass, shattercane, wild proso millet, burcucumber, and giant ragweed, with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 64 fluid ounces per acre.

MID-ATLANTIC/SOUTHEAST RECOMMENDATIONS

Narrow row or drilled soybeans: A single in-crop application of this product will provide effective control of labeled weeds.

Weed Height (inches)	Rate (fluid oz/A)
3-6	24
6-12	32
12-18	48

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this product at 16-32 fluid ounces per acre may be necessary to control late flushes of weeds.

Wide row soybeans: A single in-crop application of this product will provide effective control of initial stand of labeled weeds. If new flushes of weeds occur, they can be controlled by sequential applications of this product.

Initial Treatment

Weed Height (inches)	Rate (fluid oz/A)
3-6	24
6-12	32
12-18	48

57/4

Sequential Application*

Weed Height (inches)	Rate (fluid oz/A)
1-3	16
3-6	24
6-12	32

*Combined total application not to exceed 64 fluid ounces per acre.

Hemp sesbania: Apply 24 fl oz/A at 0-2" tall; 32 fl oz/A at 2-4" tall; 40 fl oz/A at 4-6" tall; 48 fl oz/A at 6-8" tall.

For morningglory, black nightshade, groundcherry, and Pennsylvania smartweed apply:

Weed Height (inches)	Rate (fluid oz/A)
1-3	24
3-6	32
6-12	48

Some weeds, such as black nightshade, broadleaf signalgrass, Texas panicum, burcucumber, and sicklepod, with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 64 fluid ounces per acre.

DELTA/MID-SOUTH RECOMMENDATIONS

Narrow row or drilled soybeans: An in-crop application of this product will provide effective control of initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds.

Wide row soybeans: An in-crop application of this product will provide effective control of initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds.

Initial Treatment

Weed Height (inches)	Rate (fluid oz/A)
1-3	24
3-6	32
6-12	48

6/6

Sequential Application*

Weed Height (inches)	Rate (fluid oz/A)
1-3	16
3-6	24
6-12	32

*Combined total application in-crop not to exceed 64 fluid ounces per acre.

Hemp sesbania: Apply 24 fl oz/A at 0-2" tall; 32 fl oz/A at 2-4" tall; 40 fl oz/A at 4-6" tall; 48 fl oz/A at 6-8" tall.

PERENNIAL WEEDS RATE RECOMMENDATIONS

A 32 to 64 fluid ounce per acre rate (single or sequential applications) of this product will control or suppress perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the "PERENNIAL WEED RATE TABLE" of the label booklet for Glyphosate 41%. For some perennial species, repeat application may be required to eliminate crop competition throughout the growing season.

NOTE: Non-ionic Surfactant: Use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient.

Ammonium Sulfate: Ammonium sulfate may be mixed with this product for applications to Roundup Ready soybeans. Refer to the "ADDITIVES" section of the label booklet for use instructions for ammonium sulfate.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

SUPPLEMENTAL LABELING


READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

WARNING: USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx

Roundup Ready® is a registered trademark of the Monsanto Company.



FOR COTTON WITH THE ROUNDUP READY® GENE
IN-CROP APPLICATIONS

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% for essential product performance information.

GENERAL INFORMATION

WARNING: ALBAUGH, INC. RECOMMENDS THIS PRODUCT FOR USE ONLY OVER-THE-TOP OF OR DIRECTED ONTO IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE ROUNDUP READY® GENE. SEVERE INJURY OR DEATH OF COTTON WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY® GENE ARE SPRAYED WITH THIS PRODUCT. AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR

24

ANY DESIRABLE PLANTS AND TREES, OTHER THAN CROPS WITH THE ROUNDUP READY® GENE, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

ROUNDUP READY® COTTON VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE COTTON VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT. IT IS UNLAWFUL TO SELL OR PLANT SAVED SEED.

COTTON WITH THE ROUNDUP READY® GENE MAY ONLY BE USED FOR PLANTING A COMMERCIAL CROP IN A SINGLE SEASON. SEED MAY NOT BE SAVED FOR REPLANTING AND SAVED SEED MAY NOT BE SUPPLIED TO OTHERS FOR REPLANTING. ALBAUGH, INC. DOES NOT WARRANT THE SAFETY OR PERFORMANCE OF THIS PRODUCT WHEN USED ON "BROWN BAG" OR FARMER-MADE SEED.

APPLICATION INSTRUCTIONS

This product will control many troublesome weeds with over-the-top directed, hooded sprayer, or preharvest applications in Roundup Ready cotton.

<u>Maximum Allowable Yearly Rates</u>	
1. Combined total per year for all applications	8 quarts/A
2. Preplant, Preemergence applications	5 quarts/A
3. Total in-crop applications from cracking to layby	4 quarts/A
4. Maximum preharvest application rate	2 quarts/A

For ground applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

For aerial applications apply this product in 3 to 15 gallons of water per acre.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY® GENE. Do not apply during low-level inversion conditions, when winds are gusty or under any other conditions which favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

There are no rotational crop restrictions following applications of this product.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to Roundup Ready cotton. Follow the cleaning procedures specified on the label of the product(s) previously used. Cotton is very sensitive to many herbicides at extremely low concentrations and care should be taken to thoroughly clean all equipment prior to use.

In addition to uses listed in the Glyphosate 41% label booklet, the following applications can be made:

Over-the-top applications: This product may be applied by aerial or ground application equipment postemergence to Roundup Ready cotton from the ground cracking stage until the four leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Over-the-top applications made after the four leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss. Any single over-the-top broadcast application should not exceed 1 quart per acre. No more than two over-the-top broadcast applications may be made from crop emergence through the four leaf (node) stage of development. Sequential over-the-top applications of this product must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

NOTE: Always plant into a weed free seedbed. In no-till and stale seedbed systems, always burn down existing weeds before cotton emerges. Apply a preplant burndown treatment of 16-48 fluid ounces per acre of this product.

Post-directed or hooded applications: This product may be applied using precision post-directed or hooded sprayers to Roundup Ready cotton through layby. Be especially careful to minimize contact of the spray with cotton leaves. At this stage, post-directed equipment should be used which directs the spray to the base of the cotton plants. Place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row. For best results, make applications while weeds are small (less than 3 inches). Minimize spray drift onto the leaves of the cotton plants by maintaining low spray pressure (less than 30 PSI). Applications that contact the cotton leaves may result in boll loss, delayed maturity and/or yield loss. Any single post-directed application should not exceed 1 quart per acre of this product. No more than two applications should be made from the fifth leaf through layby. Sequential in-crop applications of this product must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

Salvage Treatment: This treatment may be used after the four leaf stage of development and should only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top application or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. **NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER GROWING SEASON.**

Weeds controlled: For specific rates of application and instructions for control of various annual and perennial weeds, refer to "ANNUAL WEEDS RATE TABLE" section of the label booklet. Glyphosate 41% applied at 1 quart per acre will burndown or suppress the growth of the following perennial weeds and reduce crop competition: yellow and purple nutsedge, rhizome johnsongrass, common bermudagrass, silverleaf nightshade, trumpet creeper, and Redvine. Fall preharvest applications may be required for control of these perennial weeds.

Tank mixtures with other herbicides may result in reduced weed control or crop injury and are not recommended for over-the-top applications of this product.

Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of this product for control.

Preharvest applications: This product may be applied for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready cotton after 20% boll crack. Do not apply more than 2 quarts of this product per acre for preharvest applications. Allow a minimum of 7 days between application and harvest. **THE USE OF ADDITIVES FOR PREHARVEST APPLICATION OF GLYPHOSATE 41% TO ROUNDUP READY COTTON IS PROHIBITED.**

4/4

For specific recommendations refer to the "COTTON" section of the Glyphosate 41% label booklet. NOTE: Glyphosate 41% will not enhance the performance of harvest aids when applied to Roundup Ready cotton. DO NOT APPLY GLYPHOSATE 41% TO CROPS GROWN FOR SEED.

NOTE: Non-ionic Surfactant: Non-ionic surfactants which are labeled for use with postemergence herbicides may be used. When using additional surfactant, use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

Albaugh
GLYPHOSATE 41%

EPA Reg. No. 42750-xxx



FOR DISTRIBUTION AND USE ONLY WITHIN OHIO.
FOR CONTROL OF CANADA THISTLE

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This supplemental labeling and the label booklet for Glyphosate 41% must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION", "APPLICATION EQUIPMENT AND TECHNIQUES" and "WEEDS CONTROLLED" sections of the label booklet for Glyphosate 41% for essential product performance information.

FOR CONTROL:

Apply 1 quart per acre of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre.

The addition of ammonium sulfate at the rate of 17 pounds per 100 gallons of spray solution is recommended for enhanced performance under stress conditions.

In the spring, apply to actively growing thistles when most are at or beyond the bud stage of growth.

After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth. Rosette size must be a minimum of 6 inches in diameter before treating.

Applications can be made in the fall as long as leaves are still green, and plants are actively growing at the time of application.

Allow 3 or more days after application before tillage. For best results, till the area within 14 days after application.

FOR SUPPRESSION WITH SPRING APPLICATIONS:

Apply 1 pint to 1 quart of this product per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Apply to actively growing plants that are at least 8 inches tall.

The addition of ammonium sulfate at the rate of 17 pounds per 100 gallons of spray solution is recommended for enhanced performance under stress conditions.

For improved burndown and to control other broadleaf weeds, mix this product with 0.5-pound a.i. 2,4-D per acre. Applications of 2,4-D must be made at least 7 days prior to planting corn and at least 30 days prior to planting soybeans.

Allow 3 or more days after application before tillage. A subsequent cultivation may enhance control.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

4/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



**FOR DISTRIBUTION AND USE ONLY WITHIN HAWAII.
FOR NONSELECTIVE WEED CONTROL IN AND AROUND PASSION FRUIT**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This supplemental labeling and the label booklet for Glyphosate 41% must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION", "APPLICATION EQUIPMENT AND TECHNIQUES" and "WEEDS CONTROLLED" sections of the label booklet for Glyphosate 41% for essential product performance information.

RECOMMENDATIONS

This product is recommended for weed control in and around established passion fruit, or for site preparation prior to planting or transplanting. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment.

When applying this product refer to the "Weeds Controlled" section of the label booklet for Glyphosate 41% for weeds controlled and specific rates to be used.

Allow a minimum of 14 days between last application and harvest.

Annual Weed Control – Apply 8 to 48 fluid ounces of this product per acre plus 0.5 to 1 percent nonionic surfactant in 3 to 10 gallons of water per acre as directed in the "Low-Volume Broadcast Application" section of the label booklet for Glyphosate 41%.

When using water volumes of 10 to 40 gallons per acre, apply 1 quart per acre if annual weeds are less than 6 inches tall. If weeds are greater than 6 inches tall, use 1.5 quarts of this product per acre.

Perennial Weed Control – Apply 3 to 5 quarts of this product per acre. Apply when weeds are actively growing and most have reached the early head or early bud stage of growth. Allow 7 or more days after application before mowing or tillage operations.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



**FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA.
FOR NONSELECTIVE CONTROL OF LISTED ANNUAL WEEDS IN SMALL GRAIN
CROPPING SYSTEMS.**

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.
This label must be in the possession of the user at the time of application.

ANNUAL WEED CONTROL

WEEDS CONTROLLED

Refer to the "RECOMMENDED RATES AND WEEDS CONTROLLED" section of the label booklet for Glyphosate 41% for rate recommendations and the annual weeds controlled.

APPLICATION INSTRUCTIONS

For ground applications, use 3 to 5 gallons of water per acre. For aerial applications, use 1 to 3 gallons of water per acre.

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this herbicide can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

NOTE

- To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.
- Adjust boom height on ground equipment to prevent streaked, overlapped or uneven applications. Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets.
- In aerial applications, do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.
- Ensure uniform application. Use appropriate marking devices when applying herbicides by air.
- Avoid spraying when weeds are subject to moisture stress, when dust is on foliage, or when straw canopy covers the weeds.
- Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residue of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF PART. LANDING GEAR ARE MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021

NEXT

LABEL

42750-60

4-25-2001

1/2


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% and this supplement.

**Albaugh
GLYPHOSATE 41%**

EPA Reg. No. 42750-xxx



GLYPHOSATE 41% COMBINATIONS WITH 2,4-D AMINE AND OUST™ FOR CONTROL OF TRUMPETCREEPER AND JOHNSONGRASS ON RAILROAD RIGHTS-OF-WAY.

Keep out of reach of children

CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

NONCROP USES

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of this label for essential product performance information.

See the following noncrop sections for specific recommended uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

NOTE: If spraying areas adjacent to desirable plants, use a shield made of cardboard, sheet metal or plywood while spraying to help prevent spray from contacting foliage of desirable plants.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

Glyphosate 41% does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

This product may be applied in noncrop sites as indicated in the "TANK MIXING AND APPLICATION INSTRUCTIONS" section unless otherwise directed.

2/2

RAILROAD RIGHTS-OF-WAY

Tank Mixture

Glyphosate 41% plus 2,4-D Amine

When applied as directed for noncrop uses, Glyphosate 41% when tank-mixed with 2,4-D amine, will provide burndown and control of Trumpet creeper in railroad rights-of-way sites. Apply 2 to 3 quarts of Glyphosate 41% with 1 to 2 pints of 2,4-D amine in 25 to 40 gallons of total spray solution per acre to actively growing Trumpet creeper. Application should be made any time from early postemergence to before a killing frost. Use the higher rates of these products when weed growth is heavy or dense.

Tank Mixture

Glyphosate 41% plus 2,4-D Amine plus Oust

When applied as directed for noncrop uses, Glyphosate 41% when tank-mixed with 2,4-D amine and Oust will provide burndown control of Johnson grass and trumpet creeper. Apply 2 to 3 quarts of Glyphosate 41% with 1 to 2 pints of 2,4-D amine plus 2 to 4 ounces of Oust in 25 to 40 gallons of total spray solution per acre. Application should be made any time from early postemergence to before a killing frost. Use the higher rates of these products when weed growth is heavy or dense.

TANK MIXING AND APPLICATION INSTRUCTIONS

Before using, refer to the individual product labels for precautionary statements.

Do not apply this tank mixture, drain or flush equipment on or near desirable trees or other plants, on areas where their roots may extend, or in locations where Oust or 2,4-D amine may be washed or moved into contact with their roots.

Fill the spray tank at least one-third full of clean water. Mix the recommended amount of Oust in a separate container with sufficient water to make a smooth slurry. Pour the slurry into the spray tank, fill spray tank with the required amount of 2,4-D amine and Glyphosate 41% and mix well before using. Maintain agitation until spraying is completed.

Before using, refer to individual product labels for specific cleaning instructions.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

™Oust is a trademark of E.I. duPont de Nemours & Company, Inc.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021