

## **PLEASE NOTE**

**This image contains more than one label  
approved for this product on this date.**


**SUPPLEMENTAL LABELING**

READ AND FOLLOW THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus, and this supplement.

**Albaugh**  
**GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx



---

**GLYPHOSATE 41% PLUS HERBICIDE AND TANK MIXTURES FOR NONCROP AREAS, INCLUDING RAILROAD RIGHTS-OF-WAY, UTILITY SUBSTATIONS, AIRPORTS, INDUSTRIAL PLANTS, ROADSIDES, STORAGE AREAS AND SIMILAR SITES.**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See the "GENERAL INFORMATION", "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" and "NONCROP USES" sections of the label booklet for Glyphosate 41% Plus for essential product information.

Do not allow spray mixtures of this herbicide to mist, drip, drift or splash onto desirable vegetation since injury or destruction may occur. Do not apply when wind or other conditions favor drift.

See the "WEEDS CONTROLLED" section of the Glyphosate 41% Plus label booklet for rate recommendations. For difficult to control species, where dense stands occur, or where conditions for control are not ideal, 5 to 10 quarts per acre of this product may be used for improved results.

**TANK MIXTURES**

This product provides control of the emerged weeds listed in the label booklet. When applied as a tank mixture, the following herbicides will provide preemergence and/or postemergence control of the weeds listed in the individual product labels.

The following list of products may be tank mixed with this product. Any recommended rate of this product may be used in a tank mixture with these products.

<u>Tank Mix Product</u>	<u>Rate per Acre</u>
Arsenal™	0.5 to 4 pints
Banvel	1 to 4 pints
2,4-D	0.5 to 1 pound
Garlon™ 3A	1 to 6 pints
Garlon 4	1 to 6 pints
Diuron	4 to 8 pounds
Diuron + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Diuron + Garlon 3A	4 to 10 pounds + 1 to 2 pints
Diuron + Garlon 4	4 to 10 pounds + 1 to 2 pints
Hyvar™ X	4 to 8 pounds
Hyvar X + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Hyvar X + Garlon 3A	4 to 8 pounds + 1 to 2 pints
Hyvar X + Garlon 4	4 to 8 pounds + 1 to 2 pints
Krovar™ I DF	4 to 6 pounds
Krovar I DF + 2,4-D	4 to 6 pounds + 0.5 to 1 pound
Krovar I DF + Garlon 3A	4 to 6 pounds + 1 to 2 pints
Krovar I DF + Garlon 4	4 to 6 pounds + 1 to 2 pints
Oust™	2 to 6 ounces
Oust + 2,4-D	2 to 6 ounces + 0.5 to 1 pound
Oust + Garlon 3A	2 to 6 ounces + 1 to 2 pints
Oust + Garlon 4	2 to 6 ounces + 1 to 2 pints
Spike™ 80W	2 to 5 pounds
Spike 80W + 2,4-D	2 to 5 pounds + 0.5 to 1 pound
Spike 80W + Garlon 3A	2 to 5 pounds + 1 to 2 pints
Spike 80W + Garlon 4	2 to 5 pounds + 1 to 2 pints

\*Arsenal is not approved for use in the state of California.

Refer to the individual product labels for specific noncrop sites, rates, carrier volumes and precautionary statements.

Read and carefully observe the label claims, cautionary statements, recommended use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Maintain good agitation at all times during the mixing process. Ensure that the tank mix products are well mixed with the spray solution before adding this product.

Mix only the quantity of spray solution which can be used during the same day. Tank mixtures allowed to stand overnight may result in reduced weed control.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

When used in combination as recommended by Albaugh, Inc., the liability of Albaugh, Inc. shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Albaugh product in such combination use.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus, and this supplement.

**Albaugh**  
**GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx



---

**FOR PREHARVEST APPLICATIONS IN ALFALFA**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

**RECOMMENDATIONS**

This product may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This product will control labeled annual and perennial weeds, including quackgrass, when applied prior to the harvest of alfalfa.

The treated crop and weeds can be harvested and fed to livestock. Allow a minimum of 36 hours between application and harvest.

Use up to 1 quart of this product per acre. Applications may be made at any time of the year. Make only one application to an existing stand of alfalfa per year.

For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments must be followed by deep tillage for complete control.

Do not use alfalfa grown for seed.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**




**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus, and this supplement.

**Albaugh**  
**GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx



---

**FOR WEED CONTROL PRIOR TO THE EMERGENCE OF SUNFLOWERS**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

**RECOMMENDATIONS**

This product controls annual and perennial weeds listed on this label prior to the emergence of sunflowers. Applications may be made before, during or after planting.

Apply up to 32 fluid ounces of this product per acre. Make only one preemergence application of this product per year.

Do not feed or graze sunflower forage following application of this product.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ AND FOLLOW THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus, and this supplement.

**Albaugh**  
**GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx



**GLYPHOSATE 41% PLUS FOR CONTROL OR SUPPRESSION OF *BROMUS* SPECIES AND MEDUSAHEAD FOUND IN INDUSTRIAL, RANGELAND AND PASTURE SITES IN COLORADO, IDAHO, IOWA, KANSAS, MONTANA, NEBRASKA, NORTH DAKOTA, OREGON, SOUTH DAKOTA, UTAH, WASHINGTON AND WYOMING**

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

When applied as directed in this label, there are no grazing restrictions.

**APPLICATION EQUIPMENT AND TECHNIQUES**

Applications may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre.

**BROMUS SPECIES**

This product may be used to treat downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*) and cheatgrass (*Bromus secalinus*) found in industrial rangeland and pasture sites. Apply 8 to 16 fluid ounces of product per acre on a broadcast basis.

For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses are able to become reestablished on the site.

---

**MEDUSAHEAD**

---

To treat medusahead, apply 16 fluid ounces of this product per acre as soon as plants are actively growing, and prior to the 4 leaf stage. Applications may be made in the fall or spring.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus, and this supplement.

**Albaugh**  
**GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx



---

**FOR USE IN NONBEARING GINSENG**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product may be used for general weed control in established non-bearing ginseng.

Direct applications so that there is no contact of this product with the ginseng plant.

Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high volume wands, lances, orchard guns or with wiper application equipment. Applications must be made at least one year prior to harvest.

For specific rates of application and instructions for control of various annual and perennial weeds, see the Weed Control tables in the label booklet for Glyphosate 41% Plus.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION. SPRAY, DRIFT OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT OR OTHER PARTS OF DESIRABLE PLANTS. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**




**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus, and this supplement.

**Albaugh**  
**GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx



---

**HOODED SPRAYERS ON CORN (ALL)**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product may be used through hooded sprayers for weed control between the rows of corn (all), including field corn, sweet corn and popcorn. Only hooded sprayers that completely enclose the spray pattern may be used.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way.

If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 1 quart of this product per acre per application.
- Corn must be at least 12 inches tall, measured without extending leaves.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 mph.
- Maximum wind speed: 10 mph.
- Use low-drift nozzles.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

For specific rates of application and instructions for control of various annual and perennial weeds, see the Weed Control tables in the label booklet for Glyphosate 41% Plus.

Do not graze or feed corn forage or fodder following applications of this product through hooded sprayers.

Do not apply more than 3 quarts of this product per acre per year for hooded sprayer applications.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh**  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx



---

**FOR WEED CONTROL IN CITRUS UP TO ONE DAY PRIOR TO HARVEST**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product is recommended for weed control in established groves of citrus. Applications may be made with boom equipment, shielded sprayers, hand-held and high volume wand, lances and orchard guns, CDA equipment or with wiper application equipment.

For specific rates of application and instructions, see the Weed Control Rate Tables and the Citrus Crops sections of the label booklet for Glyphosate 41% Plus.

Allow a minimum of one day between last application and harvest.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh  
GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx



---

**FOR RED RICE CONTROL IN TEXAS ONLY**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product may be used to control red rice, prior to planting rice. Apply 1-½ quarts of this product in 5 to 10 gallons of water per acre.

Flush fields prior to applications to obtain uniform germination and stand of red rice. Make applications when the majority of the red rice plants are in the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may only be partially controlled. Do not treat rice fields or levees when the fields contain floodwater.

Avoid spraying during low humidity conditions, as reduced control may result.

Do not re-flood treated fields for 8 days following application.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

Albaugh  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx



---

**FOR USE IN DORMANT RANGELAND**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product will control or suppress many weeds, including downy brome, cheat grass, cereal rye, medusahead rye and jointed goatgrass in dormant rangeland.

Apply 8 to 16 fluid ounces of this product per acre in the early spring when the weeds have greened up, but desirable grasses, such as crested and tall wheatgrass, are still truly dormant.

Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off.

Do not use ammonium sulfate when spraying dormant rangeland grasses with this product.

Do not make more than one application per year.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021



**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh  
GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx


**FOR DISTRIBUTION AND USE ONLY WITHIN ARKANSAS, LOUISIANA, MISSISSIPPI,  
MISSOURI AND TENNESSEE**

**FOR CONTROL OF EMERGED WEEDS PRIOR TO THE PLANTING OR EMERGENCE OF  
CORN, COTTON, RICE, SORGHUM AND SOYBEANS; PRIOR TO THE HARVEST OF  
COTTON AND SOYEANS; AND FOLLOWING THE HARVEST OF ANY CROP\* IN THE FALL  
VIA AERIAL APPLICATIONS.**

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMICAL RESPONSE



\*Do not plant subsequent crops other than those on the label for 30 days following application.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

**RECOMMENDATIONS**

This product controls annual and perennial weeds listed in the label booklet for Glyphosate 41% Plus prior to planting or emergence of corn, cotton, rice, sorghum and soybeans; prior to the harvest of cotton and soybeans; and following the harvest of any crop in the fall via aerial applications.

Aerial applications of this product may be made in fallow systems and conventional, reduced and zero tillage systems.

For applications via aerial equipment, use the recommended rates of this product in 3 to 10 gallons of water per acre. Do not exceed a rate of 3 quarts per acre.

AVOID DRIFT – DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser velocities, will allow spray drift to occur.

Read the "AERIAL EQUIPMENT" section of the Glyphosate 41% Plus label booklet for additional precautions and instructions.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh**  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx

---


**GLYPHOSATE 41% PLUS FOR USE IN POPLAR (*Populus spp.*) PRODUCTION**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMICAL



42750-61

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the Glyphosate 41% Plus label booklet for essential product performance information.

---

**PREPLANT**

---

This product is recommended for use prior to planting *Populus spp.* This includes, but is not limited to, hybrid poplars and hybrid cottonwoods.

See the Glyphosate 41% Plus label booklet for specific rates for the weeds being controlled.

---

**IN CROP USES**

---

**DIRECTED SPRAYS**

Use a 2 percent spray solution as a spray-to-wet application for the control of undesirable woody brush and trees. To control herbaceous weeds, use a 1 to 2 percent solution. Avoid contact of spray, drift, or mist with foliage, green bark or non-woody surface roots of the *Populus spp.*

**WIPERS**

This product may be used through wick or other suitable wiper applicators for control or partial control of grass and broadleaf weeds listed on the label.

For wick applicators, mix 1 gallon of this product with 2 gallons water to make a 33% solution. For wiper systems that can handle thicker solutions, such as force fed systems, a 33 to 100% Glyphosate 41% Plus solution may be used.

For best results ensure that the herbicide solution is allowed to contact the maximum amount of leaf surface. As weed densities increase, decrease equipment speed to allow sufficient herbicide flow to wet all weed surfaces contacted. Weeds not contacted will be unaffected.

**AVOID HERBICIDE CONTACT WITH DESIRABLE VEGETATION.** Desirable vegetation contacted by the herbicide solution may be injured or controlled. This includes foliage, fruit, or green stems.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh  
GLYPHOSATE 41% PLUS**


EPA Reg. No. 42750-xxx

**SHIELDED APPLICATOR USE IN GRASSES GROWN FOR SEED IN IDAHO, OREGON AND  
WASHINGTON ONLY**

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMICAL



**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

When applied using shielded applicator equipment designed to prevent direct contact, movement of spray droplets, or mist onto desirable grasses grown for seed production, this product may be used to control labeled weeds. Use of low spray pressure through low pressure nozzles will minimize the potential of spray drift.

Apply 1 to 3 quarts of this product as a broadcast spray in 10 to 20 gallons of total spray volume per acre.

Uniform planting in straight rows aid in shielding sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by or through the protective shields.

**GROWER ASSUMES ALL RESPONSIBILITY FOR CROP LOSSES FROM  
MISAPPLICATION.**

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021



**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh**  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx



---

**FOR WEED CONTROL IN BANANAS FOR UP TO ONE DAY PRIOR TO HARVEST**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product may be used for weed control in established plantations of bananas. Applications may be made with boom equipment, shielded sprayers, hand-held and high-volume wands, lances and orchard guns, CDA equipment or wiper application equipment.

For specific rates of application and instructions, see the Weed Control Tables and the Tree and Vine Crops sections of the label booklet for Glyphosate 41% Plus.

Allow a minimum of 1 day between last application and harvest.

FOR DIRECT APPLICATION TO BANANAS (BANANACIDE), REMOVE FRUIT PRIOR TO TREATMENT.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh**  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx



---

**FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA  
FOR NONSELECTIVE CONTROL OF LISTED ANNUAL WEEDS IN SMALL GRAIN  
CROPPING SYSTEMS**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

Refer to the Glyphosate 41% Plus label for rate recommendations and weeds controlled.

For ground applications, use 3 to 5 gallons of water per acre. For aerial applications, use 2 to 3 gallons of water per acre.

**ATTENTION**

**AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.**

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this herbicide can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the

use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

#### NOTE

- To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.
- Adjust boom height on ground equipment to prevent streaked, overlapped or uneven applications. Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets.
- In aerial applications, do not angle nozzles forward into the airstream and to not increase spray volume by increasing nozzle pressure.
- Ensure uniform application. Use appropriate marking devices when applying herbicides by air.
- Avoid spraying when weeds are subject to moisture stress, when dust is on foliage, or when straw canopy covers the weeds.
- Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residue of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh**  
**GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx



---

**FOR AERIAL APPLICATION IN ARKANSAS ONLY**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.  
This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING" sections of the Glyphosate 41% label booklet for essential product performance information.

**USE DIRECTIONS**

AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the recommended rate of this product in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are recommended.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75% of the length of the wingspan rotor. In many cases reducing this distance to 65% of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing airflow on rotary winged aircraft. Avoid the use of nozzles with wide-angle discharge.

Do not apply this product when wind speeds are in excess of 10 miles per hour.

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021



**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh  
GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx



---

**PREHARVEST APPLICATIONS IN GRAIN SORGHUM (MILO)**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product may be used prior to harvest of grain sorghum (milo) to control annual and perennial weeds listed on the label booklet for Glyphosate 41% Plus.

Apply up to 2 quarts of this product per acre.

Make applications at 30% grain moisture or less and at least 7 days prior to harvest.

This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for instructions.

Note: It is not recommended that grain sorghum grown for seed be treated because a reduction in germinate or vigor may occur.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

## SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**WARNING:** USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

### Albaugh GLYPHOSATE 41% PLUS

EPA Reg. No. 42750-xxx


#### FOR POSTEMERGENCE APPLICATIONS TO CORN WITH THE ROUNDUP READY® GENE

Keep out of reach of children

#### CAUTION!

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMICAL

Roundup Ready® is a registered trademark of Monsanto Company.



42750-61

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

#### GENERAL INFORMATION

ALBAUGH, INC. RECOMMENDS USE OF THIS PRODUCT ONLY ON CORN SEED DESIGNATED AS CONTAINING THE ROUNDUP READY® GENE.

- Applying this product to corn varieties which are not designated as Roundup Ready® will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Roundup Ready gene since severe injury or destruction will result.
- Roundup Ready varieties must be purchased from an authorized seed supplier. Crop safety and weed control performance are not warranted by Albaugh, Inc. when this product is used

in conjunction with "brown bag" or "bin run" corn seed saved from previous year's production and replanted.

- The Roundup Ready designation indicates that the corn contains a patented gene which provides tolerance to Albaugh's Glyphosate brand herbicides. Information on Roundup Ready corn may be obtained from your seed supplier.

---

### APPLICATION INSTRUCTIONS

---

This product may be applied postemergence to Roundup Ready corn during the period beginning at corn emergence and continuing through the 12-leaf stage or until corn height reaches 30 inches, whichever comes first. Single in-crop applications of this product are not to exceed 1 quart per acre. Sequential in-crop applications of this product should not exceed 2 quarts per acre per growing season. Total Glyphosate 41% Plus use should not exceed 8 quarts per acre per year.

Allow a minimum of 50 days between application of this product and harvest of corn forage and 7 days between application and harvest of corn grain. Allow a minimum of 14 days between in-crop applications of this product. **THE USE OF ADDITIVES FOR IN-CROP APPLICATIONS TO ROUNDUP READY CORN IS PROHIBITED.**

#### Maximum Yearly Rates Allowed

Preplant/Preemergence (Maximum)	5 quarts/A
Total in-crop applications from emergence to 12-leaf stage or 30 inches	2 quarts/A
Maximum preharvest rate	1 quart/A
Combined total per year for all applications	8 quarts/A

When applied as directed, this product controls annual grass and broadleaf weeds in Roundup Ready corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product. Applications should be made to actively growing weeds before they reach the maximum size listed in the "WEEDS CONTROLLED" section of the label booklet. Refer to the "MIXING" section of the label booklet for proper use instructions.

There are no rotational crop restrictions following applications of this product.

**ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.**

Do not allow the herbicide solution to mist, drip, drift, or splash onto other desirable vegetation since minute quantities of this product can cause severe damage or destruction to crop plants in non-target areas. The likelihood of plant injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions which allow spray drift to occur such as combinations of pressure and nozzle type that will result in fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR SPRAY PRESSURE.**

For ground applications: Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. See "WEEDS CONTROLLED" section of this label for specific recommended rates. Carefully select proper nozzles and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

**For aerial applications:** Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart per acre. See "WEEDS CONTROLLED" section on this label. AVOID DRIFT – DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

### **Weeds controlled**

For specific rates of application and instructions for control of various annual and perennial weeds, refer to the label booklet. Glyphosate 41% Plus at up to 1 quart per acre will burndown or suppress the growth of the following perennial weeds and reduce crop competition: nutsedge, rhizome johnsongrass, quackgrass, Canada thistle, wirestem muhly.

**Sequential Applications:** Some weeds with multiple germination times or suppressed (stunted) weeds may require a second application of this product. The second application should be made after some regrowth has occurred.

### **Tank mixtures**

A tank mixture of Glyphosate 41% Plus plus Micro-Tech or Partner may be used for postemergence and residual control of annual weeds in corn. These tank mixtures may be made during the period beginning at corn emergence and continuing until corn height reaches 5 inches.

A Glyphosate 41% Plus tank mixture with atrazine, Banvel, Clarity, Permit or 2,4-D may be used for postemergence control of additional annual weeds in corn. A Glyphosate 41% Plus tank mixture with atrazine may be made during the period beginning at corn emergence and continuing until corn height reaches 12 inches. A Glyphosate 41% Plus tank mixture with Banvel or Clarity at 0.125 to 0.25 lb per acre may be made during the period beginning at corn emergence and continuing until corn height reaches 30 inches. A Glyphosate 41% Plus tank mixture with Permit may be made during the period beginning at corn emergence and continuing until corn height reaches 30 inches. A Glyphosate 41% Plus tank mixture with 2,4-D at 0.125 to 0.25 lb per acre may be made during the period beginning at corn emergence and continuing until corn is at the five leaf stage or corn height reaches 8 inches, whichever comes first.

Refer to the specific product label and observe all precautions, mixing and application instructions for all products used in tank mixtures.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh**  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx

---

**PREHARVEST APPLICATIONS IN CORN**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMICAL SERVICE (800) 424-9300

**ACCEPTED**

2/9/01

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 42750-61

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product may be used prior to harvest of corn to control annual and perennial weeds listed on the label booklet for Glyphosate 41% Plus.

For ground applications, apply up to 3 quarts of this product per acre. For aerial applications, apply up to 1 quart of this product per year.

Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed). Allow a minimum of 7 days between application and harvest.

This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for instructions.

Note: It is not recommended that corn grown for seed be treated because a reduction in germination or vigor may occur.



Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh  
GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx


---

**FOR TRANSPLANTED TOMATOES**

---

Keep out of reach of children  
**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMICAL



---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product may be applied prior to transplanting tomatoes. Allow at least 3 days between application and planting.

For specific rates of application and instructions for control of various annual and perennial weeds, see the Weed Control tables in the label booklet for Glyphosate 41% Plus.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh**  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx

---


**GLYPHOSATE 41% PLUS FOR USE IN EUCALYPTUS (*Eucalyptus spp.*) PRODUCTION**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMICAL



42750-61

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See the label booklet for Glyphosate 41% Plus for essential product performance information.

This product is recommended for use prior to planting eucalyptus species.

**Ground Broadcast Sprays:**

Use the recommended rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified.

See the Glyphosate 41% Plus label booklet for specific rates for the weeds being controlled.

---

**IN ESTABLISHED EUCALYPTUS**

---

This product can be used around established eucalyptus trees to control undesirable vegetation.

**Directed Sprays (Spray-to-wet applications)**

Use a 1 to 2 percent spray solution to control herbaceous weeds in eucalyptus farms. Use a 2 percent spray solution for control of undesirable woody brush and trees. For "hard-to-control" weeds, use a 5 to 10 percent spray solution. Avoid contact of spray, drift, or mist with foliage, green bark or non-woody surface roots of eucalyptus plants.

**Wipers**

This product may be used through wick or other suitable wiper applicators for control or partial control of grass and broadleaf weeds listed on the product label.

For wick applicators, mix 1 gallon of this product with 2 gallons water to make a 33% solution. For wiper systems that can handle thicker solutions, such as force fed systems, a 33 to 100% Glyphosate 41% Plus solution may be used.

For best results ensure that the herbicide solution is allowed to contact the maximum amount of leaf surface. As weed densities increase, decrease equipment speed to allow sufficient herbicide flow to wet all weed surfaces contacted. Weeds not contacted will be unaffected.

**AVOID HERBICIDE CONTACT WITH DESIRABLE VEGETATION.** Desirable vegetation contacted by the herbicide solution may be injured or controlled. This includes foliage, fruit, or green stems.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**WARNING:** USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

**Albaugh  
GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx


**USE ONLY FOR SEED PRODUCTION OF SUGARBEETS WITH THE ROUNDUP READY®  
GENE**

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTEC (800) 424-9300

®Round Ready is a registered trademark of the Monsanto Company.



**GENERAL INFORMATION**

ROUNDUP READY® SUGARBEET VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE SUGARBEET VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT. IT IS UNLAWFUL TO SELL OR PLANT SAVED SEED.

SUGARBEETS WITH THE ROUNDUP READY® GENE MAY ONLY BE USED FOR PLANTING A COMMERCIAL CROP IN A SINGLE SEASON. SEED MAY NOT BE SAVED FOR REPLANTING AND SAVED SEED MAY NOT BE SUPPLIED TO OTHERS FOR REPLANTING. ALBAUGH, INC. DOES NOT WARRANT THE SAFETY OR PERFORMANCE OF THIS PRODUCT WHEN USED ON "BROWN BAG" OR FARMER-SAVED SEED.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.



AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT SUGARBEETS IN PRODUCTION FIELDS OF SUGARBEETS CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH WILL RESULT IF SUGARBEETS VARIETIES WHICH DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

---

#### USE RECOMMENDATIONS

---

This product will control non-glyphosate tolerant sugarbeets in seed production fields of sugarbeets containing the Roundup Ready gene. This product may be applied using ground spray equipment only. Apply 24 ounces of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application of 24 ounces per acre may be applied if needed to control non-glyphosate tolerant sugarbeet plants.

DO NOT EXCEED A MAXIMUM RATE OF 1 ½ QUARTS OF THIS PRODUCT PER ACRE PER SEASON.

Application timing – This product can be applied to Roundup Ready sugarbeets from emergence to canopy closure.

Treated sugarbeets may not be used for food or feed. Do not feed or graze treated sugarbeets. Do not process treated sugarbeets for food or feed.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh**  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx



**FOR PISTACHIOS UP TO THREE DAYS PRIOR TO HARVEST**

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

**RECOMMENDATIONS**

This product is recommended for weed control in established groves of pistachios. Applications may be made with boom equipment, shielded sprayers, hand-held and high volume wands, lances and orchard guns, CDA equipment or with wiper application equipment.

For specific rates of application and instructions, see the Weed Control Tables and Tree Nuts sections of the label booklet for Glyphosate 41% Plus.

Allow a minimum of 3 days between last application and harvest.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh**  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx

---

**GLYPHOSATE 41% PLUS FOR USE IN DORMANT ALFALFA**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

**ACCEPTED**

2/9/01

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 42750-61

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product will control or suppress many weeds including quackgrass, downy brome and cheatgrass in dormant alfalfa.

Apply 8 to 12 ounces per acre of this product. Apply in the spring to alfalfa that is dormant. Applications should be made after spring temperatures have warmed enough to encourage resumption of weed growth, but prior to initiation of trifoliate leaf expansion of the alfalfa. Applications made after expansion of the first trifoliate leaf of the alfalfa will cause growth reduction and reduced crop yield.

Do not use ammonium sulfate when spraying dormant alfalfa with Glyphosate 41% Plus.

Do not use this product where a slight yield reduction in the first cutting of alfalfa cannot be tolerated.

Do not make more than one application per year.

Allow 36 hours after application before grazing livestock or harvesting.

Slight discoloration of the alfalfa may occur, but the alfalfa will regreen and regrow under moist soil conditions as effects of this product wear off.

Application of this product can cause crop injury. Any crop injury is the sole responsibility of the applicator.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**WARNING:** USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

**Albaugh**  
**GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx

**USE ONLY FOR SEED PRODUCTION OF CANOLA WITH THE ROUNDUP READY® GENE**

Keep out of reach of children

**CAUTION!**

**ACCEPTED**

2/9/01

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 42750-61

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, OTHER THAN CANOLA BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

**NOTE:** THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT CANOLA IN PRODUCTION FIELDS OF CANOLA CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH WILL RESULT IF CANOLA VARIETIES WHICH DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

ROUNDUP READY® CANOLA VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE CANOLA VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT: IT IS UNLAWFUL TO SELL OR PLANT SAVED SEED.



CANOLA WITH THE ROUNDUP READY® GENE MAY ONLY BE USED FOR PLANTING A COMMERCIAL CROP IN A SINGLE SEASON. SEED MAY NOT BE SAVED FOR REPLANTING AND SAVED SEED MAY NOT BE SUPPLIED TO OTHERS FOR REPLANTING. ALBAUGH, INC. DOES NOT WARRANT THE SAFETY OR PERFORMANCE OF THIS PRODUCT WHEN USED ON "BROWN BAG" OR FARMER-MADE SEED.

---

#### USE RECOMMENDATIONS

---

This product will control non-glyphosate tolerant canola in seed production fields of canola containing the Roundup Ready gene. This product may be applied using ground spray equipment only. Apply 1 pint of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application of 1 pint per acre may be applied, if needed to control non-glyphosate tolerant canola plants.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART OF THIS PRODUCT PER ACRE PER SEASON.

Application timing – This product can be applied to Roundup Ready canola from emergence to the pre-flower (early bolting) stage.

Treated canola may not be used for food or feed. Do not feed or graze treated canola. Do not process treated canola for food or feed.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

**Albaugh  
GLYPHOSATE 41% PLUS**

EPA Reg. No. 42750-xxx

---


**FOR USE IN PASTURES FOR HAWAII ONLY**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEM-TREK (800) 424-9300



---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**RECOMMENDATIONS**

---

This product may be used in Hawaiian pastures, including kikuyu grass, pangola grass and guineagrass pastures. Spot treatment, wiper application, preplant, preemergence and pasture renovation applications may be made.

**Spot treatment**

This product may be applied as a spot treatment or with wiper applicators in pastures. Applications may be made in the same area at 30-day intervals. For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated at one time. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

**Preplant, preemergence and pasture renovation**

When applied prior to planting or emergence of forage grasses or as a direct application to the listed perennial pasture species, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

See the Weed Control Tables in the label booklet for Glyphosate 41% Plus for information about rates and timings for specific weeds

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**

**SUPPLEMENTAL LABELING**


READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus and this supplement.

WARNING: USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

**Albaugh**  
**GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx

Roundup Ready® is a registered trademark of the Monsanto Company



---

**FOR COTTON WITH THE ROUNDUP READY® GENE**  
**IN-CROP APPLICATIONS**

---

Keep out of reach of children

**CAUTION!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of application.

---

**GENERAL INFORMATION**

---

WARNING: ALBAUGH, INC. RECOMMENDS THIS PRODUCT FOR USE ONLY OVER-THE-TOP OF OR DIRECTED ONTO IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE ROUNDUP READY® GENE. SEVERE INJURY OR DEATH OF COTTON WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY® GENE ARE SPRAYED WITH THIS PRODUCT. AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS AND TREES, OTHER THAN CROPS WITH THE ROUNDUP READY® GENE, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

ROUNDUP READY® COTTON VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE COTTON VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT. IT IS UNLAWFUL TO SELL OR PLANT SAVED SEED.

COTTON WITH THE ROUNDUP READY® GENE MAY ONLY BE USED FOR PLANTING A COMMERCIAL CROP IN A SINGLE SEASON. SEED MAY NOT BE SAVED FOR REPLANTING AND SAVED SEED MAY NOT BE SUPPLIED TO OTHERS FOR REPLANTING. ALBAUGH, INC. DOES NOT WARRANT THE SAFETY OR PERFORMANCE OF THIS PRODUCT WHEN USED ON "BROWN BAG" OR FARMER-MADE SEED.

---

### APPLICATION INSTRUCTIONS

---

This product will control many troublesome weeds with over-the-top directed, post-directed, hooded sprayer, or preharvest applications in Roundup Ready cotton.

#### Maximum Allowable Yearly Rates

1. Combined total per year for all applications	8 quarts/A
2. Preplant, Preemergence applications	5 quarts/A
3. Total in-crop applications from cracking to layby	4 quarts/A
4. Maximum preharvest application rate	2 quarts/A

For ground applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

For aerial applications apply this product in 3 to 15 gallons of water per acre.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY® GENE. Do not apply during low-level inversion conditions, when winds are gusty or under any other conditions which favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

**There are no rotational crop restrictions following applications of this product.**

**Sprayer Preparation:** It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to Roundup Ready cotton. Follow the cleaning procedures specified on the label of the product(s) previously used. Cotton is very sensitive to many herbicides at extremely low concentrations and care should be taken to thoroughly clean all equipment prior to use.

In addition to uses listed in the Glyphosate 41% Plus label booklet, the following applications can be made:

**Over-the-top applications:** This product may be applied by aerial or ground application equipment postemergence to Roundup Ready cotton from the ground cracking stage until the four leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter).

Over-the-top applications made after the four leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss. Any single over-the-top broadcast application should not exceed 1 quart per acre. No more than two over-the-top broadcast applications may be made from crop emergence through the four leaf (node) stage of development. Sequential over-the-top applications of this product must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

NOTE: Always plant into a weed free seedbed. In no-till and stale seedbed systems, always burn down existing weeds before cotton emerges. Apply a preplant burndown treatment of 16-48 fluid ounces per acre of this product.

**Post-directed or hooded applications:** This product may be applied using precision post-directed or hooded sprayers to Roundup Ready cotton through layby. Be especially careful to minimize contact of the spray with cotton leaves. At this stage, post-directed equipment should be used which directs the spray to the base of the cotton plants. Place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row. For best results, make applications while weeds are small (less than 3 inches). Minimize spray drift onto the leaves of the cotton plants by maintaining low spray pressure (less than 30 PSI). Applications that contact the cotton leaves may result in boll loss, delayed maturity and/or yield loss. Any single post-directed application should not exceed 1 quart per acre of this product. No more than two applications should be made from the fifth leaf through layby. Sequential in-crop applications of this product must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

**Salvage Treatment:** This treatment may be used after the four leaf stage of development and should only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top application or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER GROWING SEASON.

**Weeds controlled:** For specific rates of application and instructions for control of various annual and perennial weeds, refer to "ANNUAL WEEDS RATE TABLE" section of the label booklet. Glyphosate 41% Plus applied at 1 quart per acre will burndown or suppress the growth of the following perennial weeds and reduce crop competition: yellow and purple nutsedge, rhizome johnsongrass, common bermudagrass, silverleaf nightshade, trumpet creeper, and Redvine. Fall preharvest applications may be required for control of these perennial weeds.

Tank mixtures with other herbicides may result in reduced weed control or crop injury and are not recommended for over-the-top applications of this product.

Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of this product for control.

**Preharvest applications:** This product may be applied for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready cotton after 20% boll crack. Allow a minimum of 7 days between application and harvest. For specific recommendations refer to the "COTTON" section of the Glyphosate 41% Plus label booklet.

NOTE: Glyphosate 41% Plus will not enhance the performance of harvest aids when applied to Roundup Ready cotton. DO NOT APPLY GLYPHOSATE 41% PLUS TO CROPS GROWN FOR SEED.



Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021

**NEXT**

**LABEL**


**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR GLYPHOSATE 41% PLUS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Label as used in this supplemental labeling refers to the label booklet for Glyphosate 41% Plus, container labels and this supplement.

WARNING: USE OF THIS PRODUCT OVER "ROUNDUP READY" OR OTHER GLYPHOSATE-TOLERANT CROPS MAY SUBJECT YOU TO THE RISK OF LOSS OF LICENSE RIGHTS TO PATENTED GLYPHOSATE TOLERANCE TECHNOLOGIES AND/OR LEGAL ACTION FOR INFRINGEMENT OF PATENTS TO THOSE GLYPHOSATE-TOLERANT TECHNOLOGIES. IF YOU ARE A LICENSED GROWER UNDER AN AGREEMENT WITH A GLYPHOSATE-TOLERANT SEED MANUFACTURER, PLEASE REFER TO YOUR LICENSE AGREEMENT TO DETERMINE WHETHER YOU MAY USE THIS PRODUCT WITHOUT RISK OF LOSING YOUR LICENSE OR OF LEGAL ACTION AGAINST YOU.

**Albaugh**  
**GLYPHOSATE 41% Plus**

EPA Reg. No. 42750-xxx



---

**FOR POSTEMERGENCE APPLICATIONS TO SOYBEANS WITH THE ROUNDUP READY®  
GENE**

---

Keep out of reach of children

**Caution!**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

Roundup Ready® Gene and Roundup Ready® are registered trademarks of the Monsanto Company.

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "GENERAL INFORMATION" and "APPLICATION INSTRUCTIONS" sections of the label booklet for Glyphosate 41% Plus for essential product performance information.

---

**GENERAL INFORMATION**

---

ALBAUGH, INC. RECOMMENDS USE OF THIS PRODUCT FOR POSTEMERGENCE APPLICATION ONLY ON SOYBEAN VARIETIES DESIGNATED AS CONTAINING THE ROUNDUP READY® GENE.

- Applying this product to soybean varieties which are not designated as Roundup Ready® will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Roundup Ready gene, since severe injury or destruction will result.
- Roundup Ready varieties must be purchased from an authorized seed supplier. Crop safety and weed control performance are not warranted by Albaugh, Inc. when this product is used in conjunction with "brown bag" or "bin run" soybean seed saved from previous year's production and replanted.
- The Roundup Ready designation indicates that the soybean contains a patented gene which provides tolerance to Albaugh's Glyphosate brand herbicides. Information on Roundup Ready soybeans may be obtained from your seed supplier.

---

### APPLICATION INSTRUCTIONS

---

This product may be applied postemergence to Roundup Ready soybeans from the cracking stage throughout flowering.

Allow a minimum of 14 days between application and harvest of soybeans.

---

### MAXIMUM ALLOWABLE YEARLY RATES

---

**Preplant:** Maximum amount of this product which can be applied prior to crop emergence is 5 quarts/A.

**In-crop:** Maximum combined total of multiple in-crop applications from cracking throughout flowering is 3 quarts/A. The maximum rate for any single in-crop application is 2 quarts/A. The maximum combined total of this product which can be applied during flowering is 2 quarts/A.

**Preharvest:** Maximum amount of this product that can be applied after loss of green color in soybean pods until 14 days before harvest is 1 quart/A. The combined total of in-crop and preharvest Glyphosate 41% Plus applications may not exceed 3 quarts/A.

**Cropping Season:** Combined total per year for all applications may not exceed 8 quarts/A.

When applied as directed, this product will control labeled annual grasses and broadleaf weeds in Roundup Ready soybeans. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product.

There are no rotational crop restrictions following applications of this product.

**For ground applications:** Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use nozzles which provide a flat fan pattern. Check for even distribution of spray droplets.

**For aerial applications:** Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart of this product per acre. DO NOT APPLY DURING LOW LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. MAINTAIN APPROPRIATE BUFFER ZONES TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION.

AERIAL APPLICATIONS ON ROUNDUP READY SOYBEANS MAY BE MADE ONLY IN THE FOLLOWING STATES: ALABAMA, ARKANSAS, FLORIDA, GEORGIA, KANSAS, LOUISIANA, MISSISSIPPI, MISSOURI (BOOT-HEEL ONLY), NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE AND TEXAS.

---

### ANNUAL WEED RATE TABLES

---

The following rate recommendations will provide control of labeled grasses and broadleaf weeds in conventional and no-till soybean production systems. Refer to the label booklet for rate recommendations for specific annual weeds.

Albaugh, Inc. will not warrant crop safety or weed control when Roundup Ready soybeans are treated with herbicides not specified on this supplemental label. Because of the potential for: 1) crop injury, 2) poor weed control from antagonism, and/or 3) rotational crop restrictions, herbicides not specified on this supplemental label should not be used, whether applied preemergence or applied postemergence as a tank mixture with Glyphosate 41% Plus.

This product may be used up to 64 fluid ounces per acre in any single application for control of annual weeds, where heavy weed densities exist. The maximum combined total of this product which can be applied during flowering is 64 fluid ounces per acre.

NOTE: The following recommendations are based on a clean start at planting by using a burn down application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burn-down treatment of 16-64 fluid ounces per acre of this product can be used to control existing weeds prior to crop emergence.

---

### MIDWEST/MID-ATLANTIC RECOMMENDATIONS

---

**Narrow row or drilled soybeans:** A single in-crop application of this product will provide effective control of labeled weeds. For best results, an initial application of 32 fluid ounces per acre (fl oz/A) on 4-8" weeds is recommended. Weeds will generally be 4-8" tall 3 to 5 weeks after planting. If the initial application is delayed and weeds are 8-18" tall, use 48 fl oz/A for best results.

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this product at 16 to 32 fluid ounces per acre may be necessary to control late flushes of weeds. The combined total application in-crop must not exceed 64 fluid ounces per acre.

**Wide row soybeans:** An in-crop application of this product will provide effective control of the initial stand of labeled weeds. For best results, an initial application of 32 fluid ounces per acre (fl oz/A) on 4-8" weeds is recommended. Weeds will generally be 4-8" tall 3 to 5 weeks after planting. If new flushes of weeds occur, they can be controlled by sequential applications of this product.

**Initial Treatment**

Weed Height (inches)	Rate (fluid oz/A)
8	32
18	48

**Sequential Application (if needed)\***

Weed Height (inches)	Rate (fluid oz/A)
1-3	16
3-6	24
6-12	32

\*Combined total application in-crop not to exceed 96 fluid ounces per acre.

Giant ragweed: Apply 32 fluid oz/A when the weed is 8-12" tall to avoid the need for sequential application.

Groundcherry, ladysthumb, Pennsylvania smartweed and morningglory: Apply 32 fl oz/A to weeds 3-6" tall.

Some weeds, such as black nightshade, woolly cupgrass, shattercane, wild proso millet, burcucumber, and giant ragweed, with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 96 fluid ounces per acre.

---

**SOUTHEAST RECOMMENDATIONS**


---

**Narrow row, drilled, or wide-row soybeans:** An in-crop application of this product will provide effective control of the initial stand of labeled weeds. For best results, an initial application of 32 fluid ounces per acre (fl oz/A) on 3-6" weeds is recommended. Weeds will generally be 3-6" tall 2 to 3 weeks after planting.

Weed Height (inches)	Rate (fluid oz/A)
3-66	32
6-12	48

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this product at 16 to 32 fluid ounces per acre may be necessary to control late flushes of weeds.

**Sequential Application (if needed)\***

Weed Height (inches)	Rate (fluid oz/A)
2-3	16
3-6	24
6-12	32

\*Combined total application in-crop not to exceed 96 fluid ounces per acre.

Florida pusley, hemp sesbania and spurred anoda: Apply 32 fl oz/A to weeds 2-4" for the initial application. Apply 32 oz/A when these weeds are 3-6" tall if a sequential application is necessary.

Morningglory, black nightshade, groundcherry, and Pennsylvania smartweed: Apply 24 fl oz/A on 1-3" weeds, 32 fl oz/A on 3-6" weeds, or 48 fl oz/A on 6-12" weeds for the initial application.

Some weeds, such as black nightshade, broadleaf signalgrass, Texas panicum, burcumber, and sicklepod, with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 96 fluid ounces per acre.

---

**DELTA/MID-SOUTH RECOMMENDATIONS**


---

**Narrow row, drilled, or wide row soybeans:** An in-crop application of this product will provide effective control of the initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds. For best results, an initial application of 32 fluid ounces per acre (fl oz/A) on 2-4" weeds is recommended. Weeds will generally be 2-4" tall 2 to 3 weeks after planting.

**Initial Treatment**

Weed Height (inches)	Rate (fluid oz/A)
2-4	32
5-12	48

**Sequential Application\***

Weed Height (inches)	Rate (fluid oz/A)
2-3	16
3-6	24
6-12	32

\*Combined total application in-crop not to exceed 96 fluid ounces per acre.

Hemp sesbania and spurred anoda: Apply a sequential treatment of 32 fl oz/A at 3-6" tall weeds if necessary.

Some weeds, such as black nightshade, broadleaf signalgrass, Texas panicum, burcumber, and sicklepod, with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 96 fluid ounces per acre.

### **PERENNIAL WEEDS RATE RECOMMENDATIONS**

A 32 to 64 fluid ounces per acre rate (single or sequential applications) of this product will control or suppress perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, maretail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly.

For best results, allow perennial weed species to achieve at least 6" of growth before spraying with Glyphosate 41% Plus. For additional information on perennial weeds, see the "PERENNIAL WEED RATE TABLE" of the label booklet for Glyphosate 41% Plus. For some perennial species, repeat applications may be required to eliminate crop competition throughout the growing season.

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for Glyphosate 41% Plus before using this product. These terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Albaugh, Inc.  
121 NE 18<sup>th</sup> Street  
Ankeny, IA 50021