



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 22, 2020

Carolyn Miter
Registrations Specialist
Albaugh, LLC
P.O. Box 2127
Valdosta, GA 31604

Subject: Label Amendment – Update primary brand name, update Directions for Use, and manufacturer name on label
Product Name: DDG 3-Way Herbicide
EPA Registration Number: 42750-76
Application Date: May 20, 2019
Decision Number: 557450

Dear Ms. Miter:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 42750-76
Decision No. 557450

with FIFRA section 6. If you have any questions, please contact Theresa Gerber at 703-347-8583 or by email at gerber.theresa@epa.gov.

Sincerely,



Emily Schmid, Product Manager 25
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

GLYPHOSATE	GROUP	9	HERBICIDE
2,4-D	GROUP	4	HERBICIDE
DICAMBA	GROUP	4	HERBICIDE

DDG 3-WAY HERBICIDE

DDG 3-Way Herbicide is a postemergence herbicide for control or suppression of emerged weeds in fallow and reduced tillage systems, industrial and noncrop areas. DDG 3-Way Herbicide can also be applied prior to planting or emergence of small grains (wheat, barley, oats) corn, sorghum (grain or forage), through hooded sprayers between rows of corn, as a spot treatment in corn, sorghum, forage grasses, as a preharvest treatment in wheat, and as a postharvest treatment following grain harvest.

AVOID CONTACT WITH FOLIAGE OF CROP OR OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

Read the "CONDITIONS OF SALE AND WARRANTY" before buying or using. If terms are not acceptable, return at once unopened.

ACTIVE INGREDIENTS*:

Glyphosate (N-(phosphonomethyl))glycine, in the form of its isopropylamine salt.....	23.56%
2,4-D, 2,4-dichlorophenoxyacetic acid, in the form of its isopropylamine salt	14.00%
Dicamba (3,6 dichloro-o-anisic acid), in the form of its isopropylamine salt.....	4.90%
OTHER INGREDIENTS:	57.54%
TOTAL	100.00%

*Contains: 2.28 lbs per gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt;
1.36 lbs per gallon of the active ingredient, 2,4-D, in the form of its isopropylamine salt; 0.5 lb per gallon of the active ingredient, dicamba, in the form of its isopropylamine salt.
Equivalent to 1.69 lbs per gallon of glyphosate acid; 1.07 lbs per gallon of 2,4-D acid; 0.4 lb per gallon of dicamba acid.

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted un detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN - Probable mucosal damage may contraindicate the use of gastric lavage.	

See inside booklet for additional PRECAUTIONARY STATEMENTS


FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE CALL CHEMTREC (800) 424-9300

EPA Reg. No. 42750-76

EPA Est. No. _____

NET CONTENTS

MANUFACTURED BY:
Albaugh, LLC
1525 NE 36th Street
Ankeny, IA 50021



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing.

PERSONAL PROTECTIVE EQUIPMENT: (PPE)

All mixers, loaders, applicators, flaggers, and other handlers must wear:

Long-sleeved shirt and long pants,

Shoes and socks,

Face shield or goggles,

Waterproof gloves (except for applicators using groundboom equipment, pilots and flaggers), and

Chemical resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements and exceptions.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

ENGINEERING CONTROLS STATEMENTS:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240 (d)(6)].

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in groundwater contamination of drinking water or groundwater.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.

This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Keep people and pets out of the area during application. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to EPA Web Site: <http://www.epa.gov/espp>.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, are:

1. Coveralls worn over short-sleeved shirt and short pants,
2. Chemical-resistant footwear plus socks,
3. Waterproof gloves,
4. Chemical-resistant headgear for overhead exposure,
5. Protective eyewear.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to users of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

PRODUCT INFORMATION

Read the entire label before using this product.

Use only according to label instructions.

DDG 3-Way Herbicide is a postemergence herbicide for control or suppression of emerged weeds in fallow and reduced tillage systems, pasture and rangeland, airports, dry canals, industrial plant sites, parking areas, parks, schools, storage areas, farmsteads, ditches, fencerows, roadside and highway rights-of-way, other public areas, and similar industrial and noncrop sites. DDG 3-Way Herbicide can also be applied prior to planting or emergence of wheat, barley, corn, oats, sorghum (grain or forage), through hooded sprayers between rows of corn, as a spot treatment in corn, sorghum, forage grasses, as a preharvest treatment in wheat, and as a postharvest treatment following grain harvest.

This product enters the plant through the foliage and moves throughout the plant. Visual effects of control are a gradual wilting or yellowing of the plant, which advances to complete browning of aboveground growth and deterioration of affected underground plant parts. Visible symptoms will usually develop on labeled weeds within 2 to 4 days after application, but may not occur for 7 or more days. Extremely cool or cloudy weather following treatment may slow activity of this product and delay the visual effects of control.

Keep people and domestic animals off treated areas until foliage has dried.

APPLICATION RESTRICTIONS

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

- Delay planting of wheat, barley, oats, or sorghum (grain or forage) for 15 days after application of this product.
- Do not plant any crop other than wheat, barley, corn, oats or sorghum (grain or forage) for 3 months after treatment or until disappearance of this product from the soil is confirmed with a Field Bioassay.
- Field Bioassay Instructions: In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application. The test area should sample field conditions such as soil texture, soil pH, drainage, and any other variable that could affect the seed bed of the new crop. The field bioassay can be initiated at any time between harvest of the treated crop and the planting of the rotational crop. Observe the test crop for herbicidal activity, such as poor stand (effect on seed germination) chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the test rotational crop; plant only a labeled crop or crop listed in the table above for which the rotational interval has clearly been met.
- Do not harvest, feed or forage treated vegetation for 8 weeks after application.
- Following spot treatment in forage grasses, allow 37 days before harvesting or 30 days before grazing domestic livestock.
- Do not use aerial equipment to apply this product when sensitive crops and plants are growing in the vicinity of area to be treated.
- DDG 3-Way Herbicide is subject to all state and county regulations for 2,4-D and dicamba.

The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate, dicamba or 2,4-D as the active ingredient, whether applied as mixtures or separately. Calculate application rates and ensure that the total use of this and other glyphosate, dicamba or 2,4-D containing products does not exceed the stated maximum use rate.

APPLICATION PRECAUTIONS

- This product can be used for control of emerged weeds prior to establishment of labeled crops. Large amounts of green or decaying vegetation left standing or incorporated into the seedbed may enhance the development of disease in newly planted crops. This may result in poor emergence and/or stands, especially under cool and/or wet conditions.
- Applications of this product after planting and prior to crop emergence may cause crop injury if rainfall or environmental conditions delaying crop emergence are experienced.
- Spraying early to control young weeds before dense stands develop or light cultivation to assist weed decay will favor preparation of suitable seedbeds.
- In reduced tillage and no-till systems, ensure good seed to soil contact and proper seeding depth.
- This product may cause injury to desirable trees, plants or crops, especially sensitive broadleaf plants such as beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tomatoes and other broadleaf plants if brought into contact with their roots, stems or foliage. These plants are most sensitive during their active growth and development stage.
- Applications should be made only when there is no hazard from spray drift, since very small quantities of spray, which may not be visible, may severely injure susceptible crops or desirable vegetation.
- Movement of this product on soil particles during windstorms may cause damage to susceptible plants that are contacted. This hazard is reduced if rainfall occurs shortly after application.
- Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

TIMING OF APPLICATION

This product should be applied postemergence to vigorously growing weeds when they have reached the recommended size given in the "RATES AND WEEDS CONTROLLED" section of this label. Application should be delayed until maximum emergence of the target weeds, but before weeds exceed the maximum size recommended. For annual weeds, allow 1 day after treatment before tillage. For field bindweed, allow at least 7 days after treatment before tillage.

Reduced control may result if applications are made during poor growing conditions such as drought stress, disease or insect damage or if weeds have been mowed, grazed or cut. Heavy dust on foliage or an overstory canopy covering targeted weeds may also reduce control.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash this product off the foliage and a repeat application may be required.

APPLICATION EQUIPMENT

This product may be applied using either ground or aerial spray equipment. Use extreme care to avoid misting or drifting of herbicide solution onto foliage, green stems or fruit of desirable crops, trees, or plants during both growing and dormant periods since even very small quantities of spray can cause severe plant injury.

GROUND APPLICATION: Apply specified rates of this product in 3 to 10 gallons of water per acre as a broadcast spray unless otherwise directed in this label. For optimum spray distribution and coverage, use flat fan or low-volume flood nozzles. When using flood nozzles, space them no more than 40 inches apart and ensure double overlap of spray pattern. Refer to the manufacturer's recommendations for correct pressure and nozzle height above the target canopy. Avoid pressure and nozzles which produce fine droplets or mist.

Use appropriate marking devices to ensure uniform spray coverage and best results from this product.

HAND-HELD and HIGH-VOLUME EQUIPMENT (use coarse sprays only): Mix this product in clean water and apply to foliage of vegetation to be controlled. Do not spray to the point of runoff.

AERIAL APPLICATION: Apply the specified rates of this product in 3 to 5 gallons of water per acre as a broadcast spray unless otherwise directed on this label. **DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT WAS NOT INTENDED.**

HOODED SPRAYERS: This product may be used through hooded sprayers for weed control between the rows of corn (field, pop, seed, and silage). DDG 3-Way Herbicide is not registered for use on sweet corn. Do not apply DDG 3-Way Herbicide to seed corn or popcorn without first verifying with your local seed corn company (supplier) the DDG 3-Way Herbicide selectivity on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties. Only hooded sprayers that completely enclose the spray pattern may be used.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 30 ounces of this product per acre per application.
- Corn must be at least 12 inches tall, measured without extending leaves.
- Do not apply after tasseling.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 mph.
- Maximum wind speed: 10 mph.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

Do not graze or feed corn forage or fodder following applications of this product through hooded sprayers.

Do not apply more than 102 fl. oz. of this product per acre per year for hooded sprayer applications. Allow two or more weeks between applications of DDG 3-Way Herbicide.

Do not apply DDG 3-Way Herbicide when soybeans are growing nearby if any of these conditions exist:

- Corn is more than 24 inches tall
- Soybeans are more than 10 inches tall
- Soybeans have begun to bloom

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial equipment and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For aerial equipment, the boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made in a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

For ground boom application, do not apply with a nozzle height greater than 4 feet above the crop canopy.

MIXING INSTRUCTIONS

DDG 3-Way Herbicide ALONE

Fill the spray tank to about $\frac{3}{4}$ of the desired volume with clean water. Add the specified amount of this product, then complete the filling process while maintaining agitation. Remove the hose from the mix tank immediately after filling

to avoid siphoning back into the carrier source. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, terminate by-pass and return lines at the tank bottom and/or use an agriculturally approved antifoam or defoaming agent.

NOTE: Reduced results may occur if water containing soil is used, such as water from ponds and unlined ditches.

TANK MIXTURES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Always predetermine the compatibility of labeled tank mixtures of this herbicide with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of DDG 3-Way Herbicide with water as follows:

1. Place a 20 to 35-mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it SLOWLY through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
6. Continue filling the sprayer tank with water and add the required amount of DDG 3-Way Herbicide herbicide near the end of the filling process.
7. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water-soluble liquid.

Maintain good agitation at all times, until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle should be no finer than 100 mesh and in-line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles.

Check label of all products used in tank mix for cleaning instructions. Clean as per the hardest material to remove.

ADDITIVES

To improve burndown of emerged weeds, use surfactants or crop oil concentrate with DDG 3-Way Herbicide or DDG 3-Way Herbicide tank mixes and apply after weeds have emerged. Crop oil concentrate is for non-food/feed crop uses only. Do not apply tank mixes that include ammonium sulfate or crop oil concentrate to any food/feed crop use listed on this label. For food/feed crop use, do not use liquid fertilizers that contain ammonium sulfate (AMS) as a source of nitrogen as tolerances in commodities derived from the crop may contain residues that exceed established tolerances.

Oil Concentrate

A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be non-phytotoxic
- contain only EPA-exempt ingredients
- provide good mixing quality in the jar test, and
- be successful in local experience

The exact composition of suitable products will vary; however, vegetable oil and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

Adjuvants containing crop oil concentrates may be used for preplant, pre-emergence and between cropping applications. Do not use crop oil concentrate for postemergence applications in food/feed crops (i.e. grass (hay or silage), pastures, rangeland, and wheat.)

Non-ionic Surfactant

The addition of a non-ionic surfactant labeled for use with herbicides is recommended. The standard label recommendation is 2-4 pints of an 80% active non-ionic spray surfactant per 100 gallons of water (4 pints per 100 gallons water is equal to 1/2% non-ionic spray surfactant by volume). Rate will vary with the size and condition of weeds to be controlled. Use lowest rate per 100 gallons when weeds are small and actively growing. As weeds increase in size and or become hardened off, the rate of non-ionic surfactant will have to be increased to give optimum coverage and control.

Additive ¹	Rate Additive Per Acre
Non-ionic Surfactant	2-4 pints per 100 gallons ²
Crop Oil Concentrate	1 quart

¹ See manufacturer's label for specific rate directions.

² Use lowest rate per 100 gallons when weeds are small and actively growing. As weeds increase in size and or become hardened off, the rate of non-ionic surfactant will have to be increased to give optimum coverage and control.

Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight (or liquid equivalent) or 8.5 to 17 pounds per 100 gallons of water may increase the performance of DDG 3-Way Herbicide and DDG 3-Way Herbicide tank mixtures on annual weeds. The improvement in performance may be apparent where environmental stress is a concern. Use the higher rate of ammonium sulfate with DDG 3-Way Herbicide when treating large or dense populations of annual weeds. Low-quality ammonium sulfate may contain material that will not readily dissolve which could result in nozzle tip plugging. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet line. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. Observe all precautionary statements on the ammonium sulfate product label.

NOTE: Compatibility problems may occur at carrier volumes below 5 GPA.

SPRAYER CLEANUP

The steps listed below are suggested for thorough cleaning of spray equipment following applications of this product. Failure to clean the sprayer thoroughly may result in injury to desirable vegetation subsequently sprayed with the equipment.

1. Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full with water. Flush by operating the sprayer until the system is purged of the rinse water.
2. Fill the tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Remove the nozzles and screens and flush the system with two full tanks of water.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C 38413 may prevent corrosion.

RATES AND WEEDS CONTROLLED

For best results, apply this product after most weed seeds have germinated but before seedhead formation in grasses or flower bud formation in broadleaves.

When applied as directed, this product will provide control or suppression of the grass and broadleaf weed species listed below. Rates listed are for maximum weed height at treatment time.

PERENNIAL WEED SPECIES	DDG 3-WAY HERBICIDE	
	RATE PER ACRE (FLUID OUNCES)	LENGTH OF VINES/TREATMENT HEIGHT
Bindweed, field** Convolvulus arvensis	30 - 42 fl. oz. (suppression only)	6" to 18"
Spurge, leafy	30 - 42 fl. oz. (suppression only)	Post bloom

ANNUAL WEED SPECIES	DDG 3-WAY HERBICIDE	
	RATE PER ACRE	MAXIMUM HEIGHT
Foxtail, green Setaria viridis	22 - 32 fl. oz.	12"
Barley Hordeum vulgare Brome, downy* Bromus tectorum Cheat* Bromus secalinus Foxtail Setaria spp. Kochia* Kochia scoparia Lettuce, prickly* Lactuca serriola Oats, wild Aven fatua Puncturevine Tribulus terrestris Purslane, common Portulaco oleracea	22 - 42 fl. oz.	6"

ANNUAL WEED SPECIES	DDG 3-WAY HERBICIDE	
	RATE PER ACRE	MAXIMUM HEIGHT
Cocklebur Xanthium strumarium Lambsquarters Chenopodium album Mustard, tansy Descurainia pinnata Mustard, tumble Sisymbrium altissimum Pigweed Amaranthus spp. Rye Secale cereale Sandbur, field Cenchrus spp. Stinkgrass Eragrostis ciliaris Thistle, Russian Salsola kali Wheat Triticum aestivum Witchgrass Panicum capillare	22 - 42 fl. oz.	12"
Barnyardgrass Echinocha crus-galli Buffalobur Solanum rostratum Goatgrass Aegilops cylindrical Mustard, blue Chorispora tenella Panicum, fall Panicum dichotomiflorum	30 - 42 fl. oz.	6"
Oats, wild Aven fatua	30 - 42 fl. oz.	12"

*For improved control in no-till systems or heavy infestations or overwintered stands, use 30 fluid ounces. For best results on light kochia infestations, treat after the plant has passed through the woolly stage of growth and is 3 to 6 inches in height.

** Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Prepare the desired volume of spray by mixing the amount of DDG 3-Way Herbicide in clean water as shown in the following table:

SPRAY SOLUTIONS

DESIRED VOLUME	AMOUNT OF DDG 3-WAY HERBICIDE
1 gallon	1 1/3 fluid ounces
25 gallons	1 quart
100 gallons	1 gallon

2 tablespoons = 1 fluid ounce

SPOT TREATMENT

Applications in growing crops must be made prior to heading of small grains (wheat, barley, oats), grain sorghum (milo) and silking of corn.

For applications in forage grasses, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre may be treated at one time. Further applications may be made in the same area at 30-day intervals. Remove domestic livestock before application and wait 30 days after application before grazing livestock and 37 days before harvesting.

ECOFARMING SYSTEMS

The Ecofarming System consists of the following rotation: winter wheat, corn/sorghum, ecofallow.

Use the following tank mixtures for control of emerged annual weeds before planting corn or sorghum in the Ecofarming System.

DDG 3-Way Herbicide at 30 to 34 fluid ounces per acre

Plus

ATRAZINE at 0.75 to 1 pound active ingredient per acre

Plus

A preemergent grass herbicide labeled for these uses (follow label directions for specified application rates)

The above tank mixtures should be applied in 28-0-0 or 32-0-0 liquid fertilizer carrier at 20 to 30 gallons per acre. The liquid fertilizer may be diluted with water to achieve the desired carrier volume.

WEEDS CONTROLLED – The following weeds, up to a maximum height of 4 inches, will be controlled:

Brome, downy

Bromus tectorum

Cheat

Bromus secalinus

Foxtail, green

Setaria viridis

Foxtail, yellow

Setaria lutescens

Kochia

Kochia scoparia

Lettuce, prickly

Lactuca serriola

Pigweed, redroot

Amaranthus retroflexus

Thistle, Russian

Salsola kali

Wheat, volunteer

Triticum aestivum

Risk of crop injury from 2,4-D and dicamba can be reduced by applying this treatment 7 to 14 days before planting.

Refer to the specific product labels for preemergence weed control achieved by these tank mixtures.

Refer to the specific product labels for crop rotation restrictions and cautionary statements for all products in these mixtures.

ECOFARMING RESTRICTIONS:

- Do not make more than 2 applications per year.
- Minimum spray interval between applications is 30 days.

PREHARVEST APPLICATIONS TO WHEAT

This product provides weed control when applied prior to harvest of wheat. Apply when wheat is in the hard dough stage and the green cooler is gone from the nodes (joints) of the stem and at least 7 days prior to harvest.

This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section for instructions for ground and aerial applications.

NOTE: It is not recommended that wheat grown for seed be treated because a reduction in germination or vigor may occur.

WHEAT PREHARVEST RESTRICTIONS:

- Do not apply more than 62.5 fluid ounces per acre per application.
- Do not make more than 1 preharvest application per crop cycle.
- Do not feed treated straw to livestock. Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.
- Preharvest Interval: 14 days

AID TO TILLAGE

This product used in combination with preplant and conventional fallow tillage practices will provide control of downy brome, cheat, volunteer wheat, tansy mustard and foxtail.

Apply 15 fluid ounces of this product in 3 to 10 gallons of water per acre to weeds that are actively growing. Treat when weeds are less than 6 inches in height. Application must be followed by conventional tillage practices before regrowth of the treated plant occurs. Allow at least 1 day after application before tillage.

POSTHARVEST APPLICATIONS FOLLOWING GRAIN HARVEST

This product will provide control of weeds following grain harvest. Weeds should be allowed to regrow after damage incurred during harvesting operations, and to recover from environmental stress, before application of this product. Weeds should be treated prior to the heading stage of annual grasses and before broadleaf weeds exceed 24 inches in height. Ammonium sulfate will improve performance on annual weeds under stress conditions.

Weeds controlled at the 22-42 fluid ounces-per-acre rate include downy brome, foxtail, pigweed, puncturevine, stinkgrass, Russian thistle and volunteer wheat.

Weeds controlled at the 30-42 fluid ounces-per-acre rate include field bindweed, leafy spurge, barnyardgrass, buffalobur, blue mustard, fall panicum and wild oats.

DDG 3-WAY HERBICIDE PLUS ATRAZINE

Tank mixtures of DDG 3-Way Herbicide plus atrazine will provide postemergence control of listed annual weeds in fallow and reduced tillage systems. In addition, these tank mixtures will provide soil residual control of weeds listed on the atrazine label.

Annual Weeds Controlled

Barnyardgrass Echinochloa crus-galli	Lambsquarters Chenopodium album	Sandbur, field Cenchrus spp.	Witchgrass Panicum capillare
Brome, downy Bromus tectorum	Lettuce, prickly Lactuca serriola	Stinkgrass Eragrostis cilianensis	
Foxtail, green Setaria viridis	Mustard, tansy Descurainia pinnata	Thistle, Russian Salsola kali	
Kochia Kochia scoparia	Pigweed, redroot Amaranthus retroflexus	Wheat Triticum aestivum	

Apply 22 fluid ounces of this product plus 1 pound or less of the active ingredient, atrazine, per acre. Use 30 fluid ounces of this product plus 2 pounds or less of the active ingredient, atrazine, per acre. Use 34 fluid ounces of this product plus 1 to 3 pounds of the active ingredient, atrazine, per acre. In Oregon and Washington, do not exceed 1 pound of the active ingredient, atrazine, per acre. Barnyardgrass will be suppressed at the 34 fluid ounce per acre rate and will require up to 45 ounces per acre for control. The addition of ammonium sulfate is recommended to increase the performance of DDG 3-Way Herbicide plus atrazine tank mixtures.

Consult the atrazine labels for use rates, soil type, planting, cropping and other restrictions, as well as other precautionary statements and use according to the most restrictive label.

These tank mixtures may be applied with ground or aerial equipment. See the "APPLICATION EQUIPMENT" section for instructions.

For ground application, apply in 3 to 10 gallons of water per acre.

For aerial application, apply in 3 to 5 gallons of water per acre.

FOR NONSELECTIVE CONTROL IN SMALL GRAIN CROPPING SYSTEMS IN SOUTH DAKOTA ONLY

Weeds Controlled

Refer to the "RATES AND WEEDS CONTROLLED" section of this label for rate specifications and annual weeds controlled.

Application Instructions

For ground applications, use 3 to 5 gallons of water per acre.

For aerial applications, use 1 to 3 gallons of water per acre.

GRAIN POSTHARVEST RESTRICTIONS:

- Do not apply more than 2 applications per year.
- The minimum retreatment interval is 30 days.

NON-CROPLAND USES

FARMSTEADS, DITCHES AND FENCEROWS

When applied as directed, this product will control downy brome, bulbous bluegrass, kochia, tumble mustard, tansy mustard and prickly lettuce, and provide suppression of crested wheatgrass, smooth brome and field bindweed.

Spray coverage should be uniform and complete. Do not spray to the point of runoff. Treat when weeds are small, actively growing and free of dust. Use 58 fluid ounces of this product in 5 to 10 gallons of water per acre for broadcast boom applications.

ROADSIDE AND HIGHWAY RIGHTS-OF-WAY

When applied as directed, DDG 3-Way Herbicide alone and tank mixtures of DDG 3-Way Herbicide plus hexazinone and/or sulfometuron will reduce vegetative growth of bahiagrass turf as well as control or partially control the following annual and perennial weeds (Refer to the "RATES AND WEEDS CONTROLLED" section of this label for application rates and annual weeds controlled):

Bindweed, field	Brome, downy
Convolvulus arvensis	Bromus tectorum
Spurge, leafy	Cheat
	Bromus secalinus
Foxtail, green	Foxtail
Setaria viridis	Setaria spp.
Cocklebur	Kochia
Xanthium strumarium	Kochia scoparia
Lambsquarters	Lettuce, prickly
Chenopodium album	Lactuca serriola
Mustard, tansy	Oats, wild
Descurainia pinnata	Avena fatua
Mustard, tumble	Puncturevine
Sisymbrium altissimum	Tribulus terrestris
Pigweed, redroot	Purslane, common
Amaranthus retroflexus	Portulaco oleracea
Pigweed, smooth	Barnyardgrass
Amaranthus hybridus	Echinocha crus-galli
Rye	Buffalobur
Secale cereale	Solanum rostratum
Stinkgrass	Goatgrass
Eragrostis cilianensis	Aegilops cylindrical
Thistle, Russian	Mustard, blue
Salsola kali	Chorisporea tenella
Wheat	Panicum, fall
Triticum aestivum	Panicum dichotomiflorum
Oats, wild	Witchgrass
Avena fatua	Panicum capillare
Barley	
Hordeum vulgare	

Make applications to bahiagrass prior to seedhead emergence or after the bahiagrass has been mowed to a uniform height of 4 to 5 inches. For best results, make applications when both bahiagrass and weeds are healthy and actively growing.

Apply 8 to 18 fluid ounces of DDG 3-Way Herbicide as a broadcast spray in 10 to 25 gallons of spray solution per acre.

OR

Apply 8 to 18 fluid ounces of DDG 3-Way Herbicide plus the manufacturer's specified rate of hexazinone plus the manufacturer's specified rate of sulfometuron as a broadcast spray in 10 to 25 gallons of spray solution per acre.

OR

Apply 8 to 18 fluid ounces of DDG 3-Way Herbicide plus 8 to 16 fluid ounces (1/2 to 1 pint) of hexazinone as a broadcast spray in 10 to 25 gallons of spray solution per acre.

OR

Apply 8 to 18 fluid ounces of DDG 3-Way Herbicide plus the manufacturer's specified rate of sulfometuron as a broadcast spray in 10 to 25 gallons of spray solution.

Read and carefully observe the label claims, precautionary statements and all information on the labels of each product used in this tank mixture. Use according to the most restrictive label directions for each product in the mixture.

INDUSTRIAL AREAS, AIRPORTS, RAILROADS, DRY CANALS, PLANT SITES, PARKING AREAS, UTILITY POWER LINES, STORAGE AREAS, AND PUBLIC AREAS AND NONCROP SITES

This product provides weed control when used as directed under the "RATES AND WEEDS CONTROLLED" section of this label, when applied to industrial areas, airports, dry canals, plant sites, parking areas, parks, schools, storage areas, other public areas, and noncrop sites.

This product is also for musk thistle control in industrial areas. Best control of musk thistle is obtained when applications are made while plants are in the rosette stage of growth. Applications should be made in the spring prior to bolting or in the fall prior to soil freeze-up. Partial control or suppression may be obtained with treatments made from bolting through flowering. For best results, plants should be actively growing and not under stress and free of dust layers on leaves.

For broadcast applications, apply 8 to 18 fluid ounces per acre of this product alone or as a tank mixture with either picloram or metsulfuron. Consult the picloram and metsulfuron labels for specified use rates and restrictions.

Read and carefully observe the label directions, precautionary statements and all other information on the labels of each product used in tank mixtures.

NON-CROPLAND RESTRICTIONS:

- Postemergence (annual & perennial weeds):
 - Do not make more than 2 applications per year.
 - Minimum spray interval between applications is 30 days.
 - "Maximum of 2.0 lbs. 2,4-D ae/acre per application.
- Postemergence (woody plants):
 - Do not make more than 1 application per year.
- Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- DDG 3-Way Herbicide contains 0.134 pounds of 2,4-D acid equivalent per pint (0.008 pounds of 2,4-D acid equivalent per fluid ounce). When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of 2,4-D acid equivalent per acre per year.
- DDG 3-Way Herbicide contains 0.05 pounds of dicamba acid equivalent per pint (0.003 pounds of dicamba acid equivalent per fluid ounce). When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pounds of dicamba acid equivalent per acre per year.
- DDG 3-Way Herbicide contains 0.21 pounds of glyphosate acid equivalent per pint (0.013 pounds of glyphosate acid equivalent per fluid ounce). When tank mixing with products that contain glyphosate, do not exceed a combined total of 8.0 pounds of glyphosate acid equivalent per acre per year.

PASTURE AND RANGELAND

This product is may be used for leafy spurge and musk thistle control in rangeland and pasture areas. It is also allowed for the control of those weeds listed in the "RATES AND WEEDS CONTROLLED" section of this label.

Best control of leafy spurge is obtained when applications are made during the seed-set growth stage. This typically occurs in late June. Desirable grass species may be injured with this application. However, injury is usually temporary and normal growth resumes soon after application. Usually colonial bentgrasses are more tolerant than creeping types. Velvetgrasses are most easily injured. Treatments will kill or injure alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes. Do not use this application on the same area for two consecutive years because grass injury will increase.

Apply 30 fluid ounces of this product per acre as a broadcast application in 3 to 10 gallons of water per acre.

Best control of musk thistle is obtained when applications are made while plants are in the rosette stage of growth. Applications should be made in the spring prior to bolting or in the fall prior to soil freeze-up. Partial control or suppression may be obtained with treatments made from bolting through flowering. For best results, plants should be actively growing and not under stress and free of dust layers on leaves.

For broadcast applications, apply 8 to 18 fluid ounces per acre of this product alone or as a tank mixture with either picloram or metsulfuron. Consult picloram and metsulfuron labels for specific use rates and restrictions.

When using tank mixtures, refer to the tank mix product label for grazing restrictions for that product. Follow the most restrictive label. Read and carefully observe the label directions, precautionary statements and all other information on the labels of each product used in these tank mixtures.

PASTURE & RANGELAND USE RESTRICTIONS:

- Do not make more than 2 broadcast applications per year.
- Minimum spray interval between applications is 30 days.
- If grass is to be cut for hay, Agricultural Use requirements for the Worker Protection Standard are applicable.
- DDG 3-Way Herbicide contains 0.134 pounds of 2,4-D acid equivalency per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of 2,4-D acid equivalency per acre per year.
- DDG 3-Way Herbicide contains 0.05 pounds of dicamba acid equivalency per pint (0.003 pounds of dicamba acid equivalency per fluid ounce). When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pounds of dicamba acid equivalency per acre per year.
- DDG 3-Way Herbicide contains 0.21 pounds of glyphosate acid equivalent per pint (0.013 pounds of glyphosate acid equivalent per fluid ounce). When tank mixing with products that contain glyphosate, do not exceed a combined total of 8.0 pounds of glyphosate acid equivalent per acre per year.
- Do not graze lactating dairy animals on treated grass within 7 days after application. Animals being finished for slaughter that are grazing in the treated area within 30 days of treatment must be removed from the treated area 3 days before slaughter.
- Do not cut forage for hay within 37 days of application. When this product is used for spot treatments where less than 10 percent of the total grazed area will be treated, there is no grazing restriction.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not store near fertilizers, seeds, insecticides or fungicides. STORE ABOVE 40°F to keep product in solution. If crystals form, place in a warm room (72°F), allow the product to reach room temperature and roll or shake periodically until crystals have dissolved. Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by the use

according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous waste representative at the nearest EPA Regional Office for guidance.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

CONTAINER HANDLING: Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable <5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable >5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container (250 gallon & bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH or the Seller. To the fullest extent permitted by law, all such risks shall be assumed by the Buyer.

ALBAUGH warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALBAUGH MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims

based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above.

TO THE FULLEST EXTENT PERMITTED BY LAW, IN NO CASE SHALL ALBAUGH OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALBAUGH and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of ALBAUGH. No employee or agent of ALBAUGH or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

All product names, trademarks, and registered trademarks are the property of their respective owners.

060920

LABEL HISTORY

Not Part of Final printed Label

File Name	Version Mark	Comment
042750-00076.20200602.DRAFT	060220	EPA Comment Revisions
042750-00076.20200609.DRAFT	060920	EPA Comment Revisions