

42750-80 1.28.2009

1/23



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JAN 28 2009

Mr. Morris Gaskins
Registrations Manager
Albaugh, Inc.
1525 NE 36th Street
Ankeny, IA 50021

Subject: Label Notification for Pesticide Registration Notice 2007-4

Dear Mr. Gaskins,

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated August 6, 2008 and your label resubmission emailed January 15, 2009 for the product Picloram + 2,4-D Rangeland (EPA Registration Number 42750-80). The Registration Division (RD) has conducted its review of this request for its applicability under PRN 2007-4 and finds that the label changes requested fall within the scope of PRN 2007-4. The label submitted with the application has been stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please call me directly at 703-305-6249 or Steve Schaible of my staff at 703-308-9362.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 42750-80	2. EPA Product Manager J. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Picloram + 2,4-D Rangeland	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Albaugh Inc. P.O. Box 2127 Valdosta, GA 31604-2127 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: NOTIFICATION EPA Reg. No. _____ Product Name _____ JAN 28 2009	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container	<input type="checkbox"/> Glass
					<input type="checkbox"/> Paper
					Other (Specify) _____
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 2.5, 30, 250 bulk		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Morris Gaskins	Title Registrations Manager	Telephone No. (Include Area Code) 229-244-3288
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registrations Manager	
4. Typed Name Morris Gaskins	5. Date August 6, 2008	

CORPORATE OFFICE
1525 NE 36th Street
Ankeny, IA 50021
515.964.9444 - Office
800.247.8013 - Toll Free
515.964.7813 - Facsimile

ALBAUGH, INC.

Valdosta Office 3/23
P.O. Box 2127
304 Janet Street, Suite H
Valdosta, GA 31604
229.244.3288 - Office
229.244.5841 - Facsimile

FED-X

August 6, 2008

NOTIFICATION

JAN 28 2009

Document Processing Disk (NOTIFY)
Office of pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

RE: Label notification for "Pesticide Management and Disposal; Standards for Pesticide Containers and Containment"

Dear Sirs,


The enclosed submissions are draft labels submitted in response to Pesticide Regulation Notice 2007-4 for the following Albaugh registrations:

PRODUCT	EPA REG. NO.
Picloram + 2,4-D Rangeland	42750-80


Changes are noted in strikethrough for deleted text and underline/bold for added text.

Please call if you have any questions.

Regards,



Morris Gaskins
Registrations Manager
Albaugh, Inc.



Agri Star[®]
By Albaugh, Inc.

PREMIER SUPPLIER OF OFF-PATENT CROP PROTECTION PRODUCTS
www.albaughinc.com

CORPORATE OFFICE

1525 NE 36th Street
Ankeny, IA 50021
515.964.9444 - Office
800.247.8013 - Toll Free
515.964.7813 - Facsimile

ALBAUGH, INC.

Valdosta Office

P.O. Box 2127
304 Janet Street, Suite H
Valdosta, GA 31604
229.244.3288 - Office
229.244.5841 - Facsimile

4/
23

Via E-mail

January 15, 2009

Mr. Steve Schiable
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

RE: Picloram + 2,4-D Rangeland
EPA Reg. No. 42750-80


Dear Mr. Schiable,

The enclosed submission is a draft label for the above referenced registration in response to the Agency's PR Notice 2007-4 for Container Disposal revision. This notification replaces the original notification made 8/6/08 that inadvertently deleted the "puncture before disposal" instructions.

Changes are noted in ~~strikeout~~ for deleted text and underline/bold for added text.

Thank you for bringing this oversight to our attention.

Regards,



Morris Gaskins
Registrations Manager
Albaugh, Inc.

Agri Star
By Albaugh, Inc.

PREMIER SUPPLIER OF OFF-PATENT CROP PROTECTION PRODUCTS
www.albaughinc.com

EDITOR'S NOTE: 1/15/09 "marked" draft for PR Notice 2007-4 Container Disposal label notification to correct original notification made 8/6/08

Restricted Use Pesticide
May injure (phytotoxic) susceptible, non-target plants. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

PICLORAM + 2,4-D RANGELAND

For the control of broadleaf annual and perennial weeds, and certain woody species on Conservation Reserve Program (CRP) acres, rangeland and permanent grass pastures.

ACTIVE INGREDIENTS:

- Picloram: 4-amino-3,5,6-trichloropicolinic acid, triisopropanolamine salt 10.2%
- 2,4-dichlorophenoxyacetic acid, triisopropanolamine salt 39.6%
- OTHER INGREDIENTS: 50.2%
- TOTAL: 100.0%

Picloram + 2,4-D Rangeland contains the following acid equivalents:
picloram: 4-amino-3,5,6-trichloropicolinic acid - 5.7% (0.54 lb./gal.)
2,4-dichlorophenoxyacetic acid - 21.2% (2 lbs./gal.)

NOTIFICATION

JAN 28 2009

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN - Probable mucosal damage may contraindicate the use of gastric lavage.	

See inside booklet for additional PRECAUTIONARY STATEMENTS

EPA Reg. No. 42750-80

EPA Est. No. Xxxxxx-xx-xxx

NET CONTENTS

MANUFACTURED BY:
Albaugh, Inc.
Ankeny, IA 50021

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are butyl rubber, natural rubber, neoprene or nitrile rubber. If you want more options, follow the instructions for Category C on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

Long-sleeved shirt and long pants.

Shoes and socks.

Chemical resistant gloves when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

Chemical resistant apron when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

Protective eyewear (goggles or face shield).

For containers over 1 gallon, but less than 5 gallons: Mixers and loaders who do not use a mechanical system (such as probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to other required PPE.

See engineering controls for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this products concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

ENGINEERING CONTROL STATEMENTS

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)]

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to some plants at very low concentrations. Non-target plants may be adversely affected if pesticide is allowed to drift from areas of application.

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. Apply this product only as directed on label.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Picloram is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, picloram may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetative filter strips, and areas over-laying tile drainage systems that drain to surface water.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et. Al. v. EP, C01-0132C, (W.D. WA).

Every 2 years starting January 1, 2008, the registrant will offer training to applicators which will cover applications techniques and product stewardship particular to their use(s) of this product (Picloram + 2,4-D Rangeland, EPA Reg. No. 42750-80). Applicators of this product must be able to provide certification of such training on demand by the State, Tribal or Federal enforcement agent.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- coveralls
- shoes plus socks
- chemical-resistant gloves made of any water-proof material
- protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

GENERAL INFORMATION

Picloram + 2,4-D Rangeland is a water soluble liquid product containing picloram and 2,4-D. Use Picloram + 2,4-D Rangeland in permanent grass pastures and rangeland to selectively control many annual, biennial, and perennial broadleaf weeds and woody species listed on this label.

Herbicidal effects of Picloram + 2,4-D Rangeland occur primarily from uptake by plant foliage and translocation throughout the plant, however, secondary herbicidal activity may occur from soil uptake of picloram. Broadleaf plants can be killed or damaged by very small amounts of Picloram + 2,4-D Rangeland. To prevent damage to crops and other desirable plants, carefully follow all directions and precautions.

PRECAUTIONS AND RESTRICTIONS

Observe any special use and application restrictions and limitations, including method of application and permissible areas of use as required by state or local regulations. When used in tank mix combination with other products, follow all applicable use directions, precautions, restrictions, and limitations on the labels of each product used.

Application Rate Ranges: Use higher rates in areas with dense weed populations or for longer residual weed control. For best results, the lower rate should be used only when environmental conditions are favorable for plant growth and when the plants are in the recommended growth stage. Compared to results obtained with the higher rate, a lower rate may be slower to show activity, provide a lower level of control, and may require retreatment.

Do not rotate to crops intended for food or feed use on areas treated with this product, other than range or pasture grasses, rye, forage sorghum, sudangrass, wheat, barley or oats not underseeded with a legume.

Do not move treated soil, or use treated soil for growing other plants until soil residues of picloram are no longer detectable as indicated by an adequately sensitive bioassay or chemical test.

Do not apply Picloram + 2,4-D Rangeland in residential areas or near ornamental trees and shrubs. Untreated trees can be affected by root uptake of the herbicide through movement into the top soil or by excretion of the product from the roots of nearby treated trees. Do not apply Picloram + 2,4-D Rangeland within the area occupied by roots of desirable trees, unless such injury can be tolerated.

Established grasses are tolerant to this product, but newly seeded grasses may be injured until well established as indicated by tillering, development of a secondary root system and vigorous growth (refer to the Planting Grasses Section of this label).

Picloram + 2,4-D Rangeland may suppress certain established grasses such as smooth bromegrass, Willman's lovegrass and buffalograss. However, subsequent grass growth should be improved by release from weed competition. Smooth bromegrass and Willman's lovegrass grown for seed may be sensitive to this product if applied under adverse growing conditions (moisture stress).

If injury to existing forage legumes cannot be tolerated, do not spray pastures. Picloram + 2,4-D Rangeland may injure or kill legume plants. Forage legumes may be less sensitive to the herbicide after the seed has set and plant growth is mature. Seeding of legumes may not be successful if made within one year of application.

Do not mix with dry fertilizer.

Do not transfer livestock from treated grazing areas to broadleaf crop areas without first allowing 7 days of grazing on untreated grass pasture. Otherwise, urine may contain enough picloram to cause injury to sensitive broadleaf plants.

Do not use manure from animals grazing treated areas on land used for growing broadleaf crops, ornamentals, orchards or other susceptible, desirable plants. Manure may contain enough picloram to cause injury to susceptible plants.

Do not use hay or grass from treated areas or manure from animals being fed treated forage or hay for composting or mulching of desirable, susceptible broadleaf plants.

Do not contaminate water intended for irrigation or domestic purposes. Do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not apply to snow or frozen ground.

Do not apply or permit Picloram + 2,4-D Rangeland or sprays containing Picloram + 2,4-D Rangeland to contact crops or other desirable broadleaf plants, including but not limited to alfalfa, beans, cotton, grapes, melons, peas, potatoes, safflower, soybeans, sugar beets, sunflower, tobacco, tomatoes, and other vegetable crops, flowers, fruit plants, ornamentals and shade trees.

Do not use Picloram + 2,4-D Rangeland on sub-irrigated land.

Pasture, Rangeland and CRP Application Restrictions:

- Do not exceed 7.4 pints (0.5 lb ae picloram)/acre per application.
- Do not exceed 14.8 pints (1.0 lb ae picloram)/acre per year.
- Spot Treatment: Do not exceed 7.4 pints (1.0 lb ae picloram)/acre, with no more than 50% of an acre being treated.
- Spot treatments and broadcast treatments can be applied during the same growing season only if the total amount applied does not exceed 7.4 pints (1.0 lb ae picloram)/acre per growing season.
- Minimum of 30 days between applications.
- Maximum of two applications per year.
- If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protections Standard are applicable.
- For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.
- Picloram+2,4-D Rangeland contains 0.25 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of 2,4-D a.e. per acre per year.

Grazing Restrictions: There are no grazing restrictions for non-lactating dairy animals or other livestock including horses, sheep, goats, and other animals in the treatment area.

- Do not allow lactating dairy animals to graze treated areas within 7 days after application.
- Do not harvest grass cut for hay from treated areas for 30 days after application.
- Meat animals must be withdrawn from treated forage at least 3 days before slaughter.

SPRAY DRIFT MANAGEMENT

Avoid Injurious Spray Drift

Make applications to avoid spray drift because very small quantities of the spray that may not be visible may severely injure susceptible crops during both growing and dormant periods. To minimize spray drift:

1. Use nozzle pressures no greater than are required to obtain a proper spray pattern for adequate coverage of target plants.
2. Apply as a coarse spray.
3. Use nozzles designed for herbicide application that do not produce a fine droplet spray.
4. Spray when wind velocity is low. Follow local state regulations. Avoid application under conditions which are conducive to air inversions or conditions of atmospheric temperature inversion.

Spray drift may be further lessened when making applications near susceptible crops by using a drift control system such as Microfoil®; Thru-Valve® boom (or equivalent) or a drift control agent such as Nalco-Trol® (or equivalent). If a drift control additive is used, follow all use recommendations and precautions on the product label.

Ground Equipment: With ground equipment, spray drift may be lessened by keeping the spray boom as low as possible; by keeping the operating spray pressures at the manufacturers recommended minimum pressures for the specific nozzle types used (low pressure nozzles are available from spray equipment manufacturers). Do not apply this product with a mistblower. In hand-gun applications, spray drift may be minimized by selecting the minimum pressure that will provide adequate coverage (without forming a mist); by spraying no higher than brush tops.

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide

drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial equipment and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For aerial equipment, the boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made in a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

For ground boom application, do not apply with a nozzle height greater than 4 feet above the crop canopy.

APPLICATION DIRECTIONS

Broadcast Foliar Application (Ground or Aerial)

Unless otherwise specified, apply in water alone or in an oil-water emulsion in a total spray volume of 10 to 40 gallons per acre using ground equipment or 1 or more gallons per acre by aerial application. If aerially applied, results will be more consistent for spray volumes of 2 or more gallons per acre. Use of the lower total spray volume with ground equipment is recommended primarily where Picloram + 2,4-D Rangeland is applied simultaneously with liquid fertilizer. Good coverage is essential. For aerial application, swath width should not exceed 1 ¼ times the wingspan of the aircraft.

To provide more complete wetting and coverage of the foliage, a non-ionic surfactant may be used at recommended rates. Use a drift control additive for drift reduction and improved deposition.

Section I.

Control of Broadleaf Weeds and Woody Plants in Rangeland and Permanent Grass Pastures in:

Southwest, Southeast, and Mid-Atlantic States

1-2 Pints/Acre or 3-4 Pints/Acre: Make applications at the rate indicated by stage of growth to control the following woody plants or broadleaf weeds:	
Weed Species	Specific Use Directions
annual broomweed, bitter sneezeweed, bitterweed, buffalo bur, bull thistle, bursage (bur ragweed), camphor weed, cocklebur, common ragweed, croton, horseweed, lambsquarters, pigweed, prickly lettuce, smartweed, sunflower, tasajillo, wild carrot	<p>Early Season: Apply at a rate of 1-2 pts/acre in early to mid spring when weeds are less than 3 inches tall. Rates in the lower end of the rate range are effective only when weeds are less than 2 inches tall and conditions are favorable for plant growth.</p> <p>Mid to Late Season: Apply at a rate of 3-4 pts/acre in late spring to early summer when weeds are 3 inches tall to early flowering.</p>

2-4 Pints/Acre: Make applications at the indicated growth stage to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
aster, heath	Apply prior to bud stage when actively growing.
aster, spiny (Mexican devilweed)	Apply prior to bud stage when actively growing.
bee plant, Rocky Mountain	Apply prior to bud stage when actively growing.
bindweed, hedge	Apply prior to bud stage when actively growing.

2-4 Pints/Acre: Make applications at the indicated growth stage to control the following woody plants or broadleaf weeds:

Weed or Brush Species	Specific Use Directions
blackberry	Tank-mix 2 pints per acre of Picloram + 2,4-D Rangeland with 1 pint per acre of Triclopyr 4E herbicide plus surfactant. Apply in late May to early June during or after bloom (not before) when foliage is dark green. Do not treat blackberries in the same year after mowing, shredding, or burning. Even one year after removal of top growth, blackberry stands will be more difficult to control than undisturbed stands and will require retreatment.
buckwheat, climbing false	Apply prior to seed development when actively growing.
buckwheat, wild	Apply prior to seed development when actively growing.
bullnettle, western	Apply in spring when plants begin to flower.
bundleflower, Illinois	Apply prior to bud stage when actively growing.
burdock, common	Apply prior to bud stage when actively growing.
buttercup	Apply in early spring prior to bud stage.
chickweed, mouseear	Apply prior to bud stage when actively growing.
chicory	Apply from rosette stage to early bud stage when actively growing.
coneflower, upright prairie	Apply when plants are to 6 inches tall, but before flowering.
common goldenweed, Drummond's goldenweed (Isocoma spp.)	Apply in the spring (April-June) when favorable growing conditions result in substantial canopy development. Thorough and uniform coverage is essential. Use higher spray volumes (20-25 gpa for ground and 4-5 gpa for aerial equipment). Use of a non-ionic surfactant or oil-water emulsion is recommended (refer to the Mixing Instructions section of this label).
curly dock	Early Season: Apply 2 pints per acre prior to bolting stage of growth. Mid-to-Late Season: Apply at a rate of 3-4 pts/acre from bolting to bud stage.
devil's-claw	Apply prior to flowering when actively growing.
dogfennel (cypressweed)	Apply when plants are from 6 to 24 inches tall, but before flowering. Increase rate within the rate range as season progresses and plants become larger.
erigonum, annual	Apply prior to bud stage when actively growing.
fleabane, rough	Apply prior to bud stage when actively growing.
gray goldaster narrowleaf goldaster	Apply in the spring during the bud stage (pre-bloom) using an oil-water emulsion spray. Thorough coverage is essential.
goldenrod, Missouri	Apply prior to bud stage when actively growing.
hemlock, poison	Apply from rosette stage in spring or fall up to 36" tall.
hemlock, water (common)	Apply from rosette stage in spring or fall up to bud stage.
horsenettle, Carolina	Apply 2 pints per acre when plants are 4-6 inches tall. At 2 pints per acre, retreatment may be necessary for acceptable control. Apply 3 to 4 pints per acre when flowering or for longer residual control of later emerging plants and greater stand reduction the following year.
horehound	Apply during active growth.
jimsonweed	Apply prior to bud stage when actively growing.
morningglory, ivyleaf	Apply prior to bud stage when actively growing.
mugwort	Apply prior to bud stage when actively growing.
nightshade, silverleaf	Apply 2 pints per acre when plants are 4-6 inches tall. Apply 3 to 4 pints per acre when flowering or for longer residual control of later emerging plants and greater stand reduction the following year. Retreatment is necessary for total control.
pennycress, field	Apply when plants are to 6 inches tall, but before flowering.
plantain, buckhorn	Apply prior to bud stage when actively growing.

2-4 Pints/Acre: Make applications at the indicated growth stage to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
pricklypoppy, annual	Apply prior to bud stage when actively growing.
puncturevine	Apply prior to flowering when actively growing.
ragweed, common, giant, lanceleaf and western	Use lower rates in rate range when weeds are no more than 2 inches tall and conditions are favorable for plant growth. Use higher rates when weeds are from 3 inches tall to early flowering.
sagebrush, sand	Apply when new terminal growth reaches 6-12" and before average daytime temperature reaches 95°F. Use low rate only in early season.
snow-on-the-mountain	Apply prior to bud stage when actively growing.
sowthistle, spiny (prickly)	Apply prior to bud stage when actively growing.
stickweed	Apply 2-3 pints/acre prebloom.
thistles, biennial: including bull, musk, plumeless or scotch	Apply 2 pints/acre at rosette stage. Apply 3 to 4 pints/acre in mid to late season from bolting to bud stage.
vervain, blue vervain, hoary	Apply when plants are 6 inches tall to early flowering. Increase rate within the rate range as season progresses and weeds mature.
vetch, hairy	Apply prior to bud stage when actively growing.
wingstem	Apply 2-3 pints/acre prebloom.
yankeeweed	Apply when plants are 8 to 10 inches tall.

3-4 Pints/Acre: Make applications at the indicated growth stage to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
marshelder (sumpweed)	Apply in early season when weeds are less than 4 inches tall. Use higher rates on older plants. Thorough and uniform coverage is essential. Use higher-spray volumes (20-25 gpa for ground and 5 or more gpa for aerial equipment).
mesquite and oak sprouts (suppression of regrowth):	Delay applications of Picloram + 2,4-D Rangeland for weed control until the foliage of regrowth brush in the treatment area is fully expanded and turned from light to dark green.
milkweed	Apply 4 pints/acre to actively growing milkweeds less than 4 inches tall. Add a surfactant at the manufacturer's recommended rate to improve wetting of foliage.
mullein, common	Apply 4 pints/acre during the rosette stage in spring or fall prior to bolting. Add a surfactant at the manufacturer's recommended rate to improve wetting of foliage.
poisonous plants such as: groundsel (Senecio spp.), garbancillo (Wooton loco) and Woolly loco	Make applications in fall or winter when moisture conditions are favorable. Because locoweeds are difficult to wet, use of a surfactant (0.25-0.5% vol/vol) or oil-water emulsion is recommended (refer to the Mixing Instructions section of this label). Herbicide treatment may increase palatability of poisonous plants. Do not graze treated areas until the toxic plants are no longer palatable.
thistle, wavyleaf	Apply from rosette to late bolt stage.
tropical soda apple	Apply when plants are beginning to flower.

7.4 pints/Acre: Make applications at the indicated growth stage to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
cactus, pricklypear or cholla	Make a ground broadcast application in the spring or early summer to control a broad spectrum of broadleaf weeds in addition to pricklypear.
Chinese tallowtree	Make applications in the spring or fall when conditions are favorable for plant growth. Thorough and uniform spray coverage is required. Use higher spray volumes (20-25 gpa) for ground and 5 or more gpa for aerial equipment). Use of a non-ionic surfactant or oil-water emulsion is recommended (refer to the Mixing Instructions section of this label).
Macartney rose multiflora rose	Make applications in the spring or fall when conditions are favorable for plant growth. Thorough and uniform spray coverage is essential. Use higher spray volumes (20-25 gpa for ground and 5 or more gpa for aerial equipment). Use of a non-ionic surfactant or oil-water emulsion is recommended (refer to the Mixing Instructions section of this label). Do not make application within 9-12 months after mowing or when plants have a high percentage of new growth. Poor control will result if plants are less than 3 ft. tall.
locust (honey and black) wild plum	Make applications in the spring when leaves are fully expanded and mature. Use of a surfactant (0.25-0.5% vol/vol) is recommended.

Section II.

Control of Broadleaf Weeds and Woody Plants in Rangeland and Permanent Grass Pastures in:

North and Northwestern U.S. including Colorado, Idaho, Iowa, Kansas, Minnesota, Missouri, Montana, Nebraska, North Dakota, Oregon, South Dakota, Utah, Washington and Wyoming

For best results in terms of forage response, desirable forage grasses should be present in the area to be treated in sufficient density to provide competition to lessen weed re-establishment following treatment. Additionally, good grazing management practices are recommended, particularly in the year following treatment, to allow forage grass density to increase.

Application Rates:

Use higher rates in areas with dense weed populations or for longer residual control. For best results, the lower rate should be used only when environmental conditions are favorable for plant growth and when the plants are in the recommended growth stage. Compared to results obtained with the higher rate, a lower rate may be slower to show activity, provide a lower level of control, and may require retreatment.

2 to 4 Pints/Acre: Make applications at the indicated growth stage to control the following broadleaf plant species. Increase rate within rate range as growing season progresses:	
Weed or Brush Species	Specific Use Directions
absinth wormwood annual broomweed	Apply when actively growing in spring or early summer.
biennial thistles, such as bull, musk, plumeless or scotch	Apply 2 pints/acre at rosette stage. Apply 3 to 4 pints/acre to bolted thistle, but apply before early bud stage.
broom snakeweed	Make application after full leaf development to early bloom stage when plants are actively growing.
curly dock	Apply 2 pints/acre early season prior to bolting. Apply 3 to 4 pints/acre in mid to late season from bolting to early flower.

2 to 4 Pints/Acre: Make applications at the indicated growth stage to control the following broadleaf plant species. Increase rate within rate range as growing season progresses:

Weed or Brush Species	Specific Use Directions
curlycup gumweed	Apply when new growth and seedlings have fully emerged before bloom stage.
fringed sagebrush	Apply a minimum of 3 pints/acre after seed stalk elongation and early flowering (mid-late June) and throughout the summer under good growing conditions.
goldenrod	Apply prior to bud stage during active growth.
hemp (marijuana)	Make application from rosette stage in spring or fall up to 36" tall.
hemlock, poison	
hemlock, water (common)	Make application from rosette stage in spring or fall up to bud stage.
ironweed, western	Apply 2 to 3 pints/acre prior to bud stage during active growth. A surfactant is recommended.
locoweeds, such as silky crazyweed (white point loco) and lambert crazyweed	Apply from early bud to early bloom stage. Application of Picloram + 2,4-D Rangeland may increase palatability of these poisonous plants. Do not graze treated areas until after the toxic plants have dried up. Higher rate range should be considered to provide greater reduction of poisonous plants.
phlox, hoods	Make application during active growth.
plains pricklypear	Apply when the majority of plants are in the flower stage. The lower rate will provide a partial stand reduction. More complete control may be obtained with the higher rate. Treatment response is very slow and may continue for 2 years or longer.
ragweed, common, giant, lanceleaf and western	Use the lower rate in early season when weeds are no more than 2 inches tall. Use the higher rate when weeds range from 3 inches tall to early flowering, when conditions are favorable for plant growth.
thistles, biennial: including bull, musk, plumeless or scotch	Apply 2 pints/acre at rosette stage. Apply 3 to 4 pints/acre in mid to late season from bolting to bud stage.
vervain, blue and hoary	Apply when plants are 6 inches tall to early flowering. Increase rate within the rate range as season progresses and plants mature.
wormwood, Louisiana and absinth	Make application during active growth prior to woody stem development.
yarrow	Apply 2 pints/acre prior to bud stage. A surfactant is recommended.

4 Pints/Acre: Make applications at the indicated growth stage to control the following broadleaf weed species.

Weed or Brush Species	Specific Use Directions
dense clubmoss	Apply in early summer with a surfactant at 0.25% v/v.
geyer larkspur	Apply from rosette to flower bud formation.
hairy goldenaster	Apply at bloom stage during active growth.
houndstongue	Apply to rosettes in late fall or early summer.
larkspur, plains	Apply prior to bud stage when actively growing.
licorice, wild	Apply at bloom stage, but before bur formation.
loco, woolly	Make application from bolting to early bloom. Application of Picloram + 2,4-D Rangeland may temporarily increase palatability of this poisonous plant. Do not graze treated areas until toxic plants have dried up.
milkweed, common	Apply at bud stage when actively growing.

4 Pints/Acre: Make applications at the indicated growth stage to control the following broadleaf weed species.	
Weed or Brush Species	Specific Use Directions
mullein, common	Apply during rosette stage in spring or fall prior to bolting. Add a surfactant at the manufacturer's recommended rate to improve wetting of foliage.
oxeye daisy	Apply 3-4 pints/acre when all plants have emerged to late flowering.
pussytoes	Make application prior to bud stage when actively growing. Use a surfactant at the manufacturer's recommended rate to improve wetting of foliage.

7.4 pints/Acre: Make applications at the indicated growth stage to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
Macartney rose multiflora rose	Make application in spring or fall when conditions are favorable for plant growth. Thorough and uniform spray coverage is essential. Use higher spray volumes (20-25 gpa for ground and 5 or more gpa for aerial equipment). Use of a non-ionic surfactant or oil-water emulsion is recommended (refer to the Mixing Instructions section of this label). Do not make application within 9-12 months after mowing or when plants have a high percentage of new growth. Poor control will result if plants are less than 3 ft. tall.
locust (honey and black) wild plum	Apply in spring when leaves are fully expanded and mature. Use of a surfactant (0.25-0.5% vol/vol) is recommended.

High-Volume Foliar Applications

Spray to thoroughly wet foliage and stems. The use of an approved agricultural surfactant is recommended. Do not use more than 7.4 pints of Picloram + 2,4-D Rangeland (0.54 lb. of picloram) per acre. To minimize spray drift, use the lowest possible pressure and coarse spray to achieve good coverage. Keep sprays no higher than brush tops.

7.4 pints/100 Gallons of Spray: Make applications at the indicated growth stage to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
blackberry, elm, granjeno, locust, maple, oaks, sweetgum, sumac	Tank mix the recommended rate of Picloram + 2,4-D Rangeland with 1-2 qts./100 gallons of Triclopyr 4E and apply in late spring to early summer when leaves are fully expanded and mature. Use of a surfactant (0.25-0.5% vol/vol) is recommended. Spray to thoroughly wet foliage. For best results on blackberry, treat during or after bloom.

7.4 pints/100 Gallons of Spray: Make applications at the indicated growth stage to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
annual broomweed, bitterweed, bitter sneezeweed, bullnettle, bursage (bur ragweed), bull thistle, buffalo bur, camphorweed, cocklebur, common ragweed, croton, gray goldaster, lanceleaf ragweed, marshelder (sumpweed), musk thistle, narrowleaf goldaster, prickly lettuce, smartweed, sunflower, wild carrot, silverleaf nightshade, tasajillo, upright prairie cone flower, western hosenettle, western ragweed, yankeeweed	Make application when target weeds are 2-3 inches tall until early flowering.
flameleaf sumac honeylocust	Apply in spring when leaves are fully expanded and mature. Use of a surfactant (0.25-0.5% vol/vol) is recommended. Spray to thoroughly wet foliage.
Tropical soda apple	Apply when plant begins to flower.

*7.4-14.8 pints/100 Gallons of Spray: Make application at the indicated growth stage to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
Macartney rose multiflora rose	Make application in spring or fall when conditions are favorable for plant growth. High volume application is recommended for control of large undisturbed clumps or small regrowth.
*Do not exceed 7.4 pints of product per acre	

*14.8 pints/100 Gallons of Spray: Make application at the indicated growth stage to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
Chinese tallow tree	Make application in spring or fall when conditions are favorable for plant growth.
cactus, pricklypear or cholla	Applications may be made throughout the year. Spray to wet all pads to runoff. Use of a surfactant (0.25-0.5% vol/vol) is recommended. Water soluble dye may be added to the spray mixture to mark treated plants.
common goldenweed, Drummond's goldenweed	Apply in the spring (April-June) when favorable growing conditions result in substantial canopy development.
Poisonous plants such as: groundsel (Senecio spp.), garbancillo (Wooton loco), and Woolly loco	Make application in fall or winter when moisture conditions are favorable. Application of Picloram + 2,4-D Rangeland may increase the palatability of poisonous plants. Do not graze treated areas until the toxic plants have dried up and lost their palatability.
*Do not exceed 7.4 pints of product per acre	

Treatment After Planting Grasses, Including Conservation Reserve Program (CRP) Acres

NOTE: For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

Weed Control Prior to Seeding Planting Grasses

Picloram + 2,4-D Rangeland may be applied to control weeds prior to planting cool season grasses. Apply Picloram + 2,4-D Rangeland at 4 pints per acre or less depending on the target species. Picloram + 2,4-D Rangeland may be tank-mixed with Gly Star™ Plus (glyphosate) to control grasses prior to seeding.

- To optimize weed control, minimal disturbance of the treatment area with the seeding operation is suggested. The site should be left undisturbed for a minimum of 21 days prior to seedbed preparation or seeding. To optimize weed control and reduce the potential for injury of seeded grasses, increase the interval between application of Picloram + 2,4-D Rangeland and planting grass seed.
- Do not plant smooth brome grass for 60 days after treatment.

Perennial Grasses

Make applications of Picloram + 2,4-D Rangeland to perennial grasses only after perennial grasses are well established as indicated by vigorous growth and a well-developed secondary root system.

Sprigged Bermudagrass: Picloram + 2,4-D Rangeland at 1.5 pints per acre or less can be used on sprigged bermudagrass once the runners (stolons) have reached 6 inches in length and growing conditions are favorable.

Overseeding: Picloram + 2,4-D Rangeland at rates of 1.5 pints per acre or less can be applied to permanent pastures that have been over seeded with small grains (such as barley, forage sorghum, oats, rye, ryegrass, sudangrass or wheat) grown for pasture or hay only. Young seedling small grains or grasses are sensitive to Picloram + 2,4-D Rangeland. Do not apply Picloram + 2,4-D Rangeland until overseeded grasses are well established and at tillering stage of growth or later.

Precautions:

- Applications of Picloram + 2,4-D Rangeland to established warm season grasses such as bermudagrass during initial greenup in early spring could delay or suppress emergence of new growth. If temporary suppression of new growth cannot be tolerated, application of Picloram + 2,4-D Rangeland should be made prior to greenup or after vigorous vegetative growth has resumed.
- Do not use Picloram + 2,4-D Rangeland if legumes are a desired cover during CRP.
- Conditions unfavorable to plant growth, such as drought, will increase potential for injury to grasses at all stages of growth.
- Crop Rotation: Do not rotate to grain sorghum (milo) if greater than 4 pints per acre of Picloram + 2,4-D Rangeland has been applied. For rates below 4 pints per acre, do not plant grain sorghum for 8 months after application. This product is not intended for use on land planted to sweet sorghum. To avoid potential crop injury, planting of small grains should be delayed a minimum of 60 days of soil temperatures above 40°F following application, except in Idaho, North Dakota, Nebraska, Montana, Oregon, South Dakota, Washington and Wyoming, where the minimum interval should be 90 days.

- After CRP, do not plant broadleaf crops in treated acres until an adequately sensitive bioassay (described below) shows that no detectable picloram is present in the soil.

Field Bioassay Instructions:

In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application. The test area should sample field conditions such as soil texture, soil pH, drainage, and any other variable that could affect the seedbed of the new crop. The field bioassay can be initiated at any time between harvest of the treated crop and the planting of the rotational crop. Observe the test crop for herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the test rotational crop; plant only a labeled crop such as pasture grasses, small grains (barley, oats, rye or wheat), or, after a rotational interval of 8 months, grain sorghum.

Mixing Instructions

Ground or Aerial Application – For Use With Water Alone

Start with about half the required amount of water in the spray tank. With agitation operating, add the required amount of Picloram + 2,4-D Rangeland. If a surfactant is needed, it should be added as the remainder of the required water is added to complete the spray mix. When using a drift control additive, carefully follow the manufacturer’s directions. Complete dispersion and uniform mixing is essential to proper performance of drift control additives. This can be aided by thorough circulation through a mixing pump with moderate to high shearing action.

Use with Oil/Water Emulsions

Ground Application: Add oil to the total spray mix at a rate of 5 to 10% of the total mix, up to a maximum of 1 gallon of oil per acre, using agricultural spray emulsifiers and mixing procedures given below.

Aerial Application: Use oil and water in the spray mixture in a 1:5 ratio (1 part oil to 5 parts water), up to a maximum of 1 gallon of oil per acre using mixing procedures given below:

Mixing Instructions for Oil/Water Emulsions (Batch Mixing)

1. With continuous, vigorous agitation:
2. Add half the amount of water to be used to the spray tank
3. Add the amount of Picloram + 2,4-D Rangeland required for the total volume of spray being mixed.
4. Premix the required amount of oil with an emulsifier such as Sponto 712 or Triton X-100, using the manufacturer’s recommended rate of emulsifier per gallon of oil. Add the oil-emulsifier premix to the spray tank.
5. Finally, add the remaining amount of water required to bring the spray batch to the desired total volume.
6. Maintain agitation in the spray tank during application.

Mixing with Liquid Fertilizer for Broadcast Weed Control in Rangeland and Permanent Grass Pastures
 Picloram + 2,4-D Rangeland may be tank mixed with liquid fertilizers and used in foliar application for weed control and fertilization of rangelands and permanent grass pastures. Avoid using liquid fertilizers in applications to brush as efficacy may be reduced. Use liquid fertilizers at rates recommended by supplier or local Extension Service Specialists.

Compatibility with Liquid Fertilizer:

Prior to large scale batch mixing, conduct a "jar test" for spray mixture compatibility by mixing each component in the required order and proportion in a clear glass jar. Close the jar and agitate the mixture until evenly dispersed. Use of a compatibility agent is indicated if components of the mixture do not disperse readily or do not remain dispersed after mixing. Use of a compatibility aid such as Unite or Compex is recommended to help obtain and maintain a uniform spray solution during mixing and application. Compatibility is best with straight liquid nitrogen fertilizer solutions. Mixing with N-P-K fertilizer solutions or suspensions is more difficult and should not be attempted without first conducting a successful jar test. Agitation in the spray tank must be vigorous to compare with jar test agitation.

Suggested Mixing and Application Procedure

With continuous vigorous agitation:

1. Add half the amount of liquid fertilizer to the spray tank.
2. Add a compatibility aid such as Unite or Compex at 1 quart per 100 gallons of total spray mix.
3. First add the amount of Picloram + 2,4-D Rangeland needed for the total spray mixture. Mixing with N-P-K fertilizer solutions may be improved by premixing Picloram + 2,4-D Rangeland with water (1 part Picloram + 2,4-D Rangeland to 25-30 parts water) before adding to the spray tank.
4. Add the remaining liquid fertilizer to produce the needed total spray volume.
5. Apply as soon as mixing is complete, maintaining continuous, vigorous agitation throughout mixing and application without interruption.

Application during very cold (near freezing) weather is not advisable. The likelihood of mixing or compatibility problems with liquid fertilizer increases under cold conditions.

Do not store the spray mixture.

Note: Do not use spray equipment for application of other products to land planted, or to be planted, to susceptible crops or desirable sensitive plants, unless it has been determined that all phytotoxic herbicide residue has been removed by thorough cleaning of the equipment. See "Cleaning Instructions for Sprayer Equipment" and the PRECAUTIONS AND RESTRICTIONS sections of this label.

Cleaning Instructions for Spray Equipment

To avoid injury to susceptible plants, equipment used to apply Picloram + 2,4-D Rangeland should be thoroughly cleaned before reusing to apply any other chemicals.

1. Rinse and flush application equipment thoroughly after use. Flush the entire system at least three times with water, and dispose of rinse water in non-cropland area away from water supplies.
2. During the second rinse, add 1 qt. of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 min.). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Nozzles and screens should be removed separately.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep container tightly closed when not in use. If exposed to subfreezing temperatures, the product should be warmed to at least 40°F and thoroughly mixed before using. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

~~CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.~~

CONTAINER DISPOSAL: Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(non-refillable <5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable >5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container (250 gallon & bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, INC., its Supplemental Distributors, or the Seller. All such risks shall be assumed by the Buyer.

ALBAUGH, INC., its Supplemental Distributors and the Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER ALBAUGH, INC. NOR ITS SUPPLEMENTAL DISTRIBUTORS MAKE ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND THE EXCLUSIVE LIABILITY OF ALBAUGH, INC., ITS SUPPLEMENTAL DISTRIBUTORS AND THE SELLER FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL ALBAUGH, INC., ITS SUPPLEMENTAL DISTRIBUTORS, OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALBAUGH, INC., its Supplemental Distributors, and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of ALBAUGH, INC.

No employee or agent of ALBAUGH, INC., its Supplemental Distributor, or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

Gly Star™ is a trademark of Albaugh, Inc.

NOTIFICATION

JAN 28 2009