

42750-103

11/30/2012

1/21



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Carolyn Miter
Albaugh, Inc.
P.O. Box 2127
Valdosta, GA 31604

NOV 30 2012

Subject: Label Amendment
Product Name: Brox-M Advanced
EPA Reg. No.: 42750-103
Application Dated: August 8, 2012

Dear Ms. Miter:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

A stamped copy of the label is enclosed for your records. This label supersedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products released for shipment after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Elizabeth Fertich at (703)347-8560 or at fertich.elizabeth@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn V. Montague".

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

2/21

EDITOR'S NOTE: 8/7/12 "marked" draft label in response to label call-in for plant-back interval

BROX-M™ ADVANCED
Herbicide

ACCEPTED
NOV 30 2012
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
the Administrator of the pesticide
42750-103

FOR CONTROL OF CERTAIN BROADLEAF WEEDS IN SMALL GRAINS
(WHEAT, BARLEY, OATS AND RYE) FLAX AND GRASS GROWN FOR SOD

ACTIVE INGREDIENTS:

2-Ethylhexyl ester of 2-methyl-chlorophenoxyacetic acid*	40.0%
Octanoic acid ester of bromoxynil**(3,5-dibromo-4-hydroxybenzonitrile)	36.8%
OTHER INGREDIENTS:	23.2%
TOTAL:	100.0%

*Equivalent to or not less than 2.5 pounds MCPA acid per gallon.

**Equivalent to or not less than 2.5 pounds bromoxynil per gallon.

*** Contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN

May pose an aspiration pneumonia hazard. Contains petroleum distillate.

See additional Precautionary Statements and Directions for Use below.

EPA Reg. No. 42750-103
NET CONTENTS: 2.5 Gals

EPA Est. No. 42750-MO-001

MANUFACTURED BY:
Albaugh, Inc.
Ankeny, IA 50021

FOR CHEMICAL SPILL, LEAK, FIRE OR EXPOSURE, CALL CHEMTREC (800) 424-9300

PRECAUTIONARY STATEMENTS

WARNING

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May be fatal if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistant category selection chart.

Mixers, loaders, applicators, flaggers and other handlers must wear:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves such as barrier laminate or viton,
- Shoes plus socks

Additional PPE requirements for mixers and loaders supporting aerial application to rangelands, pasture lands, or noncropland. These mixers/loaders also must wear:

- A chemical resistant apron, and
- A NIOSH approved particulate filtering respirator equipped with N, R, or P class filter media with NIOSH approval number prefix TC-84A. It is recommended that the respirator wearer be fit tested, and trained in the use, maintenance, and limitations of the respirator.

See engineering controls for additional requirements.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)]

If you will handle a total of 60 gallons or more of this product per day, you must use a mechanical transfer system for all mixing and loading operations. If this product is packaged in a 30 gallon drum, you must use a mechanical transfer system which terminates in a drip-free hard coupling which may be used only with a spray or mix tank which has been fitted with a compatible coupling. If you do not presently own or have access to a mechanical transfer system with this type of coupling, contact your dealer for information on how to obtain such a system or to modify your present system. When using a mechanical transfer system, do not remove or disconnect the pump or probe from the container until the container has been emptied and rinsed. The pump or probe system must be used to rinse the empty container and to transfer the rinsate directly to the mixing or spray tank.

Application from a tractor with a completely enclosed cab or aerial application is required whenever this product is applied to 360 or more acres in a day. The closed systems and enclosed cabs must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)]. The handler PPE requirements may be reduced or modified as specified in the WPS.

To reduce exposure to residues, wash the spray rig, tractor, and all other equipment used to handle or apply this product with water daily or before using the equipment for any other purpose.

Chemigation: Application by chemigation must be done by fixed pipe, overhead sprinkler systems or hand moved pipe. If hand moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle.

Aerial Application: Aerial application is prohibited within 300 feet of residential areas (e.g. homes, schools, playgrounds, shopping areas, hospitals, etc.)

Do not apply with backpack or hand-held application equipment.

Apply to non-residential turf only. Do not apply to residential, playground, or schoolyard turf.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Wash thoroughly and put on clean clothing/PPE.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Combustible. Do not use or store near heat or open flame.

NOTICE: BROX-M ADVANCED contains low volatile ethyl hexyl ester of MCPA. At high air or ground surface temperatures, vapors from this product may cause injury to susceptible plants. This fact should be considered when applying BROX-M ADVANCED.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated crops during the restricted entry interval (REI). For all crops except turf, the REI is 24 hours.

The REI for harvesting sod farm turf is 26 days. For uses on turf grown for transplanting (e.g. on sod farms), notify workers of the application by warning them orally and by posting signs at entrances to treated areas.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

1. Coveralls over long-sleeved shirt and long pants
2. Shoes plus socks and
3. Protective eyewear.
4. Chemical resistant gloves such as nitrile, viton or barrier laminate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store at temperatures above 40 F and below 100° F. If allowed to freeze, remix before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable <5 gallons) Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non refillable >5 gallons) Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing

nozzle inside of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container (250 gallon & bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinseate into application equipment or rinseate collection system. Repeat this rinsing process two more times.

PRODUCT INFORMATION

BROX-M ADVANCED is formulated as an emulsifiable concentrate containing the equivalent of 2.5 lbs. per gallon of octanoic acid ester of bromoxynil and 2.5 lbs. per gallon of 2-ethylhexyl ester of MCPA.

BROX-M ADVANCED is a selective postemergence herbicide for control of important broadleaf weeds infesting wheat, barley, oats, rye, flax and grass grown for sod. Optimum weed control is obtained when BROX-M ADVANCED is applied to actively growing weed seedlings. BROX-M ADVANCED is primarily a contact herbicide, therefore thorough coverage of the weed seedlings is essential for optimum control.

BROX-M ADVANCED has little residual activity. Therefore subsequent flushes of weeds will not be controlled by the initial treatment. Generally crops that form a good canopy will help shade subsequent weed flushes.

Occasional transitory leaf burn may occur. The temporary leaf burn is similar to that seen with liquid fertilizer. Because the activity of BROX-M ADVANCED is mainly contact, recovery of the crop is generally rapid with no lasting effect. Frequency and amount of leaf burn may be greater when crops are stressed by abrasive winds, cool to cold evening temperatures or mechanical injury, such as that caused by hail, sleet or insect feeding. To reduce the potential for temporary leaf burn, applications should be made to dry foliage in the recommended spray volumes per acre when weather conditions are not extreme.

MIXING, LOADING AND HANDLING INSTRUCTIONS

2.5 Gallon Containers

Special care must be taken in mixing and loading this product. Hands should be placed on the container in such a way as to avoid possible drip or splash. Contact your local Albaugh representative if you have questions regarding the correct procedure for mixing and loading.

Bulk Containers

If you will handle a total of 48 gallons or more of this product per day, you must use a mechanical transfer system for all mixing and loading operations. If this product is packaged in a 30 gallon drum, you must use a mechanical transfer system which terminates in a drip-free hard coupling which may be used only with a spray or mix tank which has been fitted with a compatible coupling. If you do not presently own or have access to a mechanical transfer system with this type of coupling, contact your dealer for information on how to obtain such a system or to modify your present system. When using a mechanical transfer system, do not remove or disconnect the pump or probe from the container until the container has been emptied and rinsed. The pump or probe system must be used to rinse the empty container and

to transfer the rinsate directly to the mixing or spray tank.

BROX-M ADVANCED ALONE: Fill the spray tank 1/2 to 3/4 full with clean water. Begin agitation and add the specified amount of BROX-M ADVANCED. Add water to the spray tank to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

TANK MIXTURES

BROX-M ADVANCED may be tank-mixed with other pesticide products provided that these products are registered for use on the crop/use site to be treated. The tank mix must be used in accordance with the more restrictive pesticide label limitations and precautions. No label dosage rates may be exceeded. BROX-M ADVANCED cannot be mixed with any product containing a label prohibition against such mixing.

BROX-M ADVANCED can be applied in tank mixture with many other herbicides and insecticides registered for use on approved crops. Refer to the specific crop section for rate recommendations and other restrictions. To apply BROX-M ADVANCED in mixture with another product, fill the spray tank 1/2 to 3/4 full with clean water and begin agitation. If tank mixing with wettable powder, soluble powder, flowable or dry flowable products, add the powder or flowable product first. After the other herbicide is thoroughly mixed with water add the recommended amount of BROX-M ADVANCED and add water to the spray tank to the desired level. If tank mixing with other product types, add the BROX-M ADVANCED first before adding the other product. Always mix one product in water thoroughly before adding another product or compatibility problems may occur.

Maintain sufficient agitation while mixing and during application to ensure a uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application.

If tank mixing with products other than those listed within each crop section, a compatibility test should be done to ensure satisfactory spray preparation. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing. To ensure maximum crop safety and weed control, follow all cautions and limitations on this label and the labels of products used in the tank mixture with BROX-M ADVANCED.

SPRAYABLE LIQUID FERTILIZERS AND SPRAY ADDITIVES

BROX-M ADVANCED can be applied in combination with sprayable liquid fertilizer or spray additives such as surfactants or crop oil concentrate. When tank mixing with liquid fertilizer always add the fertilizer to the spray tank first and agitate thoroughly before adding BROX-M ADVANCED. Always predetermine the compatibility with liquid fertilizer by mixing small proportional quantities in advance. Agitation must be maintained during filling and application operations to ensure that BROX-M ADVANCED is evenly mixed with the fertilizer. Leaf burn may occur when BROX-M ADVANCED is applied with liquid fertilizer, but new leaves are not adversely affected.

NOTICE: Fertilizers and spray additives can increase foliage leaf burn when applied with BROX-M ADVANCED. Do not apply fertilizers or spray additives with this product if leaf burn is a major concern due to environmental conditions, crop or variety sensitivity to BROX-M ADVANCED. If BROX-M ADVANCED is mixed with liquid fertilizer, the fertilizer should compose no more than 1/2 the total spray mix.

APPLICATION PROCEDURES

BROX-M ADVANCED can be applied to registered use areas by ground, aerial and sprinkler irrigation equipment.

GROUND APPLICATION

Select a spray volume and delivery system that will ensure thorough and uniform spray coverage. For optimum spray distribution and thorough coverage use of flat fan nozzles (maximum tip size 8008) with a spray pressure of 40-60 psi are recommended. Other nozzle types and lower spray pressures that product coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control. Raindrop® nozzles and flood nozzles are not recommended as weed control with BROX-M ADVANCED may be reduced. A spray volume of 10 to 20 gallons per acre (GPA) is recommended for optimum spray coverage. A minimum of 5 GPA with a minimum spray pressure of 50 psi and a maximum ground speed of 10 mph may be used with higher speed, low volume ground application if ground terrain, crop and weed density allow effective spray distribution. When using higher speed equipment, a maximum ground speed of 10 mph is suggested if field conditions cause excessive boom movement during application which results in poor spray coverage. Ground applications made when dry, dusty field conditions exist may provide reduced weed control in wheel track areas. Applications using less than 10 gallons per acre may result in reduced weed control.

When weed infestations are heavy, use of higher spray volumes and spray pressures will be helpful in obtaining uniform weed coverage. If you are unsure of the infestation level or size of crop, consult your local agronomist or extension service.

Do not apply when winds are gusty or when other conditions favor poor spray coverage and/or off target spray movement.

AERIAL APPLICATION

Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage. Use a minimum spray volume of 5 GPA and a maximum pressure of 40 psi. A minimum spray volume of 3 gallons per acre may be used if crop canopy and weed density allow adequate spray coverage. Aerial applications using less than 5 gallons of spray volume per acre may result in reduced weed control.

Do not apply during inversion conditions, when winds are gusty or when other conditions favor poor spray coverage and/or off target spray movement. Off target spray movement can be minimized by increasing the spray volume per acre and not applying when winds exceed 10 mph.

SPRINKLER IRRIGATION APPLICATION

BROX-M ADVANCED can be applied through sprinkler irrigation systems to wheat, barley, oats, rye and grasses grown for sod.

Apply BROX-M ADVANCED through sprinkler systems including center pivot, lateral move, side (wheel) roll, solid set or hand move irrigation systems only. If hand moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle. Do not apply this product through any other type of irrigation system.

SPECIFIC REQUIREMENTS FOR APPLICATION THROUGH AUTOMATED SPRINKLER IRRIGATION SYSTEM

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to

- prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
 4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
 5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
 6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
 7. Do not apply when wind speed favors drift beyond the area intended for treatment.
 8. Agitation should be maintained in the pesticide supply tank when applying.
 9. BROX-M ADVANCED should be applied continuously for the duration of the water application with center pivot and continuous lateral move systems. Application of BROX-M ADVANCED should be made during the last 30-45 minutes of the irrigation set with other overhead sprinkler systems.
 10. For best performance, set the sprinkler system to deliver approximately 0.5 inch or less of water per acre.
 11. Remove scale, pesticide residues and other foreign matter from the supply tank and entire injector system. Flush with clean water.
 12. If BROX-M ADVANCED is diluted in the supply tank, fill the tank with half of the water amount desired, add the BROX-M ADVANCED and then add remaining water amount with agitation. Always dilute with at least 4 parts water to 1 part BROX-M ADVANCED.
 13. Start the sprinklers and then inject BROX-M ADVANCED into the irrigation line. BROX-M ADVANCED should be injected with a positive displacement pump into the main line at least 8 feet ahead of a right angle turn to insure adequate mixing. Refer to the BROX-M ADVANCED label for detailed information on application rates and timings.

CHEMIGATION USE RESTRICTIONS AND PRECAUTIONS

- Application of more than 0.5 inch/acre of irrigation water may result in decreased product performance on certain soils.
- Do not apply when conditions favor drift, when system connections or fittings leak, or when nozzles do not provide uniform distribution.
- Allow sufficient time for pesticide to be flushed through all the lines and nozzles before turning off irrigation water.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Do not connect an irrigation system used for pesticide application to a public water system.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- A person knowledgeable of the chemigation system and responsible for its operations, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when wind speed is 2-10 mph at the application site.

FOR AERIAL APPLICATION:

- The distance of the outer most nozzles on the boom must not exceed 75% of the wingspan or 90% rotor blade diameter.
- Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

FOR GROUND BOOM APPLICATION:

- Do not apply with a nozzle height greater than 4 feet above the crop canopy.

SWATH ADJUSTMENT: When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of application area by adjusting the path of the aircraft upwind.

TEMPERATURE INVERSIONS: Do not make applications during a temperature inversion because drift potential is high.

SENSITIVE AREAS: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitats for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

CONTROLLED WEED LIST

Postemergence application of BROX-M ADVANCED will control the following weeds when sprayed in the seedling stage. Maximum weed stage of growth is listed under BROX-M ADVANCED directions.

MOST SUSCEPTIBLE BROADLEAF WEED SPECIES

Annual sowthistle (<i>Sonchus oleraceus</i>)	Lanceleaf sage (<i>Salvia reflexa</i>)
Black mustard (<i>Brassica nigra</i>)	London rocket (<i>Sisymbrium irio</i>)
Black nightshade (<i>Solanum nigrum</i>)	Marshelder (<i>Iva xanthifolia</i>)
Common cocklebur (<i>Xanthium strumarium</i>)	Pennsylvania smartweed (<i>Polygonum strumarium</i>)
Common lambsquarter (<i>Chenopodium album</i>)	Pepperweed spp. (<i>Lepidium</i> spp.)
Common tarweed (<i>Hemizonia congesta</i>)	Red root pigweed (<i>Amaranthus retroflexus</i>)
Cow cockle (<i>Saponaria vaccaria</i>)	Russian thistle (<i>Salsola kali</i>)
Cutleaf nightshade (<i>Solanum triflorum</i>)	Shepherdspurse (<i>Capsella bursa-pastoris</i>)
Eastern Black Nightshade (<i>Solanum ptycanthum</i>)	Silverleaf nightshade (<i>Solanum elaeagnifolium</i>)
Coast fiddleneck (<i>Amsinckia intermedia</i>)	¹ Sunflower (<i>Helianthus annuus</i>)
Field pennycress (<i>Thlaspi arvense</i>)	Tall Waterhemp (<i>Amaranthus tubersulatus</i>)
Green smartweed (<i>Polygonum scabrum</i>)	Tartary buckwheat (<i>Fagopyrum tataricum</i>)
Hairy nightshade (<i>Solanum sarachoides</i>)	Tumble mustard (<i>Sisymbrium altissimum</i>)
Horned Poppy (<i>Glaucium corniculatum</i>)	Wild buckwheat (<i>Polygonum convolvulus</i>)
Jimsonweed (<i>Datura stramonium</i>)	Wild mustard (<i>Sinapsis arvensis</i>)
Ladysthumb (<i>Polygonum persicaria</i>)	Yellow rocket (<i>Barbarea vulgaris</i>)

¹For control of sunflower, delay application until first sunflower seedlings emerging are 4 inches in height.
SUSCEPTIBLE BROADLEAF WEED SPECIES¹

Blue (purple) mustard (<i>Chlorispora tenella</i>)	Mayweed (<i>Anthemis cotula</i>)
Common groundsel (<i>Senecio vulgaris</i>)	Prostrate knotweed (<i>Polygonum aviculare</i>)
Common ragweed (<i>Ambrosia artemisiifolia</i>)	Puncture vine (<i>Tribulus terrestris</i>)
Corn chamomile (<i>Anthemis arvensis</i>)	Redroot pigweed (<i>Amaranthus retroflexus</i>)
Corn gromwell (<i>Lithospermum arvense</i>)	Smooth pigweed (<i>Amaranthus hybridus</i>)
Fumitory (<i>Fumaria officinalis</i>)	Spiny pigweed (<i>Amaranthus spinosus</i>)
Giant ragweed (<i>Ambrosia trifida</i>)	Tall morningglory (<i>Ipomoea purpurea</i>)
Hemp sesbania (<i>Sesbania exaltata</i>)	Tansy mustard (<i>Descurainia pinnata</i>)
Henbit (<i>Lamium amplexicaule</i>)	Tarweed (<i>Hemizonia spp.</i>)
Ivyleaf morningglory (<i>Ipomoea hederacea</i>)	Velvetleaf (<i>Abutilon theophrasti</i>)
Knawel (<i>Scleranthus annuus</i>)	Wild radish (<i>Raphanus raphanistrum</i>)
Kochia (<i>Kochia scoparia</i>)	

¹Weeds germinating after spraying will not be controlled.

WEED SUPPRESSION

Canada Thistle (*Cirsium arvense*)

BROX-M ADVANCED applied at 1 1/2 pints per acre provides burn down of top growth. Regrowth may occur. Make applications when Canada thistle is 8 inches tall to the bud stage. Refer to the tank mix directions on this label for optimum suppression options.

PLANT BACK INTERVAL

Wheat, barley, oats, rye, flax and peas treated with MCPA may be replanted with any crop specified on an MCPA label or any crop for which a residue tolerance exists for MCPA. For crops not listed on an MCPA label, or on crops for which no residue tolerances for MCPA have been established, a 1-year plantback interval must be observed.

WHEAT, BARLEY, OATS AND RYE BROX-M ADVANCED DIRECTIONS

PRODUCT	BROX-M ADVANCED RATE			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
BROX-M ADVANCED	4/5	12.8	10	Fall seeded wheat, barley, oats and rye throughout the United States and spring seeded wheat, barley, oats and rye in Idaho, Oregon, Washington, Colorado,	Most Susceptible Broadleaf Weeds: Apply to weeds up to the 8 leaf stage or 4 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter.

12/21

PRODUCT	BROX-M ADVANCED RATE			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
	1 1/5 – 1 3/5	19.2 – 25.6	6.7 – 5	Wyoming and Montana. Apply to wheat, barley, oats and rye from the 3 leaf stage but before the crop reaches the boot stage.	Susceptible Broadleaf Weeds: Apply to weeds up to the 4 leaf stage or 2 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 1 inch in diameter.
	1 3/5	25.6	5		Apply to henbit, knawel and mayweed up to the 4 leaf stage or 2 inches in height, whichever comes first. Apply to kochia and tansy mustard for improved control when these weeds exceed the recommended stage of growth or are growing under cool, dry conditions.
	4/5 – 1 1/5	12.8 – 19.2	10 – 6.7	Spring seeded wheat and barley except Idaho, Oregon, Washington, Colorado, Montana, and Wyoming. Apply to wheat, barley, oats and rye from the 3 leaf stage but before the crop reaches the boot stage.	MOST SUSCEPTIBLE AND SUSCEPTIBLE BROADLEAF WEEDS: Apply to weeds that do not exceed the 8 leaf stage or 4 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter. Apply to kochia up to 2 inches in height.
	1 1/5 – 1 3/5	19.2 – 25.6	6.7 - 5	Spring seeded wheat and barley except Idaho, Oregon, Washington, Colorado, Montana, and Wyoming. Apply to wheat, barley, oats and rye from the 3 leaf stage but before the crop reaches the boot stage.	Apply to kochia that is 2-4 inches in height.
	Chemigation Only 1 3/5	25.6	5	Apply to wheat, barley, oats and rye from the 3 leaf stage but before the boot stage. Apply through automated sprinkler irrigation systems with mechanical transfer loading system only. See Engineering Controls Section for details.	Apply to MOST SUSCEPTIBLE and SUSCEPTIBLE broadleaf weeds up to the 4-leaf stage, 2 inches in height or 1 inch in diameter, whichever comes first.

PRODUCT	BROX-M ADVANCED RATE			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
	Post-harvest 3/5 – 1 3/5	9.6 – 25.6	13.3 – 5	Make applications following harvest of wheat, barley, oats and rye in the states of North Dakota, South Dakota, Minnesota, and Montana. Do not plant any rotational crop until the following use season.	Apply 3/4 to 1 pint/A to MOST SUSCEPTIBLE BROADLEAF WEEDS up to the 8 leaf stage or 4 inches in height, whichever comes first. Apply 1 1/2 to 2 pints/A to SUSCEPTIBLE BROADLEAF WEEDS up to the 4 leaf stage or 2 inches in height, whichever comes first. For control of both grasses and broadleaf weeds, tank mix BROX-M ADVANCED with Roundup® or other labeled brands of glyphosate + 2,4-D.

**BROX-M ADVANCED
TANK MIXTURE DIRECTIONS**

PRODUCT	BROX-M ADVANCED RATE			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
BROX-M ADVANCED + MCPA ester (Based on 4 lb. per gallon AI)	3/5 – 1 3/5 + ¼ - ½ pint/A	9.6 -25.6	13.3 - 5	Apply to spring seeded wheat, barley, oats and rye from tillering stage but before boot stage.	For control of MOST SUSCEPTIBLE and SUSCEPTIBLE weeds and improved control of redroot pigweed and kochia. Apply to weeds up to the 8 leaf stage, 3 inches in height or 2 inches in diameter, whichever comes first. Apply to kochia and redroot pigweed up to 2 inches in height or diameter.
BROX-M ADVANCED + Starane	3/5 – 1 3/5 + 1/3 - 2/3 pint/A	9.6 -25.6	13.3 - 5	Apply to spring seeded wheat, barley, oats and rye from the 2 leaf stage up to and including flag leaf emergence.	Enhances the control of kochia up to 4" (including ALS resistant). Apply to kochia up to 4 inches in height or diameter.

PRODUCT	BROX-M ADVANCED RATE			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
BROX-M ADVANCED + 2,4-D ester (Based on 4 lb. per gallon Ai)	3/5 – 1 3/5 + 1/4 – 1 pint/A	9.6 -25.6	13.3 - 5	Apply to spring seeded wheat, barley and rye after grain is fully tillered (usually about 4 to 8 inches high) but before it is forming joints in the stem. Do not apply to grain in boot to dough stage.	For control of MOST SUSCEPTIBLE and SUSCEPTIBLE weeds and improved control of redroot pigweed, wild buckwheat and kochia (including ALS-resistant weeds.). Apply to weeds up to the 8 leaf stage, 3 inches in height or 2 inches in diameter, whichever comes first. Apply to kochia and red root pigweed up to 2 inches in height or diameter.
BROX-M ADVANCED + Banvel® Clarity® or Dicamba DMA Salt (based on 4 lb. per gallon Ai)	3/5 – 1 1/5 + 1/8 - 1/4 pint/A	9.6 – 19.2	13.3 – 6.7	FOR USE ON WHEAT ONLY. DO NOT TREAT BARLEY, OATS OR RYE. Fall seeded wheat from the 3 leaf stage but before jointing. Spring seeded wheat from the 3 to 5 leaf stage of growth.	This tank mix improves control of broadleaves such as prostrate knotweed and kochia (including ALS-resistant weeds.). Apply to weeds up to the 8 leaf stage, 3 inches in height or 2 inches in diameter, whichever comes first. Apply to kochia up to 2 inches in height or diameter.
BROX-M ADVANCED + Glean® (refer to Glean label for adjuvant recommendation)	3/5 – 1 1/5 + 1/6 - 1/3 oz/A	9.6 – 19.2	13.3 – 6.7	Apply to wheat and barley from the 3 leaf stage but before the crop reaches the boot stage. Refer to Glean® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as Kochia, henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
BROX-M ADVANCED + Finesse® (refer to Finesse label for adjuvant recommendation)	3/5 – 1 1/5 + 1/6 - 1/3 oz/A	9.6 – 19.2	13.3 – 6.7	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Finesse® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as Kochia, henbit, tansy mustard and chickweed. Apply to weeds up to the 8 leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.

PRODUCT	BROX-M ADVANCED RATE			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
BROX-M ADVANCED + Ally® (refer to Ally label for adjuvant recommendation)	3/5 – 1 1/5 + 1/10 oz/A	9.6 – 19.2	13.3 — 6.7	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Ally® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as Kochia, henbit, tansy mustard and chickweed (including LS-resistant weeds). Apply to weeds up to the 8 leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
BROX-M ADVANCED + Peak® (refer to Peak label for adjuvant recommendation)	3/5 – 1 1/5 + 0.25 oz/A	9.6 – 19.2	13.3 — 6.7	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Peak® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as Kochia, henbit, tansy mustard and chickweed (including ALS-resistant weeds). Apply to weeds up to the 8 leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
BROX-M ADVANCED + Harmony® Extra or Harmony® GT (refer to Harmony or Harmony GT label for adjuvant recommendation)	3/5 – 1 1/5 + 3/10 - 1/2 oz/A	9.6 - 19.2	13.3 — 6.7	Winter wheat. Apply from the 3 leaf stage but before the 3 rd node is detectable. Refer to the Harmony® Extra or GT label for crop rotation and other restrictions. Spring wheat and barley. Apply from the 3 leaf stage but before the crop reaches the boot stage. Spring Oats: From the third leaf stage of crop, but before jointing. Refer to the Harmony Extra or GT label for crop Rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as Kochia, henbit, chickweed and Redroot pigweed. Apply to weeds up to the 8 leaf stage, 4 inches in height or across, whichever comes first.

PRODUCT	BROX-M ADVANCED RATE			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
BROX-M ADVANCED + Amber® (refer to the Amber label for adjuvant recommendation)	3/5 – 1 1/5 + 0.14 - 0.56 oz/A	9.6 — 19.2	13.3 — 6.7	Apply to wheat and barley from the 3 leaf stage, but before the crop reaches the boot stage. Refer to the Amber® label for crop rotation and other restrictions.	This tank mix improves control of broadleaves such as Kochia, henbit, tansy mustard, and pigweed. Apply to weeds up to the 4 leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
BROX-M ADVANCED + Express® (refer to the Express label for adjuvant recommendation)	3/5 – 1 1/5 + 1/12-1/3 oz/A	9.6 – 19.2	13.3 – 6.7	Wheat and barley. Apply from the 3 leaf stage but before the crop reaches the boot stage. Refer to the Express® label for crop rotation and other restrictions.	This tankmix improves control of broadleaf weeds such as Kochia, henbit, chickweed, redroot pigweed and suppression of Canada thistle. Apply to annual weeds up to the 8 leaf stage, 4 inches in height or across, whichever comes first and to Canada thistle 4 to 8 inches tall with 2 to 6 inches of new growth.
BROX-M ADVANCED + Curtail® or Curtail® M	3/5 – 1 1/5 + 1-2 pints/A 3/4 - 1 pints/A	9.6 – 19.2	13.3 – 6.7	Apply to wheat and barley after the crop begins to tiller up to the 1 st node detectable.	This tankmix improves suppression of Canada thistle. Apply to annual broadleaf weeds up to the 8 leaf stage, 4 inches in height or 2 inches in diameter and to Canada thistle in the rosette to prebud stage.
BROX-M ADVANCED + Metribuzin (Sencor® or Lexone®)	4/5 + 1/8 - 3/16 lb ai/A	12.8	10	Winter wheat in Idaho, Oregon and Washington. Apply in spring after growth has started and secondary roots with a minimum of 3 to 4 tillers have been established, but before the forming of joints in the stem. Avoid application when crop has experienced winter kill, frost damage, disease or drought.	This tankmix improves control of broadleaf weeds such as chickweed, filaree, henbit. Apply to weeds up to the 4 leaf stage, 2 inches in height or diameter, whichever comes first. A recognized authority should be consulted concerning the use of this mixture in your area.

PRODUCT	BROX-M ADVANCED RATE			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
BROX-M ADVANCED + Avenge®	4/5 – 1 3/5 + 2 ½ - 4 pints/A	12.8 – 25.6	10 - 5	Winter wheat. Four leaf to tillering stage. Refer to Avenge® label for varietal and other restrictions. Spring Wheat. Five to 6 leaf stage. Refer to Avenge® label for varietal and other restrictions. Barley. Three to 7 leaf stage.	This tankmix will provide wild oat control in addition to broadleaves. Apply to wild oats in the 3-5 leaf stage and broadleaves that do not exceed the 4 leaf stage or rosettes of 1.5 inches in diameter. Avenge use rates per acre are 2 ½ pints (1-10 oats per sq. ft.), 3 pints (11-25 oats per sq. ft.) or 4 pints (more than 25 oats per sq. ft.).
BRONATE Advanced ^{1M} + Assert®	3/5 – 1 1/5 + 1 - 1 1/2 pints/A	9.6 – 19.2	13.3 – 6.7	Apply to wheat and barley from the 3 leaf stage but before boot stage. Refer to Assert® label for crop rotation and other restrictions.	This tankmix will provide wild oat control in addition to broadleaf weeds. Apply to wild oats at the 1-4 leaf stage and broadleaf weeds up to the 8 leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first. Use Assert at 1 ½ pints/A west of the Rocky Mountains or if wild oats have initiated tillering. For spray volumes in excess of 10 GPA, add 0.3 fluid oz of nonionic surfactant for each gallon in excess of 10 GPA.
BRONATE Advanced + Puma®	4/5 + 1/3 - 2/3 pints/A 1 1/5 + 2/3 pints/A	12.8 19.2	10 6.7	Apply to wheat and barley from the 3 leaf stage but before boot. No closer than 60 days prior to harvest in MN, MT, ND and SD, and no closer to harvest than 70 days in all other states. Do not use this tankmix on two-row malting barley. For use on winter wheat only in the states of Washington, Oregon, and Northern Idaho.	This tankmix will control broadleaf weeds, green foxtail and foxtail millet. If the higher Puma rate is used, additional grasses controlled include wild oats and barnyard grass. Yellow foxtail will be suppressed.

PRODUCT	BROX-M ADVANCED RATE			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
BROX-M ADVANCED + Discover®	3/5 – 1 3/5 + 3.2-4 oz/A	9.6 - 25.6	13.3 - 5	Refer to the Discover® label for proper rates, crops, adjuvants, rotation restrictions and application timing information.	Use the higher end of the rate range specified for larger broadleaf weeds. Use minimum spray volume of 10 GPA by ground and 5 GPA by air.
BROX-M ADVANCED + Everest®	4/5 + 0.61 oz/A	12.8	10	Refer to the Everest label for proper rates, crops, adjuvants, rotation restrictions and application timing information.	This tank mix will control broadleaf weeds, green foxtail and foxtail millet.
BROX-M ADVANCED + Achieve®	4/5 – 1 3/5	12.8 - 25.6	10-5	Refer to the Achieve label for proper rates, crops, adjuvants, rotation restrictions and application timing information.	This tank mix will control broadleaf weeds, green foxtail and wild oats.
BROX-M ADVANCED + Maverick®	4/5 – 1 3/5 + 0.66 oz.	12.8 - 25.6	10 - 5	Refer to the Maverick label for proper rates, crops, adjuvants, rotation restrictions and application timing information.	This tank mix will control broadleaf weeds plus grasses as listed on the Maverick label.

Precautions: Wheat, Barley, Oats and Rye

- Reduced weed control may occur when weeds are stressed from lack of moisture or cold temperatures.
- Refer to labels of products used in tank mixture for additional restrictions and precautions.

Restrictions: Wheat, Barley, Oats and Rye

- Do not graze treated fields within 45 days after application.
- Do not apply more than 1.6 pints of BROX-M ADVANCED per acre in a single growing season.
- Do not plant rotational crops within 30 days following Brox-M application. Wheat, barley, oats and rye treated with MCPA may be replanted with any crop specified on an MCPA label or any crop for which a residue tolerance exists for MCPA. For crops not listed on an MCPA label, or on crops for which no residue tolerances for MCPA have been established, a 1-year plantback interval must be observed.
- Do not apply more than 0.75 lbs. of MCPA acid equivalent per year (2.4 pints of BROX-M ADVANCED) when tank mixing with products that contain MCPA.

GRASSES GROWN FOR SOD PRODUCTION
BROX-M ADVANCED DIRECTIONS
Seedling and Established Grasses

PRODUCT	RATE Per ACRE	RATE Per 1000 SQ.FT.	APPLICATION TIMING AND SPECIFIC COMMENTS	
			CROP	WEEDS
BROX-M ADVANCED	4/5 - 1 3/5 Pints	0.3 to 0.6 Fl. Oz.	Apply to established and newly seeded grasses grown for sod production before the boot stage. Established grasses tolerant to BROX-M ADVANCED include bentgrasses, Kentucky Bluegrass, Fescues, Ryegrass, Bermudagrass, St. Augustinegrass and Zoysiagrass. BROX-M ADVANCED may also be used on seedling grasses such as Merion, Park, Delta, or common Kentucky Bluegrasses, Pennlawn, Chewings, Illahee or Alta Fescues, Orchard grass, Highland, Seaside or Astoria Bentgrasses, perennial Ryegrasses, Bahiagrass and Zoysiagrass.	Refer to the GENERAL WEED LIST for a listing of susceptible broadleaf weeds. Optimal control will be attained when weeds are treated in the seedling stage (less than 4 leaf stage, 2 inches in height, or 1 inch in diameter).
BROX-M ADVANCED	Chemigation 1 3/5 pints/A only	0.6 Fl. Oz.	Apply to established and newly seeded grasses grown for sod production before the boot stage. Apply through automated sprinkler irrigation systems with mechanical transfer loading system only. See MIXING LOADING AND HANDLING INSTRUCTIONS section for complete details. Refer to the list of established grasses that are tolerant to BROX-M ADVANCED	

RESTRICTIONS: Grasses grown for sod production

- Do not allow livestock to graze in treated areas or feed treated grasses to livestock.
- Do not apply BROX-M ADVANCED to grasses grown for sod production with backpack or hand-held application equipment.
- Do not apply more than 1.6 pints of BROX-M ADVANCED per acre in a single growing season.
- Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.
- Do not apply more than 1.5 lbs acid equivalent per acre per year (4.8 pints of BROX-M ADVANCED)

FLAX (*Linum usitatissimum* only)
BROX-M ADVANCEDD DIRECTIONS

PRODUCT	BROX-M ADVANCEDD RATE -			APPLICATION TIMING AND SPECIFIC COMMENTS	
	Pints/A	Fl. Oz/A	Acres/Gal	CROP	WEEDS
BROX-M ADVANCED	5/7	11.4	11.3	Apply to flax that is 2 to 8 inches in height. Do not apply to flax during or after the bud stage.	Apply to MOST SUSCEPTIBLE weeds that do not exceed the 4 leaf stage, 2 inches in height or 1 inch in diameter, whichever comes first.

HIGHER SPRAY VOLUMES OF 15 TO 20 GALLONS PER ACRE WILL DECREASE POTENTIAL FOR FLAX INJURY.

PRECAUTIONS: Flax (*Linum usitatissimum* only)

- Unacceptable crop injury may occur following BROX-M ADVANCED application to flax grown on high organic, peat type soils.
- Application under high humidity conditions can injure flax.
- Unless otherwise instructed, do not apply BROX-M ADVANCED to flax with crop oil concentrate, surfactants or nitrogen solutions.
- Flax treated with MCPA may be replanted with any crop specified on an MCPA label or any crop for which a residue tolerance exists for MCPA. For crops not listed on an MCPA label, or on crops for which no residue tolerances for MCPA have been established, a 1-year plantback interval must be observed.

RESTRICTIONS: Flax (*Linum usitatissimum* only)

- Do not apply if temperatures are expected to exceed 85° F at application or 3 days following application or crop injury may occur.
- Do not use on ornamental flax.
- Do not apply more than 0.72 pint of BROX-M ADVANCED per acre in a single growing season.
- Do not exceed .25 lbs acid equivalent per acre per year (0.8 pints BROX-M ADVANCED)

WARRANTY LIMITATIONS AND DISCLAIMER

To the extent consistent with applicable law, all such risks shall be assumed by the buyer. Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the DIRECTIONS FOR USE when used under normal conditions. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL. Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify seller, in writing, of any claims to be eligible to receive either remedy given below. To the extent consistent with applicable law, the EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of seller will be one of the following, at the election of the seller:

1. Refund of purchase price paid by buyer or user for product bought or
2. Replacement of amount of product used.

To the extent consistent with applicable law, the seller will not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations and Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations and Disclaimer in any manner.

BROX-M ADVANCED is a trademark of Albaugh, Inc.

Assert and Avenge are registered trademarks of American Cyanamid Company.

Sencor is a registered trademark of Bayer AG, Germany.

Raindrop is a registered trademark of Delavan Corporation.

Curtail is a registered trademark of DowElanco.

Ally, Express, Finesse, Glean, Harmony and Lexone are registered trademarks of E.I. DuPont

DeNemours and Co. Banvel is a registered trademark of Sandoz Crop Protection Corporation.

Amber is a registered trademark of Ciba-Geigy.

Rhonox, WEEDAR and WEEDONE are registered trademarks of NuFarm.

Roundup is a registered trademark of Monsanto Company.