



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 17, 2024

Tasha Lott
Product Registration Manager
Albaugh, LLC
1525 NE 36th Street
Ankeny, IA 50021

Subject: Label Amendment - Registration Review Mitigation for Quinclorac
Product Name: QUINCLORAC 75 SWF
EPA Registration Number: 42750-131
Application Date: January 21, 2022
Decision Number: 581389

Dear Tasha Lott:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Quinclorac Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling

before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


If you have any questions about this letter, please contact Caleb Carr by phone at (202) 566-0636, or via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington".

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label



QUINCLORAC 75 SWF

For weed control in fallow systems, grass grown for seed, pre-plant wheat (see use directions for geographic limitations), pre-plant and in-crop sorghum, and non-crop areas in the following states: CO, DE, ID, IL, KS, MD, MN, MO, MT, ND, NE, NM, NV, OK, OR, PA, SD, UT, WA, WY, VA and designated counties in TX.

ACTIVE INGREDIENT:

3,7-dichloro-8-quinolinecarboxylic acid 75.0%

OTHER INGREDIENTS: 25.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN.

CAUTION

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
• Have person sip a glass of water if able to swallow.
• Do not induce vomiting unless told to do so by a poison control center or doctor.
• Do not give anything by mouth to an unconscious person.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
• Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
• Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
• Rinse skin immediately with plenty of water for 15-20 minutes.
• Call a poison control center or doctor for treatment advice.

IF INHALED:

- Move person to fresh air.
• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
• Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

In Case of Emergency regarding this product, call: CHEMTREC 800-424-9300

EPA Reg. No. 42750-131

EPA Est. NO. 11773-IA-001

NET CONTENTS:

Manufactured For:
ALBAUGH, INC.
ANKENY, IA 50021

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Harmful if absorbed through skin or inhaled. Avoid breathing dust or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Wash thoroughly with soap and water after handling. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Do not apply directly to water, areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and Conditions of Sales and Warranty are to be followed. This labeling must be in the user's possession during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in cool, dry and well ventilated area. Do not store containers under wet conditions.

PESTICIDE DISPOSAL: Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse non-refillable < 50 pounds as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse non-refillable > 50 pounds as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.


GENERAL INFORMATION

QUINCLORAC 75DF SWF herbicide is intended for use in fallow systems, grass grown for seed, preplant

Control Recommendations

For most effective bindweed control, apply QUINCLORAC 75DF SWF herbicide in the fall just prior to the first killing frost. Bindweed plants should be actively growing and at least 4 inches long. If tillage is a part of local post-harvest practices, allow a minimum of 30 days for bindweed plants to re-grow after tillage prior to application. For best long term bindweed control, make yearly applications of QUINCLORAC 75DF SWF at 5.3 – 8.0 ounces per acre in the Fall. Use the higher rate for dense populations or large plants.

Figure 1. Application Region for QUINCLORAC 75DF SWF



QUINCLORAC 75DF SWF may be used in the following counties of Texas: Archer, Armstrong, Bailey, Baylor, Borden, Briscoe, Brown, Callahan, Carson, Castro, Chidress, Clay, Cochran, Coke, Coleman, Collin, Collingsworth, Concho, Cooke, Cottle, Crosby, Dallam, Dawson, Deaf Smith, Denton, Dickens, Donley, Fisher, Floyd, Foard, Garza, Glasscock, Gray, Grayson, Hale, Hall, Hansford, Hardeman, Hartley, Haskell, Hemphill, Hockley, Hutchinson, Jack, Jones, Kent, King, Know, Lamb, Lipscomb, Lubbock, Lynn, McCulloch, Montague, Moore, Motley, Nolan, Chiltree, Oldham, Parmer, Potter, Randall, Roberts, Runnels, Schackelford, Scurry, Sherman, Sterling, Stonewall, Swisher, Taylor, Terry, Throckmorton, Wheeler, Wichita, Wilbarger, Wise, Yoakum, and Young.

Be sure to obtain and follow all Texas state requirements for QUINCLORAC 75DF SWF uses.

Mode of Action

QUINCLORAC 75DF SWF is a systemic herbicide with plant uptake occurring through both the foliage and roots. Resultant herbicide symptoms on susceptible plants include twisting, stunting, reddening and chlorosis. For annual plants, symptoms may take up to two weeks after application to develop with death occurring in about three weeks. For perennial weeds, symptoms may not be evident for several weeks after application and full effect may not be evident for 3 to 6 months.

Coverage

When making postemergence applications, weeds must be thoroughly covered with spray because foliar uptake of QUINCLORAC 75DF SWF by the target weed is important for optimum control. Large leaf canopies shelter smaller weeds and can prevent adequate spray coverage.

Cleaning Spray Equipment

Clean spray equipment thoroughly using a strong detergent or commercial sprayer cleaner according to

the manufacturer's directions before and after applying this product.

APPLICATION INSTRUCTIONS

Based on the uses described in this label, QUINCLORAC 75DF SWF should be applied by ground application equipment. QUINCLORAC 75DF SWF may be applied using aerial application equipment only when permitted by supplemental labeling. The supplemental labeling must be in the possession of the user at the time of aerial application.

QUINCLORAC 75DF SWF may be applied as either a broadcast or spot spray application. Applications must be made to actively growing weeds.

For most broadleaf weeds, the most effective control will result from applying QUINCLORAC 75DF SWF early, when weeds are small. Delaying application permits weeds to exceed the maximum size and may prevent adequate control. In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth.

Ground Application (Broadcast)

Water Volume:

Use 5-30 gallons of water per broadcast acre. When weed foliage is dense, higher spray volumes may be required.

Spray Pressure:

Use a maximum of 30 psi (measured at the boom, not at the pump or in the line).

Application Equipment:

Use only nozzles that will produce uniform spray patterns and thorough coverage, spaced up to 20 inches apart. Select nozzles designed to produce minimal amounts of fine spray particles. Do not use controlled droplet applicator (CDA) nozzles as erratic coverage can cause inconsistent weed control. Apply as a medium or coarser spray (ASABE standard 572). Do not use selective application equipment such as recirculating sprayers or wiper applicators. Drift reduction nozzles such as Delavan® Raindrop Drift Reduction Flat Spray Tip, RF Tips, XR Tee Jet™ Extended range Flat Spray Tips, or other brands of comparable capabilities are recommended.

SPRAY ADDITIVES

To achieve consistent weed control, the use of spray additive(s) with QUINCLORAC 75DF SWF is required. The recommended spray additive with QUINCLORAC 75DF SWF is methylated seed oil. The use of crop oil concentrate with QUINCLORAC 75DF SWF is also permitted. A nitrogen fertilizer source (AMS or UAN) can be added to enhance efficacy, but cannot be used in place of methylated seed oil or crop oil concentrate. Refer to Table 2. Spray Additive Rate Per Acre for spray additive rates.

Table 2. Spray Additive Rate per Acre

SPRAY ADDITIVE	GROUND APPLICATION
Methylated Seed Oil	1.0 – 2.0 pints ²
Crop Oil Concentrate	2.0 pints
AMS ¹	2.5 pounds
UAN Solution ¹	0.5 – 1 gallon

¹ Optional
² For best grass control, use at least 1.5 pints/acre of Methylated seed oil.

METHYLATED SEED OIL OR CROP OIL CONCENTRATE:

A methylated seed oil or crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be non-phytotoxic,
- contain only EPA-exempt ingredients,
- provide good mixing quality in the jar test, and
- be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

For additional information, see Compatibility Test for Mix Components.

For bindweed control in Oklahoma, New Mexico and the designated counties of Texas, the use of methylated seed oil plus AMS is mandatory with QUINCLORAC 75DF SWF herbicide when it is applied alone.

NITROGEN FERTILIZER SOURCE:

- Urea ammonium nitrate (UAN): Commonly referred to as 28%, 30%, or 32% nitrogen solution. Do not use brass or aluminum nozzles when spraying UAN.
- Ammonium sulfate (AMS): AMS may be substituted for UAN. Use high-quality AMS (spray grade) to avoid plugging of nozzles. Other sources of nitrogen are not as effective as those mentioned, Albaugh does not recommend applying AMS if applied in less than 10 gallons per acre because of potential problems with precipitation in reduced volumes. Use AMS only if it has been demonstrated to be successful in local experience. Because most nitrogen solutions are mildly corrosive to galvanized, mild steel, and brass spray equipment, rinse the entire spray system with water soon after use. Use high-quality AMS to avoid plugging spray nozzles. The AMS must be readily soluble in water and contain no insoluble materials. Local sources of high-quality, fine, feed-grade AMS may be better than fertilizer grade. Low-quality AMS may contain material that will not readily dissolve, which could result in nozzle tip plugging. To determine AMS quality, perform a jar test adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If any un-dissolved sediment is observed, pre-dissolve the AMS in water and filter before adding it to the spray tank. If the AMS is added directly to the spray tank, add slowly while agitating. Adding the mix too quickly may clog outlet lines.

Nonionic Surfactant:

Alternatively, an 80% active nonionic spray surfactant may only be used when QUINCLORAC 75DF SWF is tank mixed with other products that restrict the use of oil additives. However, the use of nonionic surfactant may result in reduced weed control with QUINCLORAC 75DF SWF. The standard label recommendation for nonionic surfactant is 1 quart per 100 gallons of water (0.25% vol./vol.). Applications with nonionic surfactant require the addition of a nitrogen fertilizer source.

GENERAL TANK MIXING INFORMATION

Read and follow the applicable Restrictions and Limitations and Directions For Use on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

Tank Mix Partners/Components

Use the following tank mixes to achieve control of the weeds listed as suppressed in Table 1. The following herbicides may be tank mixed with QUINCLORAC 75DF SWF herbicide according to the specific tank mixing instructions in this label and respective product labels. For all recommended tank mixes, use a rate of 5.3 – 8.0 ounces per acre of QUINCLORAC 75DF SWF.

Albaugh does not recommend using tank mixes other than those listed on Albaugh labeling. Physical incompatibility, reduced weed control, or crop injury may result from mixing QUINCLORAC 75DF SWF with other pesticides, additives, or fertilizers. Local agricultural authorities may be a source of information when using other than Albaugh recommended tank mixes.

- 2,4-D
- Atrazine
- Buctril® (bromoxynil)
- Brox 2E (bromoxynil)
- Buctril®/Atrazine (bromoxynil + atrazine)
- Brox AT (bromoxynil + atrazine)
- Clarity® (dicamba)
- Dicamba DMA (dicamba)
- Cyclone® (paraquat)
- Distinct® (diflufenzopyr + dicamba)
- Fallowmaster® (glyphosate + dicamba)
- Fallow Star® (glyphosate + dicamba)
- Frontier® (dimethenamid)
- Gly Star Plus (glyphosate)
- Guardsman® Max (dimethenamid-P+atrazine)
- Landmaster® (glyphosate + 2,4-D)
- Marksman® (dicamba + atrazine)
- Outlook® (dimethenamid-P)
- Peak® (prosulfuron)
- Range Star (dicamba + 2,4-D)
- Roundup® RT (glyphosate)
- Roundup® Ultra (glyphosate)
- Weedmaster® (dicamba + 2,4-D)

Compatibility Test for Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

1. Water: For 20 gallons per acre spray volume, use 3-1/3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
2. Products in PVA bags: Cut an opening in the water-soluble PVA bag just large enough to use a teaspoon for measuring purposes. Use the opened water-soluble PVA bag first when preparing spray solution. Cap the jar and invert 10 cycles.
3. Water-dispersible products including QUINCLORAC 75DF SWF, such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions: For the 5.3-ounce rate, use 1 teaspoon. For the 8.0-ounce rate, use 1.5 teaspoons. Cap the jar and invert 10 cycles.
4. Water-soluble products: Cap the jar and invert 10 cycles.
5. Emulsifiable concentrates: (methylated seed oil or crop oil concentrate when applicable). Cap the jar and invert 10 cycles.
6. Water-soluble additives (AMS or UAN when applicable): Cap the jar and invert 10 cycles.
7. Let the solution stand for 15 minutes.
8. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. Do not use any spray solution that could clog spray nozzles.

Mixing Order

1. Water. Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
2. Agitation. Maintain constant agitation throughout mixing and application.
3. Inductor. If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA bags: Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water-dispersible products (including QUINCLORAC 75DF SWF herbicide), dry flowables, wettable

powders, suspension concentrates, or suspo-emulsions.

6. Water-soluble products.
7. Emulsifiable concentrates (such as oil concentrate when applicable).
8. Water-soluble additives (such as AMS or UAN when applicable).
9. Remaining quantity of water, Maintain constant agitation during application.

GENERAL RESTRICTIONS AND LIMITATIONS

- Maximum application rate: Do not apply more than 16 ounces of QUINCLORAC 75DF SWF® herbicide (equivalent to 0.75 pounds quinclorac) per acre per application.
- Maximum seasonal use rate: Do not apply more than a total of 16 ounces of QUINCLORAC 75DF SWF® herbicide per acre, per calendar year.
- Restricted Entry Interval (REI): 12 hours.
- Crop Rotation Restrictions: In case of crop failure, only Spring or Winter wheat or grain sorghum may be immediately replanted. Do not plant any other crop other than Spring or Winter wheat or grain sorghum for 309 days (10 months) following application. For alfalfa, clover, dry beans, flax, peas, lentils, safflower, Solanaceous crops listed below, and sugarbeets, do not replant for 24 months and conduct a bioassay prior to planting any of these crops.
- Do not apply to weeds or grasses under stress due to lack of moisture, herbicide injury, mechanical injury or cold temperatures, as unsatisfactory control may result.
- Do not apply to crops subjected to stress conditions such as hail damage, flooding, drought, injury from other herbicides, or widely fluctuating temperatures, as crop injury may result.
- Do not use selective application equipment such as recirculating sprayers, wiper applicators, or shielded applicators.
- Rainfast period: QUINCLORAC 75DF SWF is rainfast 6 hours after application.
- Wind Speed for Ground Applications; Do not apply QUINCLORAC 75DF SWF when wind speeds are greater than 10 mph at the application site.
- Do not apply QUINCLORAC 75DF SWF using aerial equipment unless permitted by supplemental labeling.
- For ground applications: Do not release spray at a height greater than 30 inches above the ground.
- Do not apply through any type of irrigation equipment.

DRIFT:

Do not allow QUINCLORAC 75DF SWF to drift onto other desirable plants, especially sensitive crops belonging to the following plant families:

1. Solanaceae [tomato, potato, tobacco, eggplant, peppers (Capsicum), among others]
2. Umbelliferae (celery, parsley, carrots, among others)
3. Leguminosae (alfalfa, green bean, among others)
4. Convolvulaceae (sweet potato, among others)
5. Chenopodiaceae (spinach, sugar beet, among others)
6. Malvaceae (okra, among others)
7. Cucurbitaceae (watermelon, cantaloupe, squash, pumpkin, among others)
8. Compositae (lettuce, sunflowers, among others)
9. Linaceae (flax)

- Do not allow spray containing QUINCLORAC 75DF SWF to drift onto areas where tomatoes are to be planted, have been planted, or onto emerged tomatoes, as severe injury will occur.
- Do not use QUINCLORAC 75DF SWF in tank mixes not specified on this label or Albaugh technical bulletins.
- Do not premix QUINCLORAC 75DF SWF with fungicides, herbicides, insecticides, additives, or fertilizers as contamination of mixing equipment and movement of QUINCLORAC 75DF SWF to off-site mixing areas can occur.

CROP SPECIFIC INFORMATION

GRASS GROWN FOR SEED

For use in the following grasses grown for seed:

Cool Season Grasses:

Bromegrass; smooth, meadow, smooth X meadow cross
European Dunegrass
Fescue; fine, tall
Junegrass
Kentucky Bluegrass
Quackgrass
Needlegrass; green
Orchardgrass
Ryegrass; annual, indian, perennial
Wheatgrass; bluebunch, crested, fairway, fairway X crested cross, intermediate, pubescent, siberian, slender, tall, thickspike, western, bluebunch X quack cross
Wildrye; altai, basin, beardless, dariurian, mammoth, Russian

Warm Season Grasses:

Bermudagrass
Bluestem; big, little, sand
Grama; blue, side-oats
Sandreed; prairie
Switchgrass

Apply QUINCLORAC 75DF SWF herbicide at 5.3 ounces per acre for control of annual grasses and broadleaf weeds (see Table 1). Apply QUINCLORAC 75DF SWF for bindweed control after grass seed harvest and hay removal but before the first killing frost. Refer to the section entitled "Field and Hedge Bindweed Control Recommendations" for use directions,

Tank Mixing Information:

Other registered products may be tank mixed with QUINCLORAC 75DF SWF, Read and follow the applicable Restrictions and Limitations and Directions For Use on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

FALLOW SYSTEMS OR PRE-PLANT WHEAT OR PRE-PLANT SORGHUM
DO NOT use pre-plant wheat in the following states: ID, MT, NV, OR, UT, WA, or WY

QUINCLORAC 75DF SWF can be applied in fallow areas or pre-plant wheat (do not apply in ID, MT, NV, OR, UT, WA or WY) or pre-plant grain sorghum at 5.3 ounces per acre for control of annual grasses and broadleaf weeds (see Table 1). For bindweed control with QUINCLORAC 75DF SWF, refer to the section entitled "Field and Hedge Bindweed Control Recommendations" for use directions. When QUINCLORAC 75DF SWF is applied as a Pre-plant treatment in wheat, plant wheat at least 1" deep.

Shallow planting (<1" deep) may result in possible crop injury when wheat is subjected to drought or other stress conditions.

Fallow Tank Mixes:

Other registered products may be tank mixed with QUINCLORAC 75DF SWF. Read and follow the

applicable Restrictions and Limitations and Directions For Use on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

- 2,4-D
- Clarity® (dicamba)
- Dicamba DMA® (dicamba)
- Distinct® (diflufenzopyr + dicamba)
- Fallow Star® (glyphosate + dicamba)
- Landmaster® (glyphosate + 2,4-D)
- Roundup RT® (glyphosate)
- Gly Star Plus (glyphosate)
- Roundup Ultra® (glyphosate)

IN-CROP SORGHUM

Apply QUINCLORAC 75DF SWF to grain sorghum at 5.3 - 8.0 ounces per acre from preemergence to postemergence (to 12 inch tall sorghum) for control of annual grasses and broadleaf weeds. For best annual grass control, QUINCLORAC 75DF SWF should be applied at 5.3 - 8.0 ounces per acre in a tank mix with atrazine at 0.5-1.0 pound a.i. per acre when weeds are less than 2" tall. Do not use liquid fertilizer as a carrier for postemergence applications of QUINCLORAC 75DF SWF to grain sorghum.

Tabel 3. Tank Mix Use Rate Per Acre with Quinclorac 75DF SWF

Herbicide Tank Mix Partner	Fallow and Preplant Wheat	Preplant Sorghum	Post-emerge Sorghum
2,4-D	0.375 - 1.0 lb ai	0.375 - 1.0 lb ai	0.125 - 0.5 lb ai
Atrazine	—	0.5 - 1.0 lb ai	0.5 - 1.0 lb ai
Clarity®	4- 16 oz.	4- 16 oz.	8 oz.
Fallow Star®	22 - 44 oz.	22 - 44 oz.	—
Landmaster®	32 - 54 oz.	32 - 54 oz.	—
Peak®	—	—	0.25 oz.
Roundup® Ultra and Gly Star Plus	12 - 32 oz.	12 - 32 oz.	—
Buctril® and Brox 2E	—	—	16 oz.
Buctril®/Atrazine and Brox –AT	----	---	32 oz.
Guardsman Max®	—	—	40 - 64 oz.

In Oklahoma, New Mexico, and in the designated counties in Texas, apply only 8.0 ounces of QUINCLORAC 75DF SWF per acre to in-crop sorghum.

NONCROP AREAS (ROADSIDES, FENCELINES AND RIGHTS-OF-WAY)

QUINCLORAC 75DF SWF may be applied to non-crop areas such as fence lines, roadsides, highway medians, utilities, railroad and pipeline rights-of-way. QUINCLORAC 75DF SWF may be applied to non-cropland areas for the control of certain weeds in the Noxious Weed Control Programs, Districts or Areas including broadcast or spot treatments. Use 5.3 - 8.0 ounces of QUINCLORAC 75DF SWF per acre for control of annual weeds, or 8.0 – 16.0 ounces per acre for other perennial weeds (see Table I), but do not exceed a total of 16.0 ounces of QUINCLORAC 75DF SWF per acre per calendar year. For bindweed control with QUINCLORAC 75DF SWF, refer to the section entitled "Field and Hedge Bindweed Control Recommendations" for use directions.

Non-Crop Tank Mixes:

Other registered products may be tank mixed with QUINCLORAC 75DF SWF. Read and follow the applicable Restrictions and Limitations and Directions For Use on products involved in tank mixing. The

most restrictive labeling applies to tank mixes.

- 2,4-D
- Clarity® (dicamba)
- Dicamba DMA (dicamba)
- Distinct® (diflufenzopyr + dicamba)
- Roundup® RT (glyphosate)
- Roundup® Ultra (glyphosate)
- Gly Star Plus (glyphosate)

Crop-Specific Restrictions and Limitations

- Do not allow livestock to graze in treated areas.
- Do not harvest hay from treated areas within 309 days after application.
- Do not feed treated grasses, forage, hay, silage, straw, seed nor seed screenings to livestock.
- Do not apply to water or to areas where surface water is present.
- Do not apply to irrigation ditches or areas that act as a channel for water entering cropland.

Crops:

This product can be used on the following crops:

- Fallow Systems
- Grass Grown for Seed
- Grain Sorghum
- Wheat (pre-plant)

Look inside for complete Restrictions and Limitations and Application Instructions.

WEEDS LISTED IN THIS LABEL	
Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bedstraw/cleavers	<i>Galium aparine</i>
Bindweed, Field hedge	<i>Convolvulus arvensis</i> <i>Calystegia sepium</i>
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>
Clover, crimson red white	<i>Trifolium incarnatum</i> <i>Trifolium pratense</i> <i>Trifolium repens</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Dandelion	<i>Taraxacum officinale</i>
Foxtail, giant green yellow	<i>Setaria faberi</i> <i>Setaria glauca</i> <i>Setaria viridis</i>
Kochia	<i>Kochia scoparia</i>

WEEDS LISTED IN THIS LABEL	
Common Name	Scientific Name
Lambsquarters, common	<i>Chenopodium album</i>
Lettuce, prickly	<i>Lactuca seriola</i>
Morningglory	<i>Ipomea spp.</i>
Ragweed, common giant	<i>Ambrosia artemisiifolia</i> <i>Ambrosia trifida</i>
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Sowthistle	<i>Sonchus oleraceus</i>
Spurge	<i>Euphorbia spp.</i>
Sunflower, wild	<i>Helianthus annuus</i>
Thistle, Canada Russian	<i>Cirsium arvense</i> <i>Salsola iberica</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Volunteer Flax	<i>Linum sp.</i>

CONDITIONS OF SALE AND WARRANTY

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Albaugh, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Albaugh, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks, referred to above.

ALBAUGH, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT ALLOWED BY LAW, IN NO CASE SHALL ALBAUGH, INC. OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. Albaugh, Inc. and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of Albaugh, Inc.

Clarity, Distinct, Frontier, Guardsman Max, Marksman, Paramount and Weedmaster are registered trademarks and Outlook is a trademark of BASF.

Buctril is a registered trademark of Rhone-Poulenc.

Cyclone is a registered trademark of Zeneca Inc.

Peak is a registered trademark of Novartis Corporation.

Fallowmaster, Landmaster, and Roundup are registered trademarks of Monsanto Company.

Delavan is a registered trademark of Garlock International Inc.

Tee Jet is a trademark of Spraying Systems Company.

121923

LABEL HISTORY

(Not included in final printed label)

File Name	Version Mark	Comment
042750-131.20220121.DRAFT	012122	Quinclorac PID
042750-00131.20230802.DRAFT	080223	(e) Label Revisions
042750-00131.20231219.DRAFT	121923	(e) Label Revisions