UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460



JUL 0 9 2009

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Morris Gaskins Albaugh, Inc. P.O. Box 2127 304 Janet Street, Suite H Valdosta, GA 31604

Dear Mr. Gaskins:

Subject:

Supplemental Labeling; Potato Seed Piece Treatment

IMIDACLOPRID 5L ST

EPA Registration No. 42750-133 Your Submission Dated: May 8, 2009

The labeling referred to above submitted in connection with the Federal Insecticide, Fungicide and Rodenticide, is amended as acceptable.

At your next label printing, or within eighteen months of the date of this letter, whichever occurs first, you must incorporate this supplemental labeling into the Section 3 primary product label for EPA Reg. No. 42750-133.

A stamped copy of the labeling is enclosed for your records. Please submit one final printed copy of the labeling before releasing the product for shipment. If you have any questions regarding this label, please contact Autumn Metzger at (703) 305-5314 or metzger.autumn@epa.gov.

Sincerely,

Venus Eagle

Product Manager 01

Insecticide-Rodenticide Branch Registration Division (7505P)

AUCEPTED

pesticide registered under:

JUL 0.9 2009 AOT D Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the

IMIDACLOPRID 5L ST EPA Reg. No. 42750-133

PRODUCT LABEL EPA. Reg. No: 42750-133

SUPPLEMENTAL PRODUCT LABEL

This label for Imidacloprid 5L ST expires on January 9, 2011 and must not be distributed or used after this date

FOR USE AS POTATO SEED PIECE TREATMENT

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling,

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions and Conditions of Sale and Warranty are to be followed.

This labeling and the Section 3 container label must be in the user's possession during application. Follow all instructions, precautions and restrictions on the container label.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds.

APPLICATION INSTRUCTIONS

Pests Controlled	Rate: Fluid ounces per 100 lbs. of Seed	Rate: Fluid ounces per acre*
Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Potato psyllid, Wireworns (seed-piece protection)	0.16 - 0.32	3.2 - 6.4
Pest/Diseases Suppressed	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Symptoms of: Potato leaf roll virus (PLRV), Potato yellows, Net necrosis	0.32	6.4

Restrictions:

- Do not apply more than 8.0 fluid ounces per acre (0.31 lb active ingredient per acre) IMIDACLOPRID 5FL ST per crop season when making seed piece treatment applications.
- Do not use treated seed-pieces for food, feed, fodder or processing for oil.
- Do not apply any subsequent in-furrow application of any imidacloprid product following a seed-piece treatment.

Application:

Apply specified dosage as a diluted spray onto seed-pieces using a shielded spray system. Dilute with 3 parts water, or less, to 1 part IMIDACLOPRID 5FL ST. Agitate or stir spray solution as needed. Fungicidal or inert absorbent dusts may be applied after IMIDACLOPRID 5FL ST application. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Plant seed pieces as soon as possible after treating avoiding prolonged exposure of IMIDACLOPRID 5FL ST treated seed-pieces to sunlight and in accordance with the specifications of your local Extension service.

* Based on a seeding rate of 2000 lbs. per acre.

MANUFACTURED FOR:

Albaugh, Inc. Ankeny, IA 50021 229-244-3288