



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

December 1, 2016

Ms. Carolyn Miter
Registrations Specialist
Albaugh, LLC
P.O. Box 2127
Valdosta, GA 31604

Subject: Notification per PRN 98-10 – align the table of contents with the headers in the body of the label to meet California DPR’s demand
Product Name: Oxyfluorfen 2EC
EPA Registration Number: 42750-136
Application Date: November 3, 2016
Decision Number: 523546

Dear Ms. Miter:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact Eleanor Thornton at 703-305-6799 or via email at Thornton.eleanor@epa.gov.

Page 2 of 2
EPA Reg. No. 42750-136
Decision No. 523546

Sincerely,

A handwritten signature in black ink, appearing to read "Erik Kraft". The signature is fluid and cursive, with a large initial "E" and a long, sweeping tail.

Erik Kraft, Acting Product Manager 24
Fungicide and Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

OXYFLUORFEN 2EC

ACTIVE INGREDIENT

Oxyfluorfen: 2-chloro-1-(3-ethoxy-4-nitrophenoxy) 4-(trifluoromethyl)	22.3%
INERT INGREDIENTS	<u>77.7%</u>
TOTAL	100.0%

Contains 2 pounds active ingredient per gallon.
Contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison Control Center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. This product may pose an aspiration pneumonia hazard. Contains petroleum distillates.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Contact CHEMTREC toll free at 1-800-424-9300 day or night, for emergency information.

EPA Reg. No. 42750-136
~~AD051814~~

EPA Est. No. xxxxxx-xx-xxx

NET CONTENTS:

Manufactured For:
ALBAUGH, INC.
ANKENY, IA 50021

NOTIFICATION

42750-136

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

12/01/2016

TABLE OF CONTENTS

Precautionary Statements

- Hazards to Humans and Domestic Animals
- Personal Protective Equipment (PPE)
- User Safety Recommendations

—~~First Aid~~

—~~Engineering Controls~~

Physical and Chemical Hazards

- Environmental Hazards

Directions for Use

- Agricultural Use Requirements

—~~Non-Agricultural Use Requirements~~

Storage and Disposal

~~Product Use Information~~

- ~~Use Application Instructions and~~ Restrictions

~~Spray Drift Restrictions~~

~~Rotation Crop Restrictions~~

~~Weeds Listed~~

~~Application Methods and Recommended Cultural Practices~~

~~Preemergence Weed Control~~

~~Postemergence Weed Control~~

~~Ground Application~~

~~Spot Application~~

~~Aerial Application~~

Chemigation

Mixing ~~Directions~~

~~Cultural Considerations~~

~~Rate Ranges~~

~~Mixing Directions~~

Crop-Specific Use Information

~~Artichokes~~ ~~Artichoke (Globe)~~

Broccoli-/Cabbage/Cauliflower

Cacao (~~Bearing and Nonbearing~~) (~~For Use in Hawaii Only~~)

Citrus (Non-Bearing)

Clary Sage

Coffee (~~Bearing And Nonbearing~~) (~~For Use Only in Hawaii~~)

Conifer ~~and Seedbeds. Transplants. Container Stock And~~ Selected ~~Field Grown~~ Deciduous Trees

Corn

Cotton

Cottonwood

Eucalyptus

~~Use On~~ Fallow ~~Bed (Cotton/Soybeans~~ Beds (Not for use prior to planting soybeans in California)

—~~Fallow land~~

Garbanzo Beans (~~For Use Only in Arizona and California~~)

Garlic

Grapes (Non-dormant Application, ~~CA~~ ~~California~~ Only) (~~See Tree fruit/Nuts/Vines Section~~)

— (~~See Tree fruit/Nuts/Vines Section~~)

— ~~Sucker Control In Non-Dormant~~ Grapes (~~Non-dormant for wine~~ ~~Washington~~ and ~~processing, OR and~~ ~~WA~~ ~~Oregon~~ Only) (~~Grapes for Wine and Processing Only~~)

— (~~See Tree fruit/Nuts/Vines Section~~)

Guava (~~Bearing and Non-bearing~~) (~~For Use Only in~~ ~~Hawaii-Only~~)

Horseradish

Jojoba

Mint (~~Spearmint and Peppermint~~)

Non-Crop ~~Use~~

Onions

Onions Grown for Seed

Papaya (Hawaii Only)

Pistachios, Almonds and Walnuts (Non-dormant Application, ~~CA~~ California only)

~~(See Tree fruit/Nuts/Vines Section)~~

Soybeans (Not For Use in California)

Taro (For use Only in Hawaii)

Tree ~~fruit~~Fruit, Nut, Vine Crops (Dormant Application)

~~Windbreaks and Shelterbelts (MN, ND, SD) For Use Only in Minnesota, North Dakota, South Dakota and WY~~
~~Only Wyoming)~~

~~Weed Control List~~

Terms and Conditions of Use

Warranty Disclaimer

Inherent Risks of Use

Limitation of Remedies

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes Substantial But Temporary Eye Injury. Causes Skin Irritation. Harmful If Swallowed Or Absorbed Through The Skin. Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals.

Do not get in eyes, on skin or on clothing. Wear goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Mixers, loaders and applicators using engineering controls (see engineering controls requirements below), must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves when mixing and loading
- Chemical-resistant apron when mixing and loading

All other mixers, loaders, applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves
- Chemical-resistant headgear when exposed overhead
- Chemical-resistant apron when exposed to the concentrate

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Pilots must use an enclosed cockpit in a manner that meets the requirement listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

Mixer and loaders supporting aerial applications to fallow land or ground applications to corn, cotton, or soybeans must use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides[40 CFR 170.240(d)(4)], and must:

- Wear the personal protective equipment required above for mixers/loaders using engineering controls,
- Wear protective eyewear if the system operates under pressure, and
- Be provided and have immediate available for use in an emergency, such as a broken package, spill, or equipment breakdown: coveralls, and chemical-resistant footwear.

Handlers performing applications to corn must use an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(5)] for dermal protection. In addition, such applicators must:

- Wear the personal protective equipment required above for applicators using engineering controls,
- Be provided and must have immediately available for use in an emergency when they must exit the cab in the treated area: coveralls, chemical-resistant gloves, chemical-resistant footwear, and chemical-resistant headgear, if overhead exposure,
- Take off any PPE that was worn in the treated area before reentering the cab, and
- Store all such PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

PHYSICAL AND CHEMICAL HAZARDS

Combustible: Do not use or store near heat or open flame.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. See Directions for Use for additional restrictions. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS:

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours, except for the following:

- Onions, garlic and horseradish: The REI is 48 hours.
- Conifer seedlings: The REI is three (3) days.
- Conifer trees: The REI is six (6) days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material,
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS:

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep unprotected persons out of treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

PESTICIDE STORAGE: Store above 40 F.

PESTICIDE DISPOSAL: Pesticide Wastes are toxic. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable > 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container (> 55 gallons): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

Steps to be Taken In Case Material is Released or Spilled: Ventilate area. Avoid breathing vapors. Use MSHA/NIOSH self-contained breathing apparatus or airline respirator for large spills in confined areas. Dike the soil with inert material (sand, earth, etc.) and transfer the liquid or solid diking material to separate containers for recovery or disposal. Remove contaminated clothing promptly and wash exposed skin areas with soap and water. Wash clothing before reuse. Keep spill out of all sewers and bodies of water.

APPLICATION INSTRUCTIONS AND RESTRICTIONS

OXYFLUORFEN 2EC herbicide may be applied for preemergence and postemergence weed control in labeled crops as indicated in this label. All use directions as provided in the Product Use Information section and crop-specific sections of this label, must be followed.

RESTRICTIONS - THE FOLLOWING USE RESTRICTIONS APPLY TO ALL REGISTERED USES OF THIS PRODUCT:

(NOTE: SEE DIRECTIONS FOR USE FOR INDIVIDUAL CROPS FOR CROP-SPECIFIC USE RESTRICTIONS.)

- Read and observe all label directions before using. When tank mixing, read and carefully follow all applicable use directions, precautions, and limitations on the respective product labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- The annual maximum application rate for all food/feed crops is 1.5 lbs ai/Acre.
- Do not contaminate irrigation water or water used for domestic purposes.
- Do not use any plants treated with OXYFLUORFEN 2EC for feed or forage.
- Do not feed or allow animals to graze on any areas treated with OXYFLUORFEN 2EC.
- OXYFLUORFEN 2EC should be applied only by ground application equipment except as otherwise allowed or directed in specific use directions.
- Do not apply when weather conditions favor drift. Avoid drift to all non-target areas. OXYFLUORFEN 2EC is phytotoxic to plant foliage.
- Do not treat ditch banks or waterways with OXYFLUORFEN 2EC.

SPRAY DRIFT RESTRICTIONS

- A 25 ft. vegetative buffer strip must be maintained between all areas treated with this product and lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.
- For ground boom application, apply with nozzle height no more than 4 feet above the ground or crop canopy and when wind speed is 10 mph or less at the application site as measured by an anemometer.
- Use coarse spray according to ASAE 572 definition for standard nozzles or VMD of 475 microns for spinning atomizer nozzles.
- The applicator also must use all other necessary to control drift.

ROTATION CROP RESTRICTIONS

Do not rotate to small-grain crops (includes barley, buckwheat, corn, pearl millet, proso millet, oats, popcorn, rice, rye, sorghum, triticale, wheat, wild rice) within 10 months following an OXYFLUORFEN 2EC treatment.

Do not direct seed any crops, other than OXYFLUORFEN 2EC- labeled crops, within 60 days following a treatment with OXYFLUORFEN 2EC.

Do not transplant seedling crops, other than OXYFLUORFEN 2EC-labeled crops, within 30 days following treatment with OXYFLUORFEN 2EC

IMPORTANT: TREATED SOIL MUST BE THOROUGHLY INCORPORATED TO A DEPTH OF 4 INCHES AFTER HARVEST (OR ABANDONING) OF THE TREATED CROP BUT PRIOR TO PLANTING OF THE ROTATIONAL CROP. FAILURE TO ACHIEVE THIS THOROUGH AND COMPLETE INCORPORATION OR TO FOLLOW THE

REQUIRED MINIMUM PLANT-BACK INTERVAL MAY RESULT IN CROP INJURY, STAND REDUCTION AND/OR VIGOR REDUCTION OF THE PLANT-BACK CROP. See specific fallow bed labeling regarding crop planting information for applicators of OXYFLUORFEN 2EC that are made to a fallow bed or fallow field.

WEEDS LISTED

AGERATUM	<i>Ageratum conyzoides</i>
AMARANTH SPINY	<i>Amaranthus spinosus</i>
BALSAMAPPLE	<i>Momorcica charantia</i>
BARNYARD GRASS (WATERGRASS)+	<i>Echinonola crus-galli</i>
BEDSTRAW CATCHWEED	<i>Galium aparine</i>
BITTERCRESS/LESSER	<i>Cardamine oligosperma</i>
BLUEGRASS (ANNUAL)+	<i>Poa annua</i>
BUCKWHEAT WILD	<i>Polygonum convolvulus</i>
BURCLOVER	<i>Medicago hispida</i>
BUTTERCUP SMALLFLOWER	<i>Ranunculus abortivus</i>
BUTTONWEED	<i>Borreria laevis</i>
CAMPHORWEED	<i>Heterotheca subaxillaris</i>
CANARYGRASS/ANNUAL	<i>Phalaris canariensis</i>
CARPETWEED	<i>Mollugo verticillate</i>
CHEESEWEED (MALVA)	<i>Malva parviflora</i>
CLOVER, RED+	<i>Trifolium pratense</i>
CLOVER, WHITE+	<i>Trifolium repens</i>
COCKLEBUR, COMMON	<i>Xanthium pensylvanicum</i>
CRABGRASS, LARGE (HAIRY)	<i>Digitaria sanguinalis</i>
CROTALARIA	<i>Crotalaria species</i>
CROTON, TROPIC	<i>Croton glandulosus</i>
CUDWEED, NARROWLEAF	<i>Gnaphalium falcatum</i>
EVENINGPRIMROSE, CUTLEAF	<i>Oenothera laciniata</i>
FIDDLENECK, COAST+	<i>Amsinckia intermedia</i>
FILAREE, BROADLEAF	<i>Erodium botrys</i>
FILAREE, REDSTEM	<i>Erodium cicutaruim</i>
FILAREE, WHITESTEM	<i>Erodium moschatum</i>
FIREWEED (FROM SEED)	<i>Epilobium angustifolium</i>
FLIXWEED	<i>Descurainia sophia</i>
FOXTAIL, GIANT+	<i>Setaria faberi</i>
FOXTAIL, GREEN	<i>Setaria viridis</i>
FOXTAIL, YELLOW	<i>Setaria lutescens</i>
GERANIUM, CAROLINA	<i>Geranium carolinianum</i>
GOOSEGRASS+	<i>Eleusine ndica</i>
GROUNDCHERRY, CUTLEAF	<i>Physalis angulata</i>
GROUNDCHERRY, WRIGHT	<i>Physalis wrightii</i>
GROUNDSEED, COMMON	<i>Senecio vulgaris</i>
HENBIT	<i>Lamium amplexicaule</i>
HORSEWEED (MARESTAIL)	<i>Conyza canadensis</i>
JIMSONWEED	<i>Datura stramonium</i>
JOHNSONGRASS, SEEDLING	<i>Sorghum halepense</i>
KNOTWEED, PROSTRATE	<i>Polygonum aviculare</i>
LADYSTHUMB (SMARTWEED)	<i>Ploygonum persicaria</i>
LAMBSQUARTERS, COMMON	<i>Chenopodium album</i>
LETTUCE, PRICKLY (CHINA LETTUCE)	<i>Lactuca serriola</i>
MALLOW, LITTLE (MALVA)	<i>Malva parviflora</i>
MAYWEED (DOG FENNEL)	<i>Anthemis cotula</i>
MINERSLETTUCE	<i>Montia perfoliata</i>
MORNINGGLORY SPECIES, ANNUAL	<i>Ipomoea species</i>
MORNINGGLORY, IVYLEAF+	<i>Ipomoea hederacea</i>

MORNINGGLORY, TALL+	<i>Ipomoea purpurea</i>
MUSTARD, BLACK	<i>Brassica nigra</i>
MUSTARD, BLUE (PURPLE MUSTARD)	<i>Chorispora tenella</i>
MUSTARD, COMMON YELLOW	<i>Brassica campestris</i>
MUSTARD, HEDGE	<i>Sisymbrium officinale</i>
MUSTARD, TUMBLE (JIM HILL MUSTARD)	<i>Sisymbrium altissimum</i>
MUSTARD, WILD	<i>Brassica kaber</i>
NETTLE, BURNING	<i>Urtica urens</i>
NIGHTSHADE, AMERICAN BLACK	<i>Solanum americanum</i>
NIGHTSHADE, BLACK	<i>Solanum nigrum</i>
NIGHTSHADE, HAIRY	<i>Solanum sarrachoides</i>
OATS, WILD	<i>Avena fatua</i>
ORACH, RED	<i>Atriplex rosea</i>
OXALIS (BERMUDA BUTTERCUP)	<i>Oxalis pes-caprae</i>
PANICUM, FALL	<i>Panicum dichotomiflorum</i>
PEPPERWEED, VIRGINIA	<i>Lepidium virginicum</i>
PEPPERWEED, YELLOWFLOWER	<i>Lepidium perfoliatum</i>
PIGWEED, PROSTRATE	<i>Amaranthus biltoides</i>
PIGWEED, REDROOT	<i>Amaranthus retroflexus</i>
PIMPERNEL, SCARLET	<i>Anagallis arvensis</i>
POINSETTIA, WILD	<i>Euphorbia heterophylla</i>
PUNCTUREVINE	<i>Tribulus terrestris</i>
PURSLANE, COMMON	<i>Portulaca oleracea</i>
PUSLEY, FLORIDA	<i>Richardia scarbra</i>
RAGWEED, COMMON	<i>Ambrosia artemisiifolia</i>
REDMAIDS	<i>Calandrinia caulescens</i>
ROCKET, LONDON	<i>Sisymbrium irio</i>
RYEGRASS, ITALIAN	<i>Lolium multiflorum</i>
SAGE, LANCELEAF	<i>Salvia reflexa</i>
SANDBUR, FIELD	<i>Cenchrus incertus</i>
SANDSPURRY, RED	<i>Spergularia rubra</i>
SESBANIA, HEMP	<i>Sesbania exaltata</i>
SHEPHERDSPURSE+	<i>Capsella bursa-pastoris</i>
SICKLEPOD	<i>Cassia obtusifolia</i>
SIDA, PRICKLY (TEAWEED)	<i>Sida spinosa</i>
SIGNALGRASS, BROADLEAF	<i>Brachiaria platyphylla</i>
SMARTWEED, PENNSYLVANIA	<i>Polygonum pensylvanicum</i>
SORREL, RED (FROM SEED)	<i>Rumex acetosella</i>
SOWTHISTLE, ANNUAL	<i>Sonchus oleraceus</i>
SPEEDWELL, BIRDSEYE	<i>Veronica persica</i>
SPURGE, GARDEN	<i>Euphorbia hirta</i>
SPURGE, PROSTRATE++	<i>Euphorbia supina</i>
SPURGE, SPOTTED++	<i>Euphorbia maculata</i>
SPURRY, CORN	<i>Spergula arvensis</i>
TANSYMUSTARD	<i>Descurainia pinnata</i>
THISTLE, BULL++	<i>Cirsium vulgare</i>
THISTLE, RUSSIAN	<i>Salsola kali</i>
VELVETLEAF	<i>Abutilon theophrasti</i>
WITCHGRASS	<i>Panicum capillare</i>
WITCHWEED	<i>Striga asiatica</i>
WOODSORREL, COMMON YELLOW	<i>Oxalis stricta</i>

+ Highest rate and/or multiple applications may be required for acceptable control.

++Preemergence control only.

APPLICATION METHODS AND RECOMMENDED CULTURAL PRACTICES

PREEMERGENCE WEED CONTROL

Apply the recommended rate in a broadcast spray volume of 15 or more gallons of water per acre using calibrated spray equipment capable of uniform application to the soil surface. Seeding weeds are controlled as they come in contact with the soil-applied herbicide during emergence. Preemergence weed control is most effective when OXYFLUORFEN 2EC is applied to soil surfaces that are clean (free of crop or weed residues or clippings) and weed-free. Prior to application, weed or crop residues should be removed by thorough incorporation into the soil using tillage equipment or by blowing the area to be treated. At least 0.25 inch of irrigation or rainfall is required to activate OXYFLUORFEN 2EC and should occur within 3 to 4 weeks after application. For optimum results, OXYFLUORFEN 2EC should be applied to prepared beds or soil surfaces that will be left undisturbed during the time period for which weed control is desired. Cultural practices that disturb or redistribute surface soil following treatment with OXYFLUORFEN 2EC such as cutting water furrows will reduce weed control effectiveness.

Application Rates and Rate Ranges: Where rate ranges are given, use the lower rate in the rate range on coarse texture soils with less than 1% organic matter and lighter weed infestations. Use higher rates in the rate range on medium to fine texture soils, soils containing greater than 1% organic matter, heavy weed infestations, or for extended residual preemergence weed control.

POSTEMERGENCE WEED CONTROL

Apply the specified rate in a broadcast spray volume of 20 or more gallons of water per acre (a minimum 10 gallons if applying OXYFLUORFEN 2EC in tank mix with glyphosate). Because OXYFLUORFEN 2EC is a contact herbicide, complete and uniform coverage of weed foliage is essential for optimum postemergence control. Increase the spray volume to ensure complete and uniform coverage as weed height and density increases or in the presence of heavy trash (weed or crop residue). Postemergence applications of OXYFLUORFEN 2EC are most effective when made to weeds at the seedling stage. Applications made later than the 4-inch or 4 leaf stage may result in partial control or suppression. Postemergence applications should be made to seedling grasses not exceeding the 2-leaf stage. The addition of 0.25% v/v (2 pints per 100 gallons of spray) of an 80% active nonionic surfactant, labeled for application to growing crops, will enhance herbicidal effectiveness in controlling emerged weeds.

Postemergence Application Rates: Where a rate range is given, use a higher rate in the rate range for heavy weed infestations, weeds in advanced stages of growth or for extended residual preemergence weed control following control of existing emerged weeds.

GROUND APPLICATION

Ground Broadcast: Apply OXYFLUORFEN 2EC using conventional low-pressure ground spray equipment with flat fan nozzles. Follow manufacturer's recommendations for spraying pressure and boom height. An off-center (OC) nozzle positioned at the end of the boom may be desired. Check calibration of spray equipment before each use.

Directed Sprays: Apply OXYFLUORFEN 2EC as a coarse low-pressure spray in a spray volume of 20 or more gallons of spray per acre (broadcast basis). Follow manufacturer's recommendations for nozzle spacing and operating pressure. Spray should be directed toward the soil at the base of the crop. In row crops, use a minimum of 2 flat fan nozzles per row (one on each side) and for optimum spray coverage use 4 flat fan nozzles per row (two on each side). The 2 forward nozzles should point forward and downward while the rear nozzles should point to the rear and downward. With either sprayer system, nozzles should be adjusted to cover the weed foliage but minimize contact with the crop. Do not apply with hollow cone nozzles.

IMPORTANT: OXYFLUORFEN 2EC is a contact herbicide. Contact of sprays or drift with foliage or green stems can cause severe crop injury. Use directed sprays and spray shields and/or leaf lifters as necessary to minimize

contact of spray or drift with crop foliage or stems. Young green stems of woody plants are also susceptible to injury from spray contact. Potential for injury to woody stems diminishes with loss of green color and the development of relatively impervious non-living corky tissue (bark) on the surface of the stem.

Band Application: Application rates listed in this label are for broadcast application. For band application the rate per broadcast acre should be reduced according to the following formula:

$$\begin{array}{l} \text{Band Width (in inches)} \\ \text{Row Width (in inches)} \end{array} \times \begin{array}{l} \text{Rate per} \\ \text{Broadcast Acre} \end{array} = \begin{array}{l} \text{Amount Needed per Acre} \\ \text{for Banded Application} \end{array}$$

SPOT APPLICATION

For spot application, apply sprays uniformly to soil for preemergence weed control or on a spray-to-wet basis for postemergence weed control. Mix the required amount of OXYFLUORFEN 2EC with the recommended amount of water. For preemergence weed control, use one-half to one gallon of spray per 1000 sq ft. for postemergence weed control use a minimum of 1 gallon of spray per 1000 sq ft and add an 80% nonionic surfactant at the rate of 0.5 fl oz (1 Tbs) per gallon of spray. If making spot applications within an established crop, use coarse low-pressure sprays and direct the spray to the soil beneath the plants. To avoid crop injury, do not allow sprays to contact leaves and stems of herbaceous plants or leaves or green stems of woody plants.

Amount of OXYFLUORFEN 2EC Required to Treat 1000 sq ft at Specified Application Rate					
0.5 pt/acre	1.0 pt/acre	2.0 pt/acre	3.0 pt/acre	4.0 pt/acre	8.0 pt/acre
0.2 fl oz (5.5 ml)	0.4 fl oz (11 ml)	0.75 fl oz (22 ml)	1.1 fl oz (33 ml)	1.5 fl oz (44 ml)	3.0 fl oz (88 ml)

1 pint = 16 fl oz; 1 fl oz = 29.6 (30) ml

AERIAL APPLICATION

Use aerial boom equipment designed for use with herbicides and a minimum spray volume of 10 gallons per acre (5 gallons per acre if tank mixed with glyphosate). Do not aerially apply OXYFLUORFEN 2EC unless crop-specific use directions specifically allow and provide directions for aerial application.

AVOID DRIFT: Exercise extreme care to void herbicide contact with any desirable dormant or non-dormant crop, plant, tree or vegetation as severe injury may result. Extreme care must be exercised to prevent spray drift that could result in damage to other crops or desirable vegetation. Adhere to the following guidelines when aerial applications are to be made.

Spray Drift Management (Aerial Application): Avoiding spray drift at the application site is the responsibility of the applicator. The potential for spray drift is controlled by the interaction of many equipment-and-weather-related factors. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan of rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator must adhere to the following requirements when OXYFLUORFEN 2EC is aerially applied:

1. Do not apply when the wind direction is not stable, when inversion conditions exist, or when wind velocity exceeds 10 mph.

2. When wind speeds are 5 mph or less, maintain a minimum downwind buffer zone of at least ½ mile from all crops and desirable vegetation, except the following:
 - 150 feet from dormant treefruit/nut/vine crops and overwintering sugar beets.
 - 650 feet from garlic, jojoba, legumes, onions, pastures, small grains, seedling sugar beets, and non-targeted vegetable fallow beds.
3. When wind speeds are between 5 and 10 mph, downwind buffer zones in excess of those listed above are suggested.
4. For upwind and side borders, maintain buffer zone of 150 feet from any non-targeted vegetable fallow bed, crop, or desirable vegetation.

The use of drift control agent may be required by local regulations. However, the drift control agent may decrease the weed control effectiveness.

Important: Aerial applicators must be familiar with the label for OXYFLUORFEN 2EC and follow all applicable use precautions. Applying OXYFLUORFEN 2EC in a manner other than recommended in this label is done at the user's risk. Users are responsible for all loss or damage resulting from aerial spraying. In addition, aerial applicators should follow all applicable state and local regulations and ordinances. In interpreting the label and local regulations, the most restrictive limitations apply.

CHEMIGATION

Do not apply this product through any irrigation system unless the instructions for chemigation are followed. If application by chemigation is not specifically listed under the specific crop use instructions, OXYFLUORFEN 2EC herbicide may not be applied to that crop through an irrigation system.

Apply this product only through sprinkler (center pivot, solid set, portable lateral, or low-volume (micro sprinkler), drip (trickle), or flood (basin) irrigation systems. Refer to use directions for specific crops for instructions as to which type of irrigation system may be used. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SPRINKLER CHEMIGATION (FOLIAR SPRAY USES)

For sprinkler irrigation, sufficient water should be applied at the beginning of the irrigation period to ensure uniform wetting of the plant and/or soil surfaces. Meter OXYFLUORFEN 2EC at a continuous uniform rate during the middle 1/3 of the irrigation period to allow for uniform distribution to the vegetation and/or soil surface. Continue irrigation during the final 1/3 of the irrigation period to ensure proper flushing of the irrigation system. During sprinkler irrigation, sufficient water should be applied to ensure water penetration to a depth of two inches.

AVOID DRIFT: Extreme care must be exercised to prevent spray drift that could result in damage to other crops or desirable vegetation. Use the following guidelines when applications of OXYFLUORFEN 2EC are made through sprinkler irrigation:

- Do not apply when the wind direction is not stable, when inversion conditions exist, or when wind velocity exceeds 10 mph.
- When wind speeds are 5 mph or less, maintain a minimum download buffer zone of at least ½ mile from all

crops and desirable vegetation, except for the following:

- Maintain a minimum download buffer of:
 - 150 feet from dormant treefruit, dormant vines and overwintering sugar beets.
 - 650 feet from garlic, jojoba, legumes, onions, pastures, small grains, seedling sugar beets and vegetable fallow beds.
 - When wind speeds are between 5 and 10 mph, downwind buffer zones in excess of those listed above are suggested.
 - For upwind and side borders, maintain a minimum buffer zone of 150 feet from any vegetable fallow bed, crop, or desirable vegetation.

To apply a pesticide using sprinkler chemigation, the chemigation system must meet the following specifications:

- The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

FLOOD (BASIN) CHEMIGATION (SOIL DRENCH USES)

OXYFLUORFEN 2EC should be continuously metered into the water during the entire irrigation period. Agitation in the pesticide supply tank is suggested. Best weed control results from OXYFLUORFEN 2EC applied through flood (basin) irrigation systems are obtained when a uniform distribution and flow of irrigation water is maintained over level land.

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

DRIP (TRICKLE) CHEMIGATION (SOIL DRENCH USES)

Meter OXYFLUORFEN 2EC at a continuous uniform rate during the middle 1/3 of the irrigation period to allow for uniform distribution to the soil surface. For best results, OXYFLUORFEN 2EC should be uniformly positioned across the wetted area to help reduce the "ring effect" of weed escapes, as other products begin to break down around the emitter. Continue irrigation during the final 1/3 of the irrigation period to ensure proper flushing of the irrigation system.

To apply a pesticide using drip (trickle) chemigation, the chemigation system must meet the following specifications:

- The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pipe and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or (water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

CHEMIGATION CALIBRATION: FOR LOW-VOLUME SPRINKLERS (MICROSPRINKLERS) AND DRIP (TRICKLE) IRRIGATION SYSTEMS

Calculation of use rate is based on wetted area around emitters - NOT on grove acres. To determine correct amount of OXYFLUORFEN 2EC, use the following formula:

1. Treated area per each emitter = A
 $A = 3.14 \times (\text{radius} \times \text{radius})$

Example: If the average distance from emitter to perimeter of wetted area measured at the soil surface is 13 inches, then
 $A = 3.14 \times (13" \times 13")$
 $A = 3.14 \times (169)$
 $A = 530.7$ square inches

2. The area in square feet wet in each acre = B
 $B = \frac{A \times \text{emitters/acre}}{144}$

Example: If there are 300 emitters per acre, then
 $B = \frac{530.7 \times 300}{144} = B = 1105.6$ square feet wetted per acre

3. The total area (in square feet) wet by your system = C
 $C = B \times \text{acres covered by system}$

Example: If the system covers 20 acres, then
 $C = 1105.6$ square feet per acre \times 20 acres
 $C = 22,112$ square feet wetted by system

4. Amount of OXYFLUORFEN 2EC to inject = S

Rate per treated acre of OXYFLUORFEN 2EC = R

$$S = \frac{C \times R}{43,560} = \text{quarts of OXYFLUORFEN 2EC}$$

Example: If the desired application rate per treated acre is 1 quart of OXYFLUORFEN 2EC, then

$$S = \frac{22,112 \times 1.0}{43,560} = S = 0.507 \text{ quarts of OXYFLUORFEN 2EC should be injected into system.}$$

NOTE: Select the proper rate based on weed spectrum and desired length of control (See RATE RANGES section below).

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

If the chemigation system is connected to a public water supply, the following conditions must also be met:

- Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from a point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

CULTURAL CONSIDERATIONS

In order for OXYFLUORFEN 2EC to provide maximum preemergence activity:

Prior to application, the bed or soil surface should be smooth and free of crop and weed trash (decaying leaves, clippings, dead weeds, etc.). Leaves and trash may be removed by blowing the area to be treated or by thoroughly mixing the trash into the soil through cultivation prior to herbicide application. After application, at least one-quarter inch (1/4 inch) of irrigation or rainfall should occur within 3 or 4 weeks after application. The best results from OXYFLUORFEN 2EC are from applications to established beds or soil surfaces that are left undisturbed during the time period for which weed control is desired. Cultural practices that result in redistribution or disturbance of the soil surface after treatment will decrease the herbicidal effectiveness of OXYFLUORFEN 2EC. Cutting water furrows or cultivations that mix untreated soil into treated areas will also reduce the effectiveness of the treatment.

RATE RANGES

Select proper application rates based on soil conditions, weed spectrum and desired period of residual weed control.

Preemergence Application: Where rate ranges are given, use the lower rate in the rate range on coarse texture soils with less than 1% organic matter. Use higher rates in the rate range on medium to fine texture soils, soils containing greater than 1% organic matter, or where a longer period of residual weed control is desired.

Postemergence Application: Where a rate range is given, use higher rate in rate range for heavy weed infestations, weeds in advanced stages of growth or where a longer period of residual weed control is desired.

MIXING DIRECTIONS

Shake Well Before Using. Fill the spray tank at least one-third full of clean water. With the pump and agitator running, add the recommended amount of herbicides to the spray tank. The order of addition to the spray tank should be (1) wettable powders, (2) flowables and (3) soluble liquids. Complete filling of the spray tank with water.

Use of Surfactants: For all applications of OXYFLUORFEN 2EC (except garlic and onions) where postemergence weed control is desired, add 2 to 4 pints of Latron AG-98 nonionic surfactant (or comparable 80% active nonionic surfactant cleared for application to growing crops) per each 100 gallons of spray. The addition of 4 pints of Latron AG-98 per 100 gallons of spray is recommended to enhance postemergence activity when hard water (greater than 600 ppm) is used as carrier. Maintain agitation until spraying is completed.

Tank Mixing Precautions:

- Read and carefully follow all applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.
- Do not exceed specified application rates. Do not tank mix with another pesticide product that contains the same active ingredient as the product unless the label of either tank mix partner specifies the maximum dosages that may be used.

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of this product and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour, if the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Sprayer Cleanup: Thoroughly flush spray equipment (tank, pump, hoses and boom) with clean water before and after each use. Residues of OXYFLUORFEN 2EC remaining in spray equipment may damage other crops. The addition of a non-ionic surfactant to equipment flushing waters at the rate of 1 quart per 100 gallons is recommended to aid in removal of residues of OXYFLUORFEN 2EC.

CROP-SPECIFIC USE INFORMATION

ARTICHOKE (GLOBE)

Post-Directed Spray Application

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	4 – 6	<p>Application Method: Apply as a directed spray to the soil surface between the rows and at the base of artichoke plants in a minimum spray volume of 40 gallons per acre.</p> <p>Timing to Crop: Apply after completion of ditching operations. Separate applications of up to 4 pt/acre may be made 8 to 10 weeks apart or a single application of up to 6 pt/acre may be made.</p> <p>Timing to Weeds: Preemergence up to 8 leaf stage.</p>
<p>Precautions:</p> <p>Do not apply over-the-top. Contact with direct spray or drift will cause injury to artichoke fronds or severe injury to buds or flowers.</p> <ul style="list-style-type: none"> • Application of OXYFLUORFEN 2EC to artichoke plantings should be delayed a minimum of 60 days after cutting back or transplanting. 		
<p>Restrictions:</p> <ul style="list-style-type: none"> • Do not apply more than 6 pints of OXYFLUORFEN 2EC per acre per season as a result of a single application or multiple applications. • Preharvest Interval: Do not apply within 5 days of harvest. 		

Key Weeds Controlled:

Preemergence	Postemergence
cheeseweed (malva) groundsel, common lambsquarter, common mustard, common yellow oxalis (Bermuda buttercup) [†] shepherdspurse sowthistle, annual	cheeseweed (malva) groundsel, common mustard, common yellow nettle, burning oxalis (Bermuda buttercup) [†] shepherdspurse sowthistle, annual

[†] Suppression

BROCCOLI / CABBAGE / CAULIFLOWER

Pre-Transplant (Preplant) Application for Preemergence Broadleaf Weed Control

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	1 – 2	Pre-Transplant Application Only: Apply broadcast to final seedbed prior to transplanting. Use lower rate in the rate range on coarse textured soils with less than 1% organic matter. Use the highest rate in the rate range on medium to fine textured soils or soils containing greater than 1% organic matter. Transplanting should be accomplished with minimal soil disturbance and soil left undisturbed during the time weed control is desired.

Precautions:

- Pre-transplant applications may result in initial, but temporary, crop injury (leaf cupping or crinkling) and is enhanced if crop leaves come in direct contact with treated soil. Crop will rapidly outgrow this condition and develop normally. Severe crop injury may result if transplants are under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides or storage conditions. The use of transplants less than 5 weeks old or use of extremely succulent transplants grown in containers less than 1 inch square, may increase the severity of crop injury. Hardening off, increasing the age of transplants or increasing the size of the rooting containers will lessen the possibility and/or severity of potential crop injury.
- OXYFLUORFEN 2EC will assist in early season annual grass control, however, a herbicide program for preemergence or postemergence control of annual grasses is recommended.
- Note: Do not apply OXYFLUORFEN 2EC if an acetanilide herbicide such as Dual Magnum herbicide, Lasso herbicide, or Ramrod herbicide has been applied to the field during the current growing season as severe crop injury may occur.
- Do not apply OXYFLUORFEN 2EC as a preemergence treatment to direct-seeded broccoli, cabbage or cauliflower.
- Do not apply OXYFLUORFEN 2EC post-transplant or over-the-top of broccoli, cabbage or cauliflower.
- Applications to muck soils may result in partial weed control or suppression.
- Furrow and drip irrigation immediately after transplanting and under high temperatures can result in increased crop injury. Sprinkler irrigation is recommended during early establishment of transplants. If these conditions cannot be met, OXYFLUORFEN 2EC herbicide should not be used.

Crop-Specific Restrictions:

- Do not apply more than 2 pints of OXYFLUORFEN 2EC per treated acre per season.

Key Weeds Controlled:

Preemergence
carpetweed pigweed, redroot purslane, common smartweed, Pennsylvania

CACAO (BEARING AND NONBEARING)
(For Use Only in Hawaii)

OXYFLUORFEN 2EC may be applied as a pre-transplant treatment or to established or recently transplanted cacao.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	2 – 8	Pre-transplant Application: Up to 4 pints per broadcast acre may be applied as a pre-transplant application. Application to Established Plantings: In established plantings, including recently transplanted cacao plants, apply as a directed spray to the orchard floor. Use higher rates in rate range and increase spray volume to control dense growth of existing weeds or for extended residual preemergence weed control.
Precautions: <ul style="list-style-type: none"> • Do not apply preplant or preemergence to direct-seeded cacao. • OXYFLUORFEN 2EC should be applied to only healthy growing trees/transplants of suitable size to allow directed sprays. Avoid spray contact with foliage. 		
Crop-Specific Restrictions: <ul style="list-style-type: none"> • Do not apply more than 8 pints of OXYFLUORFEN 2EC per acre as a single application or more than 24 pints per acre per year. • Preharvest Interval: Do not apply OXYFLUORFEN 2EC within 1 day of harvest. 		

Key Weeds Controlled:

Preemergence	Postemergence
ageratum buttonweed crotalaria purslane, common spurge, garden	purslane, common spurge, garden

CITRUS (NONBEARING)

Citrus, such as Calamondin, Chironja, Citrus Citron, Grapefruit, Kumquat, Lemon, Lime, Mandarin, Pummelo, Satsuma Mandarin, Sour Orange, Sweet Orange, Tangelo, Tangerine, Tangor

OXYFLUORFEN 2EC may be applied only in non-bearing citrus orchards. Apply only as a directed spray to the orchard floor avoiding contact with citrus foliage.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	6	Preemergence Weed Control: Up to 6 pt/acre may be applied for residual preemergence weed control.
Postemergence	2 – 6	Postemergence Weed Control: The 6 pint/acre rate will control weeds up to 4 inches tall. Weeds greater than 4-inch or 4 inches tall may be partially controlled. Use sufficient spray volume for complete and uniform coverage of weeds. Increase the spray volume with increased weed height and density to ensure complete coverage.
<p>Tank Mixing: Refer to Mixing Directions section for Tank Mixing Precautions.</p> <ul style="list-style-type: none"> • Preemergence Use: For residual control of grass weeds, OXYFLUORFEN 2EC may be tank mixed with grass herbicides labeled for use in citrus. • Postemergence Use: For broader spectrum postemergence control of emerged grass and broadleaf weeds, OXYFLUORFEN 2EC may be tank mixed with paraquat (Gramoxone herbicide) or glyphosate. 		
<p>Precautions:</p> <ul style="list-style-type: none"> • Do not apply during periods of new citrus foliage growth. Applications should be made after foliage has fully expanded and hardened off. Avoid direct spray contact with citrus foliage. 		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> • Apply OXYFLUORFEN 2EC only to nonbearing citrus (trees that will not bear fruit for one year). • Do not apply more than 6 pints of OXYFLUORFEN 2EC per acre per year as a result of a single or multiple applications. 		

Key Weeds Controlled:

(Arizona and California)		(Florida, Louisiana and Texas)	
Preemergence	Postemergence	Preemergence	Postemergence
burclover	cheeseweed (malva)	cutweed, narrowleaf	balsamapple
cheeseweed (malva)	fiddleneck, coast	eveningprimrose, cutleaf ^{††}	cudweed, narrowleaf ^{†††}
fiddleneck, coast	filaree, broadleaf [†]	groundcherry, cutleaf	eveningprimrose, cutleaf ^{††}
filaree, broadleaf	filaree, redstem [†]	lambsquarters, common	groundcherry, cutleaf
filaree, redstem	filaree, whitestem [†]	nightshade, American black	groundcherry, Wright
filaree, whitestem	groundsel, common	nightshade, black	lambsquarters, common
groundsel, common	henbit	pepperweed, Virginia	morningglory, annual
henbit	minerslettuce	pigweed, redroot	nightshade, American black
knotweed, prostrate	nettle, burning	poinsettia, wild	nightshade, black
lambsquarters, common	pigweed, redroot	pusley, florida	pepperweed, Virginia
lettuce, prickly	redmaids	sida, prickly (teaweed)	pigweed, redroot
pigweed, redroot	shepherdspurse	smartweed, Pennsylvania	poinsettia, wild
purslane, common	sowthistle, annual	sowthistle, annual	purslane, common
redmaids		spurge, prostrate	pusley, florida
rocket, London		spurge, spotted	sida, prickly (teaweed)
shepherdspurse			smartweed, Pennsylvania
sowthistle, annual			sowthistle, annual
spurge, prostrate			
spurge, spotted			

† OXYFLUORFEN 2EC at the 6 pt/acre will provide control of filaree and other weeds up to 4-inch stage. Applications to weeds beyond the 4-inch stage may result in partial control.
 †† Highest rate and/or multiple applications may be required for acceptable control.
 ††† Maximum 0.5-inch diameter.

CLARY SAGE

Clary Sage (*Salvia sclarea*) Grown and Utilized in the Essence Industry
 (For Use Only in North Carolina)

Weed Control	Rate (pt/acre)	Specific Use Directions
Postemergence	0.5 - 1	<p>OXYFLUORFEN 2EC may be applied to established clary sage for control of henbit (<i>Lamium amplexicaule</i>) and other winter annual broadleaf weeds during the winter and spring season.</p> <p>Apply shortly after the first flush of henbit is in the 2- to 4-leaf stage of growth. Additional applications may be required to control subsequent weed flushes through the spring season. After treatment, henbit will stop growing and slowly die. Increase the spray volume if weed growth is dense.</p>
<p>Precautions:</p> <ul style="list-style-type: none"> Clary sage may respond to the topical application of this product with some marginal leaf burn, but recovery is rapid. 		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> Do not apply more than 6 pints per acre per year. 		

COFFEE (BEARING AND NONBEARING)
(For Use Only in Hawaii)

OXYFLUORFEN 2EC may be applied to established coffee, recently transplanted coffee, or as a pre-transplant treatment. In established non-dormant coffee, apply as a directed spray avoiding contact with crop foliage. Newly established transplants should be healthy and well established and of sufficient size to allow use of directed sprays without contacting crop foliage.

OXYFLUORFEN 2EC may be applied over-the-top of dormant coffee transplants. Transplants are considered to be dormant when active terminal growth has ceased and terminal buds have formed. Application over-the-top of coffee plants after buds start to swell (a sign that new growth has resumed) may result in crop injury and is not recommended.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	2 – 8	<p>Preemergence Weed Control:</p> <ul style="list-style-type: none"> • Apply as a directed spray to the orchard floor beneath established coffee plants. • Up to 4 pints per acre may be applied as a pre-transplant application prior to transplanting coffee plants. <p>Postemergence Weed Control:</p> <ul style="list-style-type: none"> • Increase the spray volume when weed growth is dense or trash is present; or use a higher rate within the rate range for extended residual preemergence weed control.
<p>Tank Mixing: Refer to Mixing Directions section for Tank Mixing Precautions. Apply tank mixes only as a directed sprays.</p>		
<p>Precaution:</p> <ul style="list-style-type: none"> • To prevent foliar injury, do not apply during periods of rapid new growth or allow spray or drift to contact actively growing foliage. 		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> • Do not apply preplant or preemergence to direct-seeded coffee. • Do not apply more than 8 pints per broadcast acre of OXYFLUORFEN 2EC in a single application or 24 pints per broadcast acre per year. • Preharvest Interval: Do not apply OXYFLUORFEN 2EC within one (1) day of harvest. 		

Key Weeds Controlled:

Preemergence	Postemergence
ageratum buttonweed crotalaria purslane, common spurge, garden	purslane, common spurge, garden

CONIFER SEEDBEDS. TRANSPLANTS. CONTAINER STOCK AND
SELECTED FIELD GROWN DECIDUOUS TREES

Use Precautions and Restrictions:

- Not for conifer release in forest management programs or for forest regeneration applications.
- Do not apply OXYFLUORFEN 2EC in an enclosed greenhouse structure as injury to plant foliage may result.
- Do not store or transport treated container stock in an enclosed structure until completion of 4 irrigations (minimum 21 days) as injury to non-labeled plants may occur.
- Apply OXYFLUORFEN 2EC only to healthy conifer stock. Do not apply OXYFLUORFEN 2EC to conifers that are under stress from excessive fertilizer or soil salts, disease, nematodes, frost, drought, flooding, previously applied pesticides, soil insects, or winter injury, as severe injury may result.
- Do not graze or harvest livestock forage from treated areas.

Key Weeds Controlled: When OXYFLUORFEN 2EC is applied preemergence or postemergence at recommended dosages and weed stages.

barnyardgrass [†]	mustard, blue
bedstraw, catchweed	mustard, tumble
bittercress, lesser	mustard, wild
bluegrass, annual [†]	nettle, burning
buckwheat, wild	nightshade, black
burclover	nightshade, hairy
carpetweed	oats, wild
clover, red [†]	orach, red
clover, white [†]	pepperweed, yellowflower
cocklebur, common	pigweed, prostrate
crabgrass, large [†]	pigweed, redroot
fiddleneck, coast [†]	pimpernel, scarlet
filaree, broadleaf	purslane, common
filaree, redstem	redmaids
fireweed (from seed)	rocket, London
flixweed	sandspurry, red
foxtail, giant [†]	shepherdspurse [†]
goosegrass [†]	sida, prickly
groundcherry, cutleaf	smartweed, Pennsylvania
groundcherry, wright	sorrel, red (from seed)
groundsel, common	sowthistle, annual
henbit	speedwell, birdseye
jimsonweed	spurge, prostrate ^{††}
knotweed, prostrate	spurge, spotted ^{††}
ladysthumb	spurry, corn
lambsquarters, common	tansymustard
lettuce, prickly	thistle, bull ^{††}
mallow, little	thistle, Russian
mayweed	velvetleaf
minerslettuce	witchgrass
morningglory, ivyleaf [†]	woodsorrel, yellow ^{††}
morningglory, tall [†]	

[†] Highest rate and/or multiple applications may be required for acceptable control.

^{††} Preemergence control only.

CONIFER SEEDBEDS

Agricultural Use Requirements: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 3 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

Coveralls

- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

OXYFLUORFEN 2EC provides both postemergence and residual preemergence control of many broadleaf weeds and annual grass species.

Seeded conifers are tolerant to preemergence and postemergence applications of OXYFLUORFEN 2EC. For weed control during the establishment of conifer seedlings, OXYFLUORFEN 2EC can be applied after seeding of conifers, but prior to emergence. For weed control in emerged conifers, OXYFLUORFEN 2EC may be applied over-the-top, but application should be delayed a minimum of 5 weeks after seedling emergence. If application is made during cool, cloudy weather, make certain that seedlings have hardened-off prior to spraying.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	1 – 4	Application after planting, but prior to emergence of conifer seedlings: Where grass weeds are present, apply 2 to 4 pints of OXYFLUORFEN 2EC per acre. In known areas of high weed competition, apply 4 pints of OXYFLUORFEN 2EC per acre. Broadcast to beds and irrigate with 1/2 to 3/4 inch of sprinkler irrigation before weed emergence. OXYFLUORFEN 2EC is most effective on annual grasses when applied preemergence.
Postemergence	1 – 2	Application after emergence of conifer seedlings: Application should be made to seedling weeds less than 4 inches in height (seedling grasses not exceeding the 2-leaf stage). Depending of subsequent weed flushes, multiple applications may be necessary to achieve season-long weed control.
Chemigation: OXYFLUORFEN 2EC may be applied at labeled rates through sprinkler irrigation systems. For center pivot irrigation systems, apply the specified dosage of OXYFLUORFEN 2EC per acre metered at a continuous uniform rate during the entire irrigation period, otherwise meter OXYFLUORFEN 2EC at a continuous uniform rate during the middle 1/3 of the irrigation period. When applying by sprinkler irrigation, follow directions given in the Chemigation Instructions section of this label.		
Precautions:		
<ul style="list-style-type: none"> • Occasionally spotting, crinkling, or flecking may appear on leaves of conifers. Leaves that receive direct spray or drift may be injured, but typically outgrow this condition rapidly and develop normally. 		
Crop-Specific Restrictions:		
<ul style="list-style-type: none"> • Do not apply more than 8 pints of OXYFLUORFEN 2EC per acre per year. 		

OXYFLUORFEN 2EC may be applied to conifer seedbeds of the following species:

Important: When applied as directed, the conifer species listed on this label have shown tolerance to OXYFLUORFEN 2EC. It is impossible, however, to evaluate this product on all varieties, biotypes and cultivars of listed species under all possible growing conditions. Until familiar with results under local growing conditions, the user should exercise reasonable judgement and caution with this product. Limit application of this product to a few

plants in a small area to determine plant tolerance and extent of injury if such occurs, prior to initiating large-scale applications.

Douglas Fir	<i>Pseudotsuga menziesii</i>
Fir	Fraser (<i>Abies fraseri</i>) Grand (<i>Abies grandis</i>) Noble (<i>Abies procera</i>)
Hemlock	Eastern hemlock (<i>Tsuga canadensis</i>)
Pine	Austrian (<i>Pinus nigra</i>) Eastern White (<i>Pinus strobus</i>) Himalayan (<i>Pinus wallichiana</i>) Jack (<i>Pinus banksiana</i>) Loblolly (<i>Pinus taeda</i>) Lodgepole (<i>Pinus contorta</i>) Longleaf (<i>Pinus palustris</i>) Monterey (<i>Pinus radiata</i>) Mugho (<i>Pinus mugo</i>) Ponderosa (<i>Pinus ponderosa</i>) Scotch (<i>Pinus sylvestris</i>) Shortleaf (<i>Pinus echinata</i>) Slash (<i>Pinus elliotii</i>) Virginia (<i>Pinus virginiana</i>)
Spruce	Blue (<i>Picea pungens</i>) Dwarf (<i>Picea glauca Conica</i>) Alberta (<i>Picea abies</i>) Norway (<i>Picea sitchensis</i>)

CONIFER TRANSPLANTS AND CONTAINER STOCK
(INCLUDES 2-0 SEEDLING AND CHRISTMAS TREE PLANTINGS)

Agricultural Use Requirements: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 6 days.
 PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Many container-grown conifers and conifer transplants are tolerant to preemergence and postemergence applications of OXYFLUORFEN 2EC. Applied postemergence, OXYFLUORFEN 2EC provides postemergence control of emerged weeds and preemergence residual control of many broadleaf weeds and grasses (see Key Weeds Controlled) at the beginning of this section.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	4 – 8	Transplanted and Container Grown Conifers: For best results, preemergence applications should be made immediately after transplanting seedlings or to weed-free container stock. Postemergence applications should be made to weeds less than 4 inches in height. Two applications may be necessary, in fall-transplanted conifer fields, for season-long weed control. The addition of a non-ionic surfactant (0.25% v/v) labeled for application to growing crops, enhances the activity of OXYFLUORFEN 2EC on emerged weeds.
Precautions: <ul style="list-style-type: none"> • Do not make over-the-top applications during periods of active conifer growth. Apply only before bud break or after new terminal growth has hardened off. 		
Crop-Specific Restrictions: <ul style="list-style-type: none"> • Do not apply more than 8 pints of OXYFLUORFEN 2EC per acre in a single application or more than 16 pints per acre per year. 		

In addition to those conifer species listed under the Conifer Seedbed section, the following conifer species have been shown to be tolerant to OXYFLUORFEN 2EC:

Arborvitae	<i>Thuja occidentalis</i> <i>Thuja orientalis</i>
Juniper	<i>Juniperus chinensis</i> <i>Juniperus horizontalis</i> <i>Juniperus procumbens</i> <i>Juniperus sabina</i> <i>Juniperus scopulorum</i>
Red cedar	<i>Juniperus virginiana</i>
Western Hemlock	<i>Tsuqa heterophylla</i>
Yew	<i>Taxus species</i>

SELECTED FIELD-GROWN DECIDUOUS TREES

Listed field-grown deciduous trees are tolerant only to directed spray applications of OXYFLUORFEN 2EC. OXYFLUORFEN 2EC provides both preemergence and postemergence control of listed broadleaf weeds and grasses.

Timing to Crop: Apply OXYFLUORFEN 2EC to established deciduous trees or after transplanting. For optimum weed control, applications should be made prior to weed germination. Apply only as a directed spray to soil beneath the trees.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Early postemergence	2 - 6	OXYFLUORFEN 2EC may be applied to established deciduous trees or after transplanting as a single or split application. Apply as a directed spray to the soil surface. Use of spray shields to reduce exposure of foliage and bark is recommended. The addition of a non-ionic surfactant (0.25% v/v) labeled for application to growing crops, will enhance herbicidal activity on emerged weeds. Spot Application: Spot treatments at recommended rates may be used to control localized weed infestations. See use directions for Spot Application in the Application Methods and Recommended Cultural Practices section.
Tank Mixing: For broader spectrum control, OXYFLUORFEN 2EC may be tank mixed with other preemergence or postemergence herbicides registered for this use in deciduous trees. Refer to Mixing Directions section for Tank Mixing Precautions.		
Precautions: <ul style="list-style-type: none"> For maximum crop safety, directed applications should be prior to budbreak in the spring or after trees have initiated dormancy in the fall. Avoid contact of spray or drift with foliage or stems with green bark. Application after bud swell may result in crop injury and is not recommended. If a non-dormant application is required due to weed competition, apply only after foliage has fully expanded and hardened off. Use only directed sprays and spray shields to prevent spray contact with stems with green bark or foliage. Do not apply OXYFLUORFEN 2EC to trees that have been weakened or are under stress from excessive fertilizer or soil salts, disease, nematodes, frost, wind injury, drought, flooding, previously applied pesticides, insects, or winter injury as severe injury may result. 		
Crop-Specific Restrictions: <ul style="list-style-type: none"> Do not apply more than 6 pints of OXYFLUORFEN 2EC per acre per year. Do not apply to bearing treefruit, nut and vine crops. For selected bearing treefruit, nut and vine crops, refer to Treefruit/Nut/Vine section of this label for use directions. Do not graze or feed livestock forage cut from areas treated with OXYFLUORFEN 2EC. 		

OXYFLUORFEN 2EC may be applied to the following deciduous tree species:

Almond ^{††}	<i>Prunus spp.</i>
Apple ^{††}	<i>Malus X domestica</i>
Apricot ^{††}	<i>Prunus spp.</i>
Ash, Green	<i>Fraxinus pennsylvanica</i>
Ash, White	<i>Fraxinus Americana</i>
Birch, River	<i>Betula nigra</i>
Cherry ^{††}	<i>Prunus spp.</i>
Chestnut ^{††}	<i>Castanea spp.</i>
Crabapple ^{††}	<i>Malus spp.</i>
Cottonwood	<i>Populus spp.</i>
Dogwood	<i>Cornus florida</i>
Eucalyptus	<i>Eucalyptus viminalis</i> <i>Eucalyptus pulverulenta</i> <i>Eucalyptus camaldulensis</i>
Filbert ^{††}	<i>Corylus spp.</i>
Lilac	<i>Syringa vulgaris</i>

Locust, Black	<i>Robinia pseudoacacia</i>
Maple, Black †	<i>Acer nigrum</i>
Maple, Red †	<i>Acer rubrum</i>
Maple, Sugar †	<i>Acer saccharum</i>
Myrtle, Crepe	<i>Lagerstroemia indica</i>
Nectarine ††	<i>Prunas spp.</i>
Nut, Hickory ††	<i>Carya spp.</i>
Nut, Macadamia	<i>Macadamia ternifolia</i>
Oak, Chestnut	<i>Quercus prinus.</i>
Oak, Cherrybark	<i>Quercus pagoda</i>
Oak, Nutt All	<i>Quercus nuttallii</i>
Oak, Pin	<i>Quercus palustres</i>
Oak, Red	<i>Quercus rubra</i>
Oak, Water	<i>Quercus nigra</i>
Oak, Willow	<i>Quercus phellos</i>
Olive, Russian	<i>Elaeagnus angustifolia</i>
Poplar	<i>Populus spp.</i>
Poplar, Tulip	<i>Liriodendron tulipifera</i>
Peach ††	<i>Prunas persica</i>
Pear ††	<i>Pyrus spp.</i>
Pecan ††	<i>Carya spp.</i>
Pistachio ††	<i>Pistacia vera</i>
Plum ††	<i>Prunas spp.</i>
Prune ††	<i>Prunas spp.</i>
Redbud	<i>Cercis canadensis</i>
Sweetgum	<i>Liquidambar styraciflua</i>
Sycamore	<i>Platanus occidentales</i>
Walnut, Black ††	<i>Juglans nigra</i>

† Do not apply to maple trees used for production of maple cap or maple syrup.

†† Apply only to nonbearing trees. For bearing treesfruit, nut and vine crops, refer to specific use directions in the Treefruit/Nut/Vine section of this label.

CORN

FOR USE ONLY ON FIELD CORN IN CONJUNCTION WITH THE USDA WITCHWEED ERADICATION PROGRAM IN NORTH CAROLINA AND SOUTH CAROLINA

Apply OXYFLUORFEN 2EC only as a directed spray from May through August for preemergence and postemergence control of witchweed (*Striga asiatica*). Corn must be a minimum of 24 inches tall. Examine witchweed infested fields during the early part of the growing season to determine uniformity of corn stand and grass weed pressure. If necessary, cultivate weed-infested fields prior to initial application of OXYFLUORFEN 2EC to allow for optimum soil coverage during the initial application. Fields treated with OXYFLUORFEN 2EC should be inspected regularly for any breakthrough of witchweed. If breakthrough occurs, a second application should be made as soon as possible after appearance of witchweed. Repeat treatments should occur prior to bloom stage to prevent seed set.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	2 – 3	Initial Application: Apply as a directed spray over the entire row surface at the rate of 2 pints per acre. Use up to 3 pints per acre in areas of heavy witchweed infestation. Use a minimum spray volume of 20 gallons per acre and a non-ionic surfactant at the time of 2 pints per 100 gallons of spray.
Postemergence	1 - 2	Repeat Applications: In case of witchweed breakthrough a repeat application may be made at 1 to 2 pints per acre.
Precautions: <ul style="list-style-type: none"> • Do not spray over the top of the corn, as this may result in severe corn injury. Spray should contact only the lower 3 to 8 inches of the corn stalk and any leaves in this zone. Spray droplets contacting the lower leaves will cause necrotic spotting or streaking of sprayed tissue. 		
Crop-Specific Restrictions: <ul style="list-style-type: none"> • Do not apply more than 5 pints (1.25 lb active) of OXYFLUORFEN 2EC per acre during the growing season. • Do not apply any application within 60 days of harvest. • Do not use corn plants from a treated field for green chop, ensilage, forage, or fodder. 		

COTTON

Application Methods and Equipment: OXYFLUORFEN 2EC may be applied as a post-direct spray to cotton a minimum of 6 to 8 inches tall. Care must be exercised to avoid spray contact with the cotton leaves. Use rigid precision ground spray equipment and spray shields to prevent spray contact with cotton foliage. Use branch lifters or shields, as necessary, to avoid contact of directed sprays with cotton plant.

Accurate, placement of spray nozzles is essential for uniform coverage of weeds and to minimize injury to cotton plants. Use a minimum broadcast spray volume of 20 gallons per acre and operate the sprayer at the minimum spray pressure recommended by the spray nozzle manufacturer. OXYFLUORFEN 2EC may be applied as a post-direct spray with only 2 flat fan nozzles per row (1 nozzle on each side of the row). For optimum coverage, use 4 flat fan nozzles per row (2 nozzles on each side of the row). The 2 forward nozzles should point forward and downward while the rear nozzles should point to the rear and downward. With either sprayer setup, nozzles should be carefully adjusted to cover the weed foliage with minimum contact to cotton plants. OXYFLUORFEN 2EC may also be applied as a band application. Do not use hollow cone nozzles.

Tank Mixing: For control of additional broadleaf and grass weeds, OXYFLUORFEN 2EC may be applied as a postemergence directed spray in tank mix combination with other herbicides registered for postemergence use in cotton (see Tank Mixing Precautions under Mixing Directions).

Weed Control	Rate (pt/acre)	Specific Use Directions
Postemergence	1 – 2	Apply as a post-directed spray. For optimum control, use the 2 pint per acre rate on actively growing weed seedlings with no more than 4 true leaves (not counting cotyledon leaves). Effective control of succulent weeds at the 2- to 3-leaf stage can usually be obtained at the 1 pint per acre rate. See Mixing Directions for surfactant recommendations. Where available, irrigation may be applied prior to application of OXYFLUORFEN 2EC to encourage maximum weed emergence. Irrigation following application will improve preemergence activity of OXYFLUORFEN 2EC against nightshade and groundcherry species.
Precautions:		
<ul style="list-style-type: none"> • Do not apply to cotton less than 6 inches tall or severe crop injury will result. • Exercise care to avoid spray contact with cotton leaves. Leaves accidentally sprayed will exhibit necrotic (dead) spots and may be dropped from the plant. Crop injury may be enhanced if application is made when excessive soil moisture is present or rainfall occurs immediately after application, however, cotton will outgrow this condition and develop normally. 		
Crop-Specific Restrictions:		
<ul style="list-style-type: none"> • Western Cotton (AZ and CA): Do not apply more than 2 pints (0.5 lb active) of OXYFLUORFEN 2EC per acre in a single application, or more than a total of 4 pints (1 .0 lb active) of OXYFLUORFEN 2EC per broadcast acre per season as a result of multiple applications. Do not apply within 75 days of harvest. • Southern Cotton (All other states): Do not apply more than 2 pints (0.5 lb active) of OXYFLUORFEN 2EC per acre of per season as a result of a single application or multiple applications. Do not apply within 90 days of harvest. 		

Key Weeds Controlled:

Postemergence	
cocklebur, common	nightshade, hairy
croton, tropic	pigweed, redroot
groundcherry, cutleaf	poinsettia, wild [†]
groundcherry, Wright	purslane, common
jimsonweed	sesbania, hemp
lambsquarter, common	sicklepod ^{††}

morningglory, annual (up to 6 leaf) nightshade, American black nightshade, black	sida, prickly (teaweed) [†] smartweed, Pennsylvania velvetleaf
---	---

[†] Multiple applications may be required for acceptable control.

^{††} Post-direct applications of OXYFLUORFEN 2EC will control or suppress seedlings not exceeding the one true leaf stage.

COTTONWOOD

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	4 – 6	OXYFLUORFEN 2EC may be applied as a single or split application. Apply as a directed spray to soil at the base of cottonwood trees. Use the higher rate in the rate range for extended preemergence weed control or for Postemergence control of weeds up to the 6 leaf stage. The addition of a non-ionic surfactant at 2 pints per 100 gallons of spray will enhance the Postemergence activity of OXYFLUORFEN 2EC on emerged weeds.
Precautions: <ul style="list-style-type: none"> Apply OXYFLUORFEN 2EC immediately after transplant only to dormant healthy cottonwood stock. In established stands, do not allow sprays of OXYFLUORFEN 2EC to contact cottonwood foliage. In newly established cottonwood plantings, use spray shields, if necessary, to prevent exposure of green bark and foliage. 		
Crop-Specific Restrictions: <ul style="list-style-type: none"> Do not apply more than 6 pints per acre of OXYFLUORFEN 2EC in a single application or more than 18 pints per acre per year. 		

Key Weeds Controlled:

groundsel, common knotweed, prostrate lambsquarter	mustard, hedge shepherdspurse smartweed, Pennsylvania
--	---

EUCALYPTUS

Apply OXYFLUORFEN 2EC for preemergence and postemergence control of listed broadleaf weeds in established eucalyptus plantings.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	4 – 6	<p>Directed Spray: OXYFLUORFEN 2EC may be applied as a single or split application. Apply as a directed spray to soil at the base of eucalyptus trees. Use the higher rate in the rate range for extended preemergence weed control or for postemergence control of weeds up to the 6 leaf stage.</p> <p>The addition of a non-ionic surfactant at the rate of 2 pints per 1 00 gallons of spray will enhance the postemergence activity of OXYFLUORFEN 2EC on emerged weeds.</p> <p>Over-the-Top Application: In new plantings, apply OXYFLUORFEN 2EC just before or immediately after transplanting eucalyptus seedlings that are in a dormant condition (i.e., leaves may be present, but terminal growth has hardened off and terminal buds have formed). In established plantings, OXYFLUORFEN 2EC may be applied as an over-the-top spray when plants are in a dormant condition.</p>
<p>Precautions:</p> <ul style="list-style-type: none"> At transplant, apply OXYFLUORFEN 2EC only to healthy "dormant" healthy eucalyptus stock. In established plantings, use spray shields, if needed, to prevent exposure of foliage and bark of small and/or actively growing plants. To avoid phytotoxicity, make over-the-top applications only to eucalyptus trees in a dormant condition. Do not make over-the-top applications after bud break and resumption of active growth. 		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> Do not apply more than 6 pints of OXYFLUORFEN 2EC per acre in a single application or more than 18 pints per acre per year. 		

Key Weeds Controlled:

Preemergence	Postemergence
burclover cheeseweed (malva) fiddleneck, coast filaree, broadleaf filaree, redstem filaree, whitestem groundsel, common lettuce, prickly pigweed, redroot purslane, common redmaids rocket, London shepherdspurse sowthistle, annual spurge, prostrate spurge, spotted	cheeseweed (malva) fiddleneck, coast filaree, broadleaf † filaree, redstem † filaree, whitestem † groundsel, common henbit minerslettuce nettle, burning pigweed, redroot redmaids shepherdspurse sowthistle, annual

† At the 6-pint rate, OXYFLUORFEN 2EC will provide control of filaree up to the 6-leaf stage.

USE ON FALLOW BEDS
(Not for use prior to planting soybeans in California)

Used alone or in tank mix combination with glyphosate, OXYFLUORFEN 2EC provides preemergence and/or postemergence control of winter annual broadleaf weeds on land to be planted to crops.

Prior to planting, treated fallow beds should be thoroughly tilled (incorporated) to a depth of at least 2.5 inches. OXYFLUORFEN 2EC is no longer herbicidally effective once the active layer in the soil surface is disrupted by soil incorporation.

Aerial Application: OXYFLUORFEN 2EC may be aerially applied for weed control in fallow beds. Follow requirements for Aerial Application in the Product Use Information section of this label.

Minimum Treatment to Planting Intervals for listed crops:

Direct Seeded Crops	Minimum Treatment-to-Planting Interval	
	OXYFLUORFEN 2EC (up to 1 pint/acre)	OXYFLUORFEN 2EC (>1 to 2 pints/acre)
carrot	90 days	90 days
cotton	7 days	7days
potato	60 days	60 days
sugar beet	60 days	90 days
other root/tuber crops	90 days	90 days
onions	180 days	180 days
other bulb vegetables	180 days	180 days
cabbage	90 days	90 days
cauliflower	90 days	90 days
other brassica crops	120 days	120 days
lettuce	90 days	120 days
other leafy vegetables (except brassica crops)	120 days	120 days
pepper	90 days	120 days
tomato	60 days	120 days
other fruiting vegetables	120 days	120 days
cantaloupe	60 days	90 days
squash	90 days	120 days
watermelon	60 days	60 days
other cucurbits	90 days	120 days
dry beans	60 days	60 days
Peanut	60 days	60 days
other legume vegetables	60 days	60 days
safflower	60 days	60 days
Soybeans (Except California)	7days	7 days
cereal grains: Including barley, buckwheat, corn, proso millet, pearl millet, oats, popcorn, rice, rye, sorghum, triticale, wheat, and wild rice	10 months	10 months
cotton and soybean	(see specific labeling for fallow beds to be planted to cotton or soybeans)	

Transplanted Crops	Minimum Treatment-to-planting Interval	
	OXYFLUORFEN 2EC (up to 1 pint/acre)	OXYFLUORFEN 2EC (>1 to 2 pints/acre)
celery	30 days	30 days
conifer	0 days	0 days
garlic	0 days	30 days
grape/kiwi	0 days	0 days
onion	0 days	30 days
pepper	30 days	30 days
strawberries	30 days	30 days
tomato	30 days	30 days
treefruit/nut/citrus	0 days	0 days

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	1 – 2	<p>Use 20 or more gallons of spray volume per acre and increase spray volume for dense weed growth.</p> <p>Use the 1 pint per acre rate for up to 4 weeks of preemergence control and postemergence control of susceptible weeds up to 4-leaf stage. Use the 2 pint per acre rate for up to 8 weeks of preemergence control and postemergence control of susceptible weeds up to 6-leaf stage. Best preemergence control is achieved when irrigation or rainfall occurs within 3 or 4 weeks after application.</p> <p>A tank mix with glyphosate is recommended if the treatment area contains dense weed populations, oversized weed seedlings, volunteer grains, annual grasses or under unfavorable environmental conditions.</p> <p>Outside of California: For enhanced contact activity (burndown/suppression) tank mix 6.5 fl oz of OXYFLUORFEN 2EC with the labeled rate of either glyphosate or paraquat (Gramoxone). Apply at the application rate and weed growth stages recommended in the respective tank mix product label.</p>
<p>Precautions:</p> <ul style="list-style-type: none"> • Failure to achieve thorough and complete incorporation, or to follow the recommended treatment-planting interval, may result in stand reduction and/or vigor reduction of the planted crop. • Crop injury may be enhanced if newly seeded crops or transplants are under stress due to drought, flooding, excessive fertilizer or soil salts, low soil temperatures, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects or diseases. • Exercise extreme care to avoid herbicide contact with any desirable dormant or non-dormant crop, plant, tree or vegetation as severe injury may result. 		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> • Do not apply more than 2 pints of OXYFLUORFEN 2EC per acre per fallow season. 		

Key Weeds Controlled: OXYFLUORFEN 2EC provides preemergence and postemergence control of the following weeds on fallow beds: †

buttercup, smallflower cheeseweed (malva) eveningprimrose, cutleaf †† fiddleneck, coast filaree, broadleaf filaree, redstem geranium, Carolina	mustard species nettle, burning oxalis pigweed, redroot purslane, common redmaids rocket, London
--	--

groundcherry, cutleaf groundsel, common henbit ladysthumb minerslettuce	shepherdspurse sida, prickly sowthistle, annual velvetleaf (wild common)
---	---

† Thorough spray coverage is essential to maximize the postemergence activity of OXYFLUORFEN 2EC. For postemergence control when applied by air, a tank mixture of OXYFLUORFEN 2EC with either glyphosate or paraquat (Gramoxone) is recommended.

†† Requires maximum rate and/or multiple applications for effective control.

GARBANZO BEANS
(For Use Only in Arizona and California)

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	1	Apply after planting but prior to weed or crop emergence as a single broadcast application using a spray volume of 20 or more gallons of water per acre.
Precautions: <ul style="list-style-type: none"> Garbanzo beans are tolerant to preemergence application of OXYFLUORFEN 2EC, however, under certain conditions, severe but temporary crop injury may occur. A heavy splashing rain shortly after crop emergence or wet soil conditions during early growth stages can cause leaf cupping, crinkling, stunting or defoliation of the garbanzo seedlings. Injury, when it occurs, it is usually limited to the first few leaves that develop after plants emerge from the soil. Delays in crop development and/or maturity may result, but Garbanzo beans do recover with little to no impact on yield. 		
Crop-Specific Restrictions: <ul style="list-style-type: none"> Do not apply more than 1 pint per acre of OXYFLUORFEN 2EC in a single application. Do not use bean vines for livestock feed or hay. 		

Key Weeds Controlled:

Preemergence
groundsel, common mallow, little rocket, London Shepherdspurse

GARLIC

Agricultural Use Requirements: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

Coveralls

- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Cultural Considerations: For optimum preemergence weed control, the soil surface should be smooth and free of excessive trash (clippings, plant residues, etc.). Following application, treated beds should be left undisturbed during the time period for which weed control is desired. Cultural practices that result in soil disturbance or redistribution or untreated soil can result in reduced weed control.

Direct Seeded Garlic (Postemergence Application):		
Weed Control	Rate (per acre)	Specific Use Directions
Postemergence	2 - 4 fl oz	Northeastern States Including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont: Apply OXYFLUORFEN 2EC at 2 to 4 fl oz per acre to seeded garlic that has at least 3 true leaves using ground equipment. Multiple treatments at 2 to 4 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre pre use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing. Application at later than 4-leaf growth stage may result in reduced weed control.
Postemergence	0.5 - 1 pt	Western States Including Arizona, Colorado, Idaho, Nevada, New Mexico, Oregon, Texas, Utah and Washington: Apply OXYFLUORFEN 2EC at 0.5 to 1 pt per acre to seeded garlic that has at least 2 true leaves using ground equipment. Multiple treatments at 0.5 to 1 pt per acre may be applied up to a maximum of 2.0 pints per acre pre use season. For optimum postemergence weed control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing. Application at later than 4-leaf growth stage may result in reduced weed control.
Postemergence	0.5 pt	All Other States: Apply OXYFLUORFEN 2EC at 0.5 pt per acre to seeded garlic that has at least 2 true leaves using ground equipment. Multiple treatments at 0.5 pt per acre may be applied up to a maximum of 2 pints per acre pre use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing. Application at later than 4-leaf growth stage may result in reduced weed control.

Direct Seeded Garlic (California Only)		
Weed Control	Rate (per/acre)	Specific Use Directions
Preemergence Postemergence	1 pt	<p>Application after planting but Prior to Garlic Emergence: Apply OXYFLUORFEN 2EC after planting, but prior to crop emergence, for preemergence control of listed broadleaf and grass weeds using ground, air or sprinkler irrigation (chemigation). Aerial Application: Apply in a minimum spray volume of 10 gallons per acre. Follow Aerial Application instructions and precautions in the Product Use Information section of this label.</p> <p>Postemergence and Directed Application: Apply OXYFLUORFEN 2EC as a directed or over-the-top spray to garlic that is at least 12 inches tall. Accurate, uniform placement of directed postemergence sprays is essential for effective weed control and to minimize injury to garlic. Use low-pressure sprays and a minimum spray volume of 20 gallons per acre. Adjust nozzles for minimum spray contact with garlic plants, directing the spray to the soil at the base of garlic plants and adjacent bed top and furrow area. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing. Application at later than 4-leaf growth stage may result in reduced weed control.</p> <p>Sprinkler Irrigation (Portable Lateral or Solid Set) Preemergence or Postemergence: Apply OXYFLUORFEN 2EC at the recommended broadcast application rate using sufficient irrigation to wet soil to a depth of 2 inches. Apply after planting but prior to garlic emergence or postemergence when garlic is at least 12 inches tall. Follow the application directions and precautions for "Sprinkler Chemigation" given in the Chemigation section of this label.</p>
<p>Precautions:</p> <ul style="list-style-type: none"> Garlic Response to Preemergence Applications of OXYFLUORFEN 2EC: Following a preemergence application of OXYFLUORFEN 2EC, a chlorotic band around some of the leaves may be observed after the first irrigation (or rainfall) following garlic emergence. 		

Transplanted Garlic: Postemergence Application Immediately after Planting		
Weed Control	Rate (per/acre)	Specific Use Directions
Postemergence	up to 2 pt	All States Except Northeastern States: Transplanted garlic is most tolerant of a postemergence application immediately after transplanting. An application of up to 2 pints per acre may be made within two days after transplanting. If less than 2 pints per acre is applied, a second application can be made two weeks or more after transplanting. Do not exceed the maximum use rate of 2 pints per acre of OXYFLUORFEN 2EC per season as a result of multiple applications.
Postemergence	2 - 4 fl oz	Northeastern States, including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont: Multiple treatments at 2 to 4 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre per use season.

Key Weeds Controlled:

<p>canarygrass (annual) eveningprimrose, cutleaf groundsel, common mallow, little (malva) nightshade, black pigweed, prostrate † pigweed, redroot †</p>	<p>puncturevine purslane, common † rocket, London sage, lanceleaf shepherdspurse † sowthistle, annual</p>
---	--

† Key weeds controlled at recommended rates in Northeastern States.

Garlic - Crop-Specific Precaution (Postemergence Application):

Postemergence applications of OXYFLUORFEN 2EC may cause chlorotic leaf banding, necrotic lesions, or stunting of the garlic plants. Symptoms may be more severe if garlic emerged under cool, wet, overcast, or foggy weather. These conditions are temporary and should not affect the vigor or development of garlic plants.

Garlic - Crop-Specific Restrictions (Applicable to All Methods of Application):

- In all states except Northeastern states, do not apply until direct seeded garlic plants have two (2) fully developed true leaves. In the Northeastern states, do not apply until direct seeded garlic plants have three (3) fully developed true leaves. Application made prior to the specified growth stage may result in serious crop injury and is not recommended.
- Do not apply more than a total of 2 pints per acre of OXYFLUORFEN 2EC per use season as a result of multiple applications.
- Do not apply within 60 days of harvest.
- In direct seeded garlic (except in California), do not apply OXYFLUORFEN 2EC as a preemergence treatment.
- Use only on dry bulb garlic.
- Do not apply to garlic grown for seed.
- For weed control in Garlic, do not mix OXYFLUORFEN 2EC with oils, surfactants, liquid fertilizers or pesticides except as specified on approved Albaugh Supplemental Labeling.
- Do not apply to garlic plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects, nematodes or diseases.

GUAVA (Bearing and Non-bearing)
(For Use Only in Hawaii)

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	5 – 8	<p>Preemergence or Postemergence: In established guava plantings, apply preemergence or postemergence to weeds. Increase the spray volume to ensure adequate coverage in high densities of emerged weeds or heavy trash. Minimize contact with guava plants by directing the spray to the soil surface. Spray shields are suggested to minimize spray contact in young plantings.</p> <p>For broader spectrum postemergence control of grass and broadleaf weeds, OXYFLUORFEN 2EC may be applied in tank mix combination with paraquat (Gramoxone) or glyphosate. Follow applicable use directions, precautions and limitations on the labels of the respective tank mix products.</p>
Postemergence	2 – 8	
<p>Precautions:</p> <ul style="list-style-type: none"> • Prevent direct spray or drift from contacting green stems, fruit or foliage, as injury may result. • Alone or in tank mix combination, OXYFLUORFEN 2EC should be applied to only healthy growing trees. • Application of OXYFLUORFEN 2EC should be made only after new foliage growth has hardened off. 		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> • Do not apply more than 8 pints per acre of OXYFLUORFEN 2EC in a single application or more than 16 pints per season. • Do not apply OXYFLUORFEN 2EC within 1 day of harvest. 		

Key Weeds Controlled:

Preemergence	Postemergence
ageratum buttonweed crotalaria purslane, common spurge, garden	purslane, common spurge, garden

HORSERADISH

Agricultural Use Requirements: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	2	Apply OXYFLUORFEN 2EC after the horseradish roots have been planted but prior to emergence of new horseradish leaves. Emerged leaves that receive direct or indirect spray (drift) contact will be injured. If necessary, cultivate before application to destroy germinated weeds.
<p>Precautions:</p> <p>Do not apply OXYFLUORFEN 2EC to horseradish plantings that have been weakened or stressed due to unfavorable temperature conditions, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture.</p>		
<p>Crop-Specific Restrictions:</p> <p>Do not apply more than 2 pints of OXYFLUORFEN 2EC per acre per crop.</p>		

Key Weeds Controlled:

lambquarters, common pigweed, redroot purslane, common	shepherdspurse smartweed, pennsylvania
--	---

JOJOBA

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	4 – 6	Initial application may be made when jojoba plants have reached a height of 6 inches or more. Use sufficient spray volume to ensure thorough coverage of dense weed growth. Sprays should be directed to the base of jojoba plants to avoid possible phytotoxicity to foliage. Spray shields are suggested for use in young plantings. Use higher rate in rate range for extended residual preemergence weed control. Make follow-up applications as necessary to maintain weed control. For early postemergence control of susceptible seedling weeds (less than 8 inches tall) apply OXYFLUORFEN 2EC at the rate of 4 pints per acre. OXYFLUORFEN 2EC may be applied at the rate of 6 pints per acre for postemergence control of weeds up to 12 inches tall. For optimum residual control, apply during the fall or winter months. Control may be unsatisfactory for weeds greater than 12 inches tall.
<p>Precautions:</p> <p>Avoid direct spray or drift contact with jojoba flowers or buds as severe injury may result. Over-the-top applications may cause burning, crinkling or bronzing of jojoba foliage, particularly to the youngest leaves, flowers, or buds present at the time of application.</p>		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> Do not apply more than 6 pints per acre per year. 		

Key Weeds Controlled:

Preemergence	Postemergence
burclover fiddleneck, coast filaree, broadleaf filaree, redstem filaree, whitestem groundsel, common henbit knotweed, prostrate lambsquarters, common lettuce, prickly mallow, little (malva, cheeseweed) pigweed, redroot purslane, common redmaids rocket, London shepherdspurse sowthistle, annual	fiddleneck, coast filaree, broadleaf ^{††} filaree, redstem ^{††} filaree, whitestem [†] groundsel, common [†] henbit mallow, little (malva, cheeseweed) minerslettuce nettle, burning pigweed, redroot [†] redmaids shepherdspurse sowthistle, annual

[†] Highest rate may be required for acceptable postemergence control.

^{††} OXYFLUORFEN 2EC at the 6-pint rate will provide control of filaree not exceeding the 4-inch stage. Applications to filaree beyond the 4-inch stage may result in partial control.

MINT (SPEARMINT AND PEPPERMINT)

Mint (Spearmint and Peppermint) Grown on Mineral Soils		
Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	4 – 6	Oregon and Washington (East of Cascades), California, Montana, Idaho, Nevada, South Dakota and Utah: Apply from December through March when mint is dormant. When used postemergence (to weeds), add an 80% active ingredient nonionic surfactant at the rate of one quart per 100 gallons of spray volume and apply before weeds exceed a height of 4 inches. Late winter applications will provide maximum activity on summer weeds, but summer grass control may be inconsistent. For best results, fall-plowed fields should be harrowed to provide a smooth surface for application. In furrow-irrigated fields, corrugating must be done prior to application. Corrugating or harrowing will result in disturbance of treated soil or movement of untreated soil into treated areas, resulting in poor weed control.
Preemergence	2 – 3	Peppermint (Western Oregon Willamette Valley): Apply OXYFLUORFEN 2EC from November through February to dormant peppermint only. Treatments in January or February generally provide better residual preemergence control of annual broadleaf weeds. Full season weed control should not be expected from this treatment.
<p>Precautions:</p> <ul style="list-style-type: none"> • Application must be made prior to emergence of new spring growth or severe crop injury may result. • In the Willamette valley, do not apply OXYFLUORFEN 2EC to mint that has been plowed. • Apply OXYFLUORFEN 2EC only to healthy stands of spearmint and peppermint. Do not apply to spearmint or peppermint weakened by disease, drought, flooding, excessive fertilizer, soil salts, previously applied pesticides, nematodes, insects, or winter injury, as severe injury may result. 		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> • Do not make more than one application of OXYFLUORFEN 2EC per season. 		

Key Weeds Controlled:

bedstraw, catchweed † bluegrass, annual flixweed goundsel, common lambsquarter, common lettuce, prickly (china lettuce) mustard, blue (purple mustard) mustard, tumble (Jim hill mustard) nightshade, hairy	† oats, wild orach, red pepperweed, yellowflower pigweed, redroot † ryegrass, Italian shepherdspurse sowthistle, annual tansymustard thistle, Russian
---	---

† Control of annual grasses is best obtained when OXYFLUORFEN 2EC is applied prior to emergence. Postemergence control of winter annual grasses is generally unsatisfactory if applications are made after the 1 to 2-leaf stage.

Mint (Spearmint and Peppermint) Grown on Muck Soils: For Use Only on Mint Grown in Indiana, Michigan, Montana, North Dakota, South Dakota, and Wisconsin		
Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	4 – 6	Note: Use directions in this section apply only to spearmint and peppermint grown on muck soils (organic matter content of 20% or greater). When used postemergence (to weeds), add an 80% active ingredient nonionic surfactant at the rate of one quart per 100 gallons of spray volume and apply before weeds exceed a height of 4 inches.
<p>Precautions:</p> <ul style="list-style-type: none"> • Application must be made prior to emergence of new spring growth or severe crop injury may result. • To avoid excessive crop injury, do not apply within 4 days of planting (sprigging) spearmint or peppermint. • Apply OXYFLUORFEN 2EC only to healthy spearmint or peppermint. Do not apply to spearmint or peppermint that has been weakened by disease, nematodes, soil insects, or winter injury, as severe injury may result. 		
<p>Crop-Specific Restrictions:</p> <p>Do not make more than one application of OXYFLUORFEN 2EC per season.</p>		

Key Weeds Controlled:

knotweed, prostrate pigweed, redroot purslane, common

NON-CROP USE

(Non-Food-Producing, Non-Cultivated Agricultural or Non-Agricultural Areas, such as Highway and Utility Rights-of-Way, Industrial Sites, Tank Farms, Storage Areas, Airports, Fencerows, and Farmsteads)

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	5 – 8	Preemergence: Use higher rate in rate range for longer residual control. Postemergence: Use the lower rate in the rate range for control of susceptible weeds in the early postemergence stage, less than 4 inches tall. Use the higher rate for weeds up to 12 inches tall. Application to weeds beyond the 4-inch stage may result in partial control.
Postemergence	2 – 8	
<p>Tank Mixing: Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.</p> <ul style="list-style-type: none"> • Preemergence: For broader-spectrum residual preemergence weed control, OXYFLUORFEN 2EC may be applied in tank mix combination diuron (Karmex) or simazine. • Postemergence: For additional postemergence control of susceptible grass and broadleaf weeds, OXYFLUORFEN 2EC may be applied in tank mix combination with paraquat (Gramoxone) or glyphosate. 		
<p>Site-Specific Restrictions:</p> <ul style="list-style-type: none"> • Do not feed or allow animals to graze on any areas treated with OXYFLUORFEN 2EC. • Do not apply more than 8 pints per acre in a single application. 		

Key Weeds Controlled:

Preemergence	Postemergence
burclover cheeseweed (malva) fiddleneck, coast filaree, broadleaf filaree, redstem groundsel, common henbit knotweed, prostrate lambsquarters, common lettuce, prickly pigweed, redroot purslane, common redmaids rocket, London shepherdspurse sowthistle, annual	cheeseweed (malva) fiddleneck, coast filaree, broadleaf filaree, redstem groundsel, common henbit minerslettuce nettle, burning pigweed, redroot purslane, common redmaids shepherdspurse sowthistle, annual

ONIONS

Agricultural Use Requirements: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Cultural Considerations: For maximum preemergence activity, the soil surface should be smooth and free of excessive trash (clippings, plant residues, etc.). Following application, cultural practices that result in redistribution or disturbance of the soil surface or move untreated soil into treated areas will reduce weed control. For best results, make applications to established beds that are left undisturbed during the time period for which weed control is desired.

Direct Seeded Onions: Postemergence Application		
Weed Control	Rate (per acre)	Specific Use Directions
Postemergence	2 - 4 fl oz	Northeastern States Including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont: Apply OXYFLUORFEN 2EC at 2 to 4 fl oz per acre to seeded onions that have at least 3 true leaves using ground equipment. Multiple treatments at 2 to 4 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre pre use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing.
Postemergence	0.5 - 1 pt	Western States Including Arizona, California, Colorado, Idaho, Nevada, New Mexico, Oregon, Texas, Utah and Washington: Apply OXYFLUORFEN 2EC at 0.5 to 1 pt per acre to seeded onions that have at least 2 true leaves using ground equipment. Multiple treatments at 0.5 to 1 pt per acre may be applied up to a maximum of 2.5 pints per acre per use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing.
Postemergence	0.5 pt	All Other States: Apply OXYFLUORFEN 2EC at 0.5 pt per acre to seeded onions that have at least 2 true leaves using ground equipment. Multiple treatments at 0.5 pt per acre may be applied up to a maximum of 2 pints per acre per use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing.
Postemergence	(see above)	Sprinkler Irrigation - All Except Northeastern States (Center Pivot, Portable Lateral or Solid Set): Apply OXYFLUORFEN 2EC at the recommended broadcast application rate using sufficient irrigation to wet soil to a depth of 2 inches. Follow the application directions and precautions for "Sprinkler Chemigation" given in the Chemigation section of this label.

Transplanted Onions: Application Immediately before Planting		
Weed Control	Rate (per/acre)	Specific Use Directions
Preemergence Postemergence	1 – 2 pt	Pre-transplant Application (Not for Use in Northeastern States or Western States: OXYFLUORFEN 2EC may be applied as a broadcast or band application after completion of tillage operations, but before transplanting of onion plants. Transplanting should be accomplished with a minimum of soil disturbance and, for optimum weed control, soil surfaces should be left undisturbed after transplanting for the period for which weed control is desired. However, timely cultivation after weed emergence will assist in weed control. If less than 2 pt per acre was applied as a pre-transplant application, postemergence applications may be made as instructed for seeded onions. Do not exceed the maximum use rate of 2 pt per acre per use season as a result of multiple applications.
Transplanted Onions: Application Immediately after Planting		
Application Timing for Target Weeds	Rate (per/acre)	Specific Use Directions
Preemergence	up to 2 pt	All States Except Northeastern States: Transplanted onions are most tolerant of a postemergence application immediately after transplanting. An application of up to 2 pints per acre may be made within two days after transplanting. If less than 2 pints per acre is applied, a second application can be made two weeks or more after transplanting. Do not exceed the maximum use rate of 2 pints per acre of OXYFLUORFEN 2EC per season as a result of multiple applications.
Preemergence	2 - 4 fl oz	Northeastern States including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont: Multiple treatments at 2 to 4 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre per use season.
Onions - Use Precautions (Applicable to All Areas and Methods of Application):		
<ul style="list-style-type: none"> • OXYFLUORFEN 2EC can cause necrotic lesions, twisting, pigtailling or stunting of the onion plants. Injury will be more severe if applications are made immediately following or during cool, wet weather and/or if applications are made prior to the specified onion growth stage of the onion plants as specified in Specific Use Directions. • Do not apply to onion plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects, nematodes or diseases. 		
Onions - Crop-Specific Restrictions (Applicable to All Areas and Methods of Application):		
<ul style="list-style-type: none"> • In all states except Northeastern states, do not apply until direct seeded onion plants have at least two (2) fully developed true leaves. In the Northeastern states, do not apply until direct seeded onion plants have at least three (3) fully developed true leaves. Application made prior to the specified growth stage may result in serious crop injury and is not recommended. • Do not apply more than a total of 2 pints per acre of OXYFLUORFEN 2EC per use season as a result of multiple applications. • Do not apply within 45 days of harvest. • Do not apply OXYFLUORFEN 2EC as a preemergence treatment to direct seeded onions. • Use only on dry bulb onions. • Do not apply to onions grown for seed, except as instructed in separate use directions. • For Arizona, California, Idaho, Oregon, New Mexico, Nevada, Utah and Washington only, tank mixtures of OXYFLUORFEN 2EC with oils, surfactants, liquid fertilizers or other pesticides may be made but could result in unexpected results including enhanced crop response or injury. • See the Inherent Risks of Use section of the Terms and Conditions of Use at the end of the label for more information. 		

Key Weeds Controlled:

Postemergence
canarygrass (annual) eveningprimrose, cutleaf ^(a) groundsel, common mallow, little (malva) nightshade, black pigweed, prostrate ^(b) pigweed, redroot ^(a, b) puncturevine purslane, common ^(a, b) rocket, London sage, lanceleaf shepherdspurse ^(b) sowthistle, annual

^(a) Weeds controlled when applied as a pre-transplant application. In addition, OXYFLUORFEN 2EC at the rate of 1 to 2 pints per acre will provide control/suppression of carpetweed, Pennsylvania smartweed, galinsoga, common lambsquarters, and wild mustard. Applications of OXYFLUORFEN 2EC to muck soils may result in partial control or suppression of the weeds listed.

^(b) Specific weeds controlled at rates recommended for use in northeastern states (see DOSAGE section).

ONIONS GROWN FOR SEED

<p>Agricultural Use Requirements: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.</p> <p>PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:</p> <ul style="list-style-type: none"> Coveralls Chemical-resistant footwear plus socks Chemical-resistant gloves made of any waterproof material Shoes plus socks
--

Weed Control	Rate (per/acre)	Specific Use Directions
Preemergence	2 fl oz	Northeastern States including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont: Multiple treatments at 2 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre pre use season. Prior to initial treatment, seeded onions must have at least four (4) true leaves. Multiple treatments at the aforementioned rate may be applied.
Preemergence	up to 0.5 pt	All other States: Apply OXYFLUORFEN 2EC at up to 0.5 pt per acre to seeded onions that have at least three (3) true leaves. Multiple treatments at 0.5 pt per acre may be applied up to a maximum of 2 pints per acre pre use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing. Sprinkler Irrigation - Portable Lateral or Solid Set: Apply OXYFLUORFEN 2EC at the recommended broadcast application rate using sufficient irrigation to wet soil to a depth of 2 inches. Follow the application directions and precautions for "Sprinkler Chemigation" given in the Chemigation section of this label.

Weed Control	Rate (per/acre)	Specific Use Directions
<p>Use Precautions:</p> <ul style="list-style-type: none"> • Notice: Some varieties or inbred lines of onions may be more susceptible to OXYFLUORFEN 2EC. Care should be taken to ensure that the particular onion variety or line being grown is tolerant to OXYFLUORFEN 2EC. It is suggested that all onion varieties or lines be tested in limited areas to ensure an adequate level of crop tolerance prior to an application for postemergence weed control. • OXYFLUORFEN 2EC can cause necrotic lesions, twisting, pigtailling or stunting of the onion plants. Injury will be more severe if applications are made immediately following or during cool, wet weather and/or if applications are made prior to the specified onion growth stage of the onion plants as specified in Specific Use Directions. • Do not apply to onion plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects or diseases. 		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> • In all states, do not apply OXYFLUORFEN 2EC until the onions have reached the minimum leaf stage specified. Application prior to the recommended stage of development may result in serious injury and is not recommended. • Do not apply more than a total of 2 pints per acre of OXYFLUORFEN 2EC during one use season. • Do not apply within 60 days of harvest. • For seeded onions, do not apply OXYFLUORFEN 2EC with oils, surfactants, liquid fertilizers or other pesticides except as specified in approved Albaugh Supplemental Labeling. 		

Key Weeds Controlled:

Postemergence
<p>canarygrass (annual) eveningprimrose, cutleaf groundsel, common mallow, little (malva) nightshade, black pigweed, prostrate † pigweed, redroot † puncturevine purslane, common † rocket, London sage, lanceleaf shepherdspurse sowthistle, annual</p>

† Specific weeds controlled at rates recommended for use in northeastern states (see DOSAGE section).

PAPAYA (Hawaii Only)

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	4	<p>The initial application should occur no sooner than 4 months after transplanting or 6 months after direct seeding, and after the papaya has reached a minimum height of 4 feet. Applications may be repeated at approximate 4-month intervals.</p> <p>Apply preemergence or postemergence to weeds. Increase the spray volume to assure adequate coverage of dense growth of emerged weeds. OXYFLUORFEN 2EC must be applied as a directed spray to the orchard floor beneath the papaya plants. Accurate, uniform placement of OXYFLUORFEN 2EC is essential for effective weed control and to minimize crop injury. OXYFLUORFEN 2EC must be applied using rigid precision ground sprayer equipment.</p> <p>Postemergence applications may be made up to the 4 leaf stage of weed growth.</p>
<p>Precautions:</p> <ul style="list-style-type: none"> Do not allow the herbicide solution, spray, drift or mist to contact green bark, stems, fruit or foliage as injury may result. Do not use OXYFLUORFEN 2EC on papaya plantings that are weak, or under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture. 		
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> Do not apply more than 4 pints of OXYFLUORFEN 2EC per broadcast acre in a single directed spray or more than 12 pints per broadcast acre per year as a result of multiple applications. Do not apply OXYFLUORFEN 2EC within 1 day of harvest. 		

Key Weeds Controlled:

amaranth, spiny purslane, common	spurge, garden
-------------------------------------	----------------

SOYBEANS
(Not for Use in California)

SOYBEANS – EARLY PREPLANT APPLICATION IN CONSERVATION TILLAGE SYSTEMS		
Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	1.5 – 3	<p>Early Preplant Application: Surface apply OXYFLUORFEN 2EC to the stale seedbed approximately 14 days before planting conservation tillage soybeans for postemergence and preemergence residual broadleaf control. Use a spray volume of 20 or more gallons per acre and increase the spray volume if growth of existing weed is dense. OXYFLUORFEN 2EC at 2 to 3 pints provides early season suppression of annual grasses, but should not be relied upon as a basic grass herbicide. A planned program utilizing herbicides registered for early preplant, preemergence or postemergence grass control in soybeans is recommended.</p> <p>Use of ridge or slot planter or a similar planting implement that causes minimal soil disturbance is recommended. Movement or redistribution of surface soil will reduce herbicidal effectiveness.</p>

SOYBEANS: NO-TILL (Double-Crop)		
Application Timing For Target Weeds	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	0.5 – 2	Preemergence Application to Soybeans: Applied preemergence, OXYFLUORFEN 2EC provides postemergence and residual preemergence control of susceptible broadleaf weeds. Apply OXYFLUORFEN 2EC within one day after planting. Later applications may result in severe crop injury and are not recommended. Apply in a minimum spray volume of 20 gallons per acre and increase spray volume if growth of existing weeds is dense.
Tank Mixing: For enhanced postemergence control of existing grass and broadleaf weeds, OXYFLUORFEN 2EC may be tank mixed with paraquat (Gramoxone) or glyphosate. For extended residual control of annual grassed no-till soybeans, OXYFLUORFEN 2EC may also be tank mixed with a residual grass herbicide such as Bronco Herbicide, Dual Magnum Herbicide, or Lasso Herbicide.		
Postemergence	1	Postemergence Directed Application: OXYFLUORFEN 2EC may be applied as a post-directed application. Optimum control is achieved when OXYFLUORFEN 2EC is applied to seedling weeds not exceeding 4 true leaves (not counting cotyledon leaves) and actively growing. Use of an 80% nonionic surfactant cleared for application to growing crops at the rate of 2 pints per 100 gallons of spray is recommended whenever postemergence weed control is desired. For Postemergence application, Soybeans must be a minimum 8 inches tall. Use a minimum of 2 flat fan nozzles per row. Use branch lifters or shields to prevent excessive spray contact to the soybean plants. Do not use hollow cone nozzles.

Soybeans: Grown Under Conventional Tillage Systems		
Application Timing For Target Weeds	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	1 – 1.5	Preemergence Application to Soybeans: OXYFLUORFEN 2EC provides preemergence control of susceptible broadleaf weeds. Apply OXYFLUORFEN 2EC within one day after planting. Later applications may result in severe crop

		injury and are not recommended. Apply in a minimum spray volume of 20 gallons per acre and increase spray volume if growth of existing weeds is dense. The 1.5 pint per acre rate will assist in early season annual grass control but should not be relied upon as a basic grass herbicide. OXYFLUORFEN 2EC may also be applied as a preemergence application following a preplant incorporated grass herbicide treatment.
<p>Preemergence Tank Mixes (To Control Additional Grass and Broadleaf Weeds): Apply preemergence tank mixes of OXYFLUORFEN 2EC within one day after planting. Later applications may result in severe crop injury and are not recommended.</p> <ul style="list-style-type: none"> • OXYFLUORFEN 2EC at 0.6 to 1.5 pints per acre may be applied preemergence to soybeans in tank mix with Dual Magnum Herbicide or Lasso Herbicide. OXYFLUORFEN 2EC may be applied alone as a preemergence application following a preplant incorporated grass herbicide application or as a tank mix in a preemergence application with Dual Magnum, or Lasso herbicides. Refer to the label of tank mix product for additional weeds controlled. • OXYFLUORFEN 2EC at 0.6 to 0.8 pints per acre may be applied preemergence to soybeans in tank mix with 1 to 1.67 pints of Command 6EC herbicide. Refer to the label for Comman 6EC for additional weeds controlled. 		
Postemergence	1	Postemergence Directed Sprays: OXYFLUORFEN 2EC may be applied as a post-directed application at 1 pint per acre. Optimum control is achieved when weeds not exceed 4 true leaves and are actively growing (do not count cotyledon leaves). Use of an 80% nonionic surfactant cleared for application to growing crops at the rate of 2 pints per 100 gallons of spray is recommended whenever postemergence weed control is desired. For postemergence application, Soybeans must be a minimum 8 inches tall. Use a minimum of 2 flat fan nozzles per row. Use branch lifters or shields to prevent excessive spray contact to the soybean plants. Do not use hollow cone nozzles.
<p>Postemergence Tank Mixes: For broader spectrum control or broadleaf weeds, OXYFLUORFEN 2EC may be applied in tank mix with Butoxone Herbicide or Butyrac 200 Herbicide. Use 1 pint of OXYFLUORFEN 2EC with 1 pint of Butoxone or 0.7 to 0.9 pint of Butyrac 200 per acre. Refer to label of tank mix product for additional weeds controlled.</p>		

Soybeans – Precautions (All Methods and Timings to Soybeans):

- Soybeans are tolerant to preemergence and post-directed applications of OXYFLUORFEN 2EC at recommended rates, however, under certain conditions injury may occur. Heavy splashing rain shortly after crop emergence or cold, wet soil conditions during early growth stages can cause leaf cupping and crinkling. When injury occurs, it is generally limited to the first few leaves that develop after crop emergence. Soybeans recover from this injury and yields are not adversely affected. Soybeans accidentally sprayed during a post-directed application will exhibit necrotic spotting and injury to the soybean plant. Exercise care to avoid spray contact with the soybean leaves.

Soybeans – Crop-Specific Restrictions:

- Tank Mixing: Read and observe all label directions before using. Follow applicable use directions, precautions and limitations on the labels of the respective tank mix products. Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on respective product labels. In interpreting the labels of tank mixed products, the most restrictive limitations must apply.
- Do not make more than two applications of OXYFLUORFEN 2EC per growing season.
- Do not apply more than 2 pints (0.5 lbs active) of OXYFLUORFEN 2EC per acre during one growing season as a result of preemergence application in no-till (double-crop) or conventional till soybeans, or post-directed in conventional till soybeans. If early preplant application is made, do not apply more than 3 pints (0.75 lb active) of OXYFLUORFEN 2EC per acre during one growing season.
- Do not apply a post-directed application of OXYFLUORFEN 2EC to soybeans after the initial appearance of blooms.

Key Weeds Controlled (OXYFLUORFEN 2EC Alone):

Preemergence	Postemergence
groundcherry, cutleaf † jimsonweed lambsquarters, common nightshade, American black † pigweed, redroot poinsettia, wild shepherdspurse sida, prickly (teaweed) smartweed, Pennsylvania sowthistle, common † velvetleaf	cocklebur, common croton, tropic groundcherry, cutleaf groundcherry, Wright jimsonweed lambsquarters, common morningglory, annual (up to 6 leaf) mustard, wild nightshade, American black nightshade, black nightshade, hairy pigweed, redroot † poinsettia, wild purslane, common sesbania, hemp shepherdspurse sicklepod †† sida, prickly (teaweed) † smartweed, Pennsylvania velvetleaf

† Multiple applications may be required for acceptable control.

†† Post-directed applications of OXYFLUORFEN 2EC will kill or suppress seedlings not exceeding the one true leaf stage.

TARO
(For Use Only in Hawaii)

For use only to dryland taro grown in Hawaii. Dryland taro is defined as taro grown without irrigation, or by using irrigation practices that do not result in run-off, irrigation return flow, or other loss of irrigation water from the production area. If irrigation is used, the water applied shall not exceed the field capacity of the soil.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	2	Preemergence to Taro and Weeds: A single application of OXYFLUORFEN 2EC at the rate of 2 pints per acre may be applied within 1 week after transplanting but prior to emergence of taro plants.
Postemergence	1	Postemergence to Taro and Weeds: OXYFLUORFEN 2EC may be applied as a post-directed or band application at the rate of 1 pint per acre. Effective control of succulent weed seedlings in the 2-to 3-leaf stage can usually be obtained. Applications to weeds beyond the 3-leaf stage may result in partial control.

Precautions:

Accurate, uniform placement of OXYFLUORFEN 2EC is essential for effective weed control and to minimize crop injury. Taro foliage receiving accidental spray or drift will be injured. OXYFLUORFEN 2EC must be applied using rigid precision ground sprayer equipment.

Occasionally, after the use of OXYFLUORFEN 2EC, spotting, crinkling or flecking may appear on the leaves of the taro. Leaves that receive direct or indirect (drift) spray contact will be injured.

Do not use OXYFLUORFEN 2EC on taro plantings that are weak, or under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture.

Crop-Specific Restrictions:

- Do not apply more than 2 pints of OXYFLUORFEN 2EC per broadcast acre as a single preemergence application.
- Do not apply more than 1 pint of OXYFLUORFEN 2EC per acre in a single post-direct spray or more than 2 pints per acre per season as a result of multiple post-directed applications.
- Do not apply more than 4 pints of OXYFLUORFEN 2EC per acre per season as a result of preemergence and post-direct applications.
- Do not apply OXYFLUORFEN 2EC within 6 months of harvest of taro (corms, leaves).

Key Weeds Controlled:

amaranth, spiny
purslane, common
spurge, garden

~~TREEFRUIT~~ TREE FRUIT/NUT/VINE CROPS (Dormant Application)

Almond, Apple, Apricot, Avocado, Beechnut, Brazil Nut, Butternut, Cashew, Cherry, Chestnut, Chinquapin, Crab Apple, Date, Feijoa, Fig, Filbert, Grapes, Hickory Nut, Kiwi, Loquat, Macadamia Nut, Mayhaws, Nectarine, Olives, Peach, Pear, Pecan, Persimmon, Pistachio, Plum, Pomegranates, Prune, Quince, and Walnut

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence (broadcast application)	5 – 6	Apply OXYFLUORFEN 2EC a minimum of 20 gallons of water per acre. Use higher spray volumes to ensure thorough coverage in high densities of emerged weeds or heavy trash. Sprays should be directed to the soil and the base of dormant trees or vines.
(banded application)	5 - 8	In California, OXYFLUORFEN 2EC may be applied as an over-the-top or directed spray to dormant nonbearing grape plantings. The use of a low-pressure sprayer is suggested. Do not apply over-the-top to grape plantings that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, injury from previously applied pesticides, or injury due to insects, nematodes, or diseases, as severe crop injury may result.
Postemergence (broadcast application)	2 – 6	Apply in a spray volume of 40 or more gallons per acre. For optimum control, apply when weeds are at seedling stage of growth.
(banded application)	2 - 8	The lower rate in the rate range (2 pints per acre) is recommended for the control of susceptible seedling weeds in the early postemergence stage up to the 4-leaf stage. Higher rates (up to 6 pints per acre) may be used for weeds up to the 6-leaf stage. Applications to weeds beyond the 6-leaf stage may result in partial control.
<p>Tank Mixing: Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply. See labels of tank mix partners to determine suitability and use rates for various crops.</p> <ul style="list-style-type: none"> • Postemergence: For broader spectrum postemergence control of listed grass and broadleaf weeds, OXYFLUORFEN 2EC may be applied in tank mix with paraquat (Gramoxone) or glyphosate. These herbicides may also be added to preemergence tank mixes for enhanced control of existing weeds. • Preemergence: For broad-spectrum preemergence control of susceptible grass and broadleaf weeds in listed treefruit, nut or vine plantings, OXYFLUORFEN 2EC may be applied in tank mix with napropamide (Devrinol herbicide), diuron (Karmex herbicide), pronamide (Kerb⁸ herbicide), simazine, norflurazon (Solicam herbicide) or oryzalin (Surflan herbicide). 		
<p>Chemigation (All States): For dormant season application using sprinkler (low-volume (micro sprinkler), drip (trickle), and flood (basin) irrigation systems, apply OXYFLUORFEN 2EC at the specified rate per acre. Follow applicable directions in the Chemigation section of this label when making applications using irrigation systems.</p>		
<p>Precautions:</p> <ul style="list-style-type: none"> • OXYFLUORFEN 2EC or any of the combinations recommended on this label should be applied to only healthy growing trees or vines. • Avoid direct plant contact. Direct spray toward the base of tree or vines unless specific use recommendations allow over-the-top application. 		

Weed Control	Rate (pt/acre)	Specific Use Directions
<p>Crop-Specific Restrictions:</p> <ul style="list-style-type: none"> In all states, unless otherwise specified, do not apply OXYFLUORFEN 2EC during the period between bud swell and completion of final harvest or when fruit/nuts are present. OXYFLUORFEN 2EC may be applied upon completion of final harvest. In Arizona and California, OXYFLUORFEN 2EC may be applied during the period following completion of final harvest up to February 15 (February 1st in the Coachella Valley, California). Applications made after these calendar dates, but prior to bud swell, may result in significant crop injury and are the responsibility of the user. For banded applications, up to 8 pints per acre of OXYFLUORFEN 2EC per use season may be applied within the treated band. Do not apply more than a maximum of 6 pints per acre per use season on a broadcast basis. Do not apply to grapes or kiwi established less than 3 years unless vines are on a trellis wire a minimum of 3 feet above the soil surface. Do not apply to grapes or kiwi that are not staked or trellised unless vines are free standing. 		

Key Weeds Controlled (Arizona and California):

Preemergence	Postemergence
burclover cheeseweed (malva) fiddleneck, coast filaree, broadleaf filaree, redstem filaree, whitestem groundsel, common henbit knotweed, prostrate lambsquarter, common lettuce, prickly pigweed, redroot purslane, common redmaids rocket, London shepherdspurse sowthistle, annual	cheeseweed (malva) fiddleneck, coast filaree, broadleaf † filaree, redstem † filaree, whitestem † groundsel, common henbit minerslettuce nettle, burning pigweed, redroot redmaids shepherdspurse sowthistle, annual

† OXYFLUORFEN 2EC at the 6-pint rate will provide control of filaree not exceeding the 4-inch stage. Applications to filaree beyond the 4-inch stage may result in partial control.

Key Weeds Controlled (All Other States Except Arizona and California):

Preemergence	Postemergence
camphorweed cudweed, narrowleaf eveningprimrose, cutleaf † groundcherry, cutleaf jimsonweed lambsquarters, common nightshade, American black nightshade, black pepperweed, Virginia pigweed, redroot poinsettia, wild sida, prickly smartweed, Pennsylvania sowthistle, annual spurge, prostrate spurge, spotted velvetleaf	balsamapple cocklebur, common cudweed, narrowleaf †† eveningprimrose, cutleaf ††† groundcherry, cutleaf groundcherry, Wright jimsonweed lambsquarters, common morningglory, annual nightshade, American black nightshade, black pepperweed, Virginia pigweed, redroot poinsettia, wild purslane, common sesbania, hemp shepherdspurse sida, prickly (teaweed) smartweed, Pennsylvania sowthistle, annual velvetleaf

† Highest rate and/or multiple applications may be required for acceptable control.

†† Maximum 0.5-inch diameter.

††† Highest rate and/or multiple applications may be required for acceptable control.

GRAPES (Non-Dormant Application)
(California Only)

OXYFLUORFEN 2EC may be applied as a directed spray or, for supplemental preemergence weed control, through low-volume sprinkler (micro sprinkler) or drip irrigation systems for control or suppression of listed broadleaf weeds in non-dormant grapes (raisin and wine grapes only). OXYFLUORFEN 2EC may also be applied to all grapes (raisin, table, and wine) as a dormant season application. Refer to Treefruit/Nut/Vine Crops (Dormant Application) section above for use directions for dormant season application to grapes.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	2	OXYFLUORFEN 2EC may be applied preemergence or postemergence to weeds either as a directed spray in a minimum spray volume of 20 gallons per acre or through low-volume sprinkler (micro sprinkler) or drip irrigation systems. Repeat applications may be required. Applications may be made from completion of bloom up to 14 days before to harvest. When applied as a postemergence directed spray, add 1 quart 80% active nonionic surfactant cleared for application to growing crops per 100 gallons of spray. Sprays should be directed to the soil and the base of vines.
Postemergence	1 – 2	
<p>Tank Mixing:</p> <ul style="list-style-type: none"> When applied as a directed postemergence spray using ground equipment, OXYFLUORFEN 2EC may be applied in tank mix with paraquat (Gramoxone) or glyphosate in a minimum spray volume of 10 gallons per acre. Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply. 		
<p>Chemigation: Follow chemigation instructions in Product Use Information section.</p> <ul style="list-style-type: none"> Low Volume Sprinkler (Microsprinkler) and Drip (Trickle) Irrigation: Apply only through low-volume sprinkler or drip systems designed to uniformly distribute irrigation water beneath the canopy. Meter OXYFLUORFEN 2EC at a continuous rate during the middle 1/3 of the irrigation period and discontinue application during the final 1/3 of the irrigation period to ensure proper flushing of the irrigation system. Use of OXYFLUORFEN 2EC through low-volume sprinklers or drip emitters helps to reduce the "ring effect" of weed escapes in areas around sprinklers or emitters where previously applied broadcast or directed treatments begin to break down. 		
<p>Precautions:</p> <ul style="list-style-type: none"> Crop Tolerance: The use of OXYFLUORFEN 2EC may result in varying degrees of injury to non-dormant grapes. Grape foliage will typically exhibit injury symptoms from direct or indirect (spray drift, soil contact) exposure. This injury may result in necrosis, reddening, cupping or crinkling of grape leaves. The grape plant will continue to grow normally. Grape leaves that are immature or expanding at the time of contact with OXYFLUORFEN 2EC are the most susceptible to foliage injury. Grapes may exhibit some small blemishes (spots or flicks) on the fruit. OXYFLUORFEN 2EC is phytotoxic to plant foliage. Avoid drift to all other crops and nontarget areas. Do not apply when weather conditions favor drift. 		

Crop-Specific Use Restrictions:

- The total amount of OXYFLUORFEN 2EC applied during one season (from completion of final harvest through dormancy to non-dormant use covered by this section) cannot exceed 6 pints per acre as a result of multiple applications in any given area (broadcast, banded, or within the wetted area of the low-volume sprinkler or drip irrigation system).
 - Do not apply within 14 days of harvest.
 - Do not initiate application of OXYFLUORFEN 2EC in non-dormant grapes until the completion of the bloom period.
 - Do not apply to grapes established less than 3 years unless vines are either on a trellis wire a minimum of 3 feet above the soil surface, or protected by grow tubes.
 - OXYFLUORFEN 2EC should be applied only by ground application equipment of through low-volume sprinkler (micro sprinkler) or drip (trickle) irrigation systems.
- Apply OXYFLUORFEN 2EC as a non-dormant application to wine grapes or raisin grapes only.

Key Weeds Controlled or Suppressed:

Preemergence	Postemergence
burclover cheeseweed, malva fiddleneck, coast groundsel, common henbit knotweed, prostrate lambsquarters, common minerslettuce mustard, black nettle, burning nightshade, black pigweed, redroot purslane, common redmaids rocket, London sowthistle, annual	cheeseweed, malva fiddleneck, coast groundsel, common henbit minerslettuce morningglory species, annual mustard, black nettle, burning nightshade, black pigweed, redroot purslane, common redmaids rocket, London sowthistle, annual

SUCKER CONTROL IN NON-DORMANT GRAPES
 (Washington and Oregon Only)
 (Grapes for Wine and Processing Only)

Application Timing for Sucker Control	Rate (pt/acre)	Specific Use Directions
Grape suckers less than 12 inches in length.	1 -2	Apply OXYFLUORFEN 2EC in a three-foot band directed towards to newly emerging suckers at the base of the grapevine. The highest rate and/or a second application may be required to achieve an acceptable level of control/suppression of grape suckers. Avoid spray contact on flowers, grape clusters, or fruit. Use mounted nozzles to deliver the spray solution. Thorough spray coverage of sucker growth is essential for optimal activity. Use a spray volume of 50 or more gallons per acre (broadcast basis).

Tank Mixing: For enhanced postemergence sucker activity, a tank mixture of OXYFLUORFEN 2EC with either glufosinate (Rely Herbicide) or paraquat (Gramoxone) can be used. Apply at the recommended rates and growth stages in a manner describe on the respective labels. Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.

Precautions:

- The use of OXYFLUORFEN 2EC may result in varying degrees of injury to non-dormant grapes. Grape foliage will typically exhibit injury symptoms from direct or indirect (spray drift or soil contact) exposure. This injury may result in necrosis, reddening, cupping or crinkling of grape leaves. The grape plant will continue to grow normally. Leaves that are immature or expanding at the time of contact with OXYFLUORFEN 2EC are the most susceptible to injury. Grape fruit may exhibit some small blemishes (spots or flecks) on the fruit.

Crop-Specific Restrictions:

- The total amount of OXYFLUORFEN 2EC applied during one crop year (dormant and non-dormant) cannot exceed 6 pints per acre as a result of multiple applications in any give area (broadcast or banded).
- OXYFLUORFEN 2EC should be applied only by ground application equipment.
- Apply OXYFLUORFEN 2EC as a non-dormant application for sucker control only to wine or processed grapes.
- Do not apply OXYFLUORFEN 2EC within 60 days of harvest.

PISTACHIOS, WALNUTS, ALMONDS (CALIFORNIA ONLY)
(Non-Dormant Application)

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	5 – 6	Preemergence: For residual weed control of listed weeds.
Postemergence	1 – 2	Postemergence (Suppression): Apply to seedling weeds less than 4 inches in height. Repeat applications may be required.
	2 – 6	Postemergence (Cleanup): Contact (postemergence) control for cleanup sprays and preharvest applications. Apply to seedling weeds less than 4 inches in height. Applications to weed seedlings beyond the 4-inch stage may result in partial control.
<p>Tank Mixing: For broader spectrum grass and broadleaf weed control in tree row middles, OXYFLUORFEN 2EC may be tank mixed with either paraquat (Gramoxone) or glyphosate. Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.</p>		
<p>Chemigation: Follow chemigation instructions in Product Use Information section.</p>		
<p>Flood (Basin) Irrigation: For flood (basin) irrigation systems, meter continuously into the water during the entire irrigation period. Best weed control results are obtained when a uniform distribution and flow of irrigation water is maintained over level land. Irrigation water treated with OXYFLUORFEN 2EC must be contained on the treated area until the water is absorbed by the soil.</p>		
<p>Low Volume Sprinkler (Microsprinkler) and Drip (Trickle) Irrigation: Apply only through low-volume sprinkler or drip systems designed to uniformly distribute irrigation water beneath the tree canopy. Applications should be made prior to weed emergence; otherwise postemergence activity may be inconsistent due to uneven coverage. Meter OXYFLUORFEN 2EC at a continuous rate during the middle 1/3 of the irrigation period and discontinue application during the final 1/3 of the irrigation period to ensure proper flushing of the irrigation system. Use of OXYFLUORFEN 2EC through low-volume sprinklers or drip emitters helps to reduce the "ring effect" of weed escapes in areas around sprinklers or emitters where previously applied broadcast or directed treatments begin to break down.</p>		
<p>Precautions:</p> <ul style="list-style-type: none"> • Direct spray toward the base of trees. Avoid direct contact with foliage or nuts. • OXYFLUORFEN 2EC should be applied only to healthy growing trees 		
<p>Crop-Specific Use Restrictions:</p> <ul style="list-style-type: none"> • When applied as a non-dormant treatment, OXYFLUORFEN 2EC can only be applied to pistachio plantings between May and 7 days prior to harvest. • When applied as a non-dormant treatment, OXYFLUORFEN 2EC can only be applied to almond plantings between April 1 and September 30 and to walnut plantings between May 1 and September 30. • Do not apply OXYFLUORFEN 2EC within 7 days of harvest of pistachios. • Do not apply OXYFLUORFEN 2EC within 30 days of harvest of almonds. • Do not apply OXYFLUORFEN 2EC within 7 days of harvest of walnuts. • Do not apply more than 6 pints of OXYFLUORFEN 2EC per acre during the non-dormant season. 		

Key Weeds Suppressed and/or Controlled

cheeseweed, malva fiddleneck, coast filaree, broadleaf filaree, redstem filaree, whitestem	morningglory species, annual mustard, black nettle, burning pigweed, redroot purslane, common
--	---

groundsel, common henbit minerslettuce	redmaids rocket, London sowthistle, annual
--	--

Additional Weeds Controlled in Tank Mix with Glyphosate or Paraquat

barnyardgrass bluegrass, annual chickweed, common	horseweed (marestail) rocket, London ryegrass, Italian
---	--

WINDBREAKS AND SHELTERBELTS
(For Use Only in Minnesota, North Dakota, South Dakota and Wyoming)

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence Postemergence	4 – 6	<p>Apply OXYFLUORFEN 2EC may be applied as a broadcast, banded or post-directed spray.</p> <p>Preemergence control is most effective when spray is applied to clean, weed-free soil surfaces. Pre-transplant applications must be made after completion of soil preparation but prior to transplanting. Transplanting should be completed with minimal soil disturbance. For optimum weed control results, treated soil surfaces should be left undisturbed during the time period for which weed control is desired.</p> <p>Postemergence Weed Control: For best results, apply before 4-leaf stage for broadleaf weeds or 2-leaf stage for grass weeds.</p> <p>Conifers: OXYFLUORFEN 2EC can be applied pre-transplant, post-directed or postemergence (over-the-top) to conifers. Postemergence or post-directed applications should be applied prior to budbreak or after new growth foliage has hardened off and new terminal buds have formed.</p> <p>Deciduous Hardwoods: OXYFLUORFEN 2EC has exhibited selectivity to many deciduous species when applied pre-transplant or as a post-directed spray prior to budbreak.</p>
<p>Precautions:</p> <ul style="list-style-type: none"> • Important: Some varieties or cultivars of conifers or deciduous species listed may be susceptible to OXYFLUORFEN 2EC. Care should be taken to ensure that the particular variety to be sprayed with OXYFLUORFEN 2EC is tolerant. For unfamiliar species, it is suggested that OXYFLUORFEN 2EC be tested on a limited number of plants prior to large-scale application. • Occasionally after the use of OXYFLUORFEN 2EC, a spotting, crinkling or flecking may appear on the leaves of the deciduous species. Leaves that receive direct or indirect (drift) spray contact will be injured. Deciduous species typically rapidly outgrow these symptoms and develop normally. • Application after budbreak may result in injury to deciduous species and is not recommended. If non-dormant application is required, apply only after foliage has fully expanded and hardened off. Avoid direct or indirect spray contact with the foliage by applying to the soil surface as a directed spray. • Apply OXYFLUORFEN 2EC only to healthy deciduous and/or conifer trees. Do not apply OXYFLUORFEN 2EC to conifers or deciduous trees that have been weakened or under stress from excessive fertilizer or soil salts, disease, nematodes, frost, drought, flooding, previously applied pesticides, soil insects, or winter injury, as severe injury may result. 		
<p>Specific Use Restrictions for Shelterbelts:</p> <ul style="list-style-type: none"> • Do not apply more than 6 pints of OXYFLUORFEN 2EC per acre in a single application or more than 18 pints per acre per year. 		

Key Broadleaf Weeds Controlled:

buckwheat, wild burclover carpetweed dock, curly groundcherry, cutleaf groundcherry, Wright groundsel, common henbit jimsonweed knotweed, prostrate kochia ladythumb lambsquarters, common lettuce, prickly mallow, little mayweed mustard, blue mustard, tumble	mustard, wild nettle, burning nightshade, black nightshade, hairy oats, wild orach, red pepperweed, yellow flower pigweed, prostrate pigweed, redroot purslane, common rocket, London shepherdspurse † smartweed, Pennsylvania sowthistle, annual tansymustard thistle, Russian (seedling) velvetleaf
---	---

† The highest rate or multiple applications may be required for acceptable control.

Key Grasses Controlled:

barnyardgrass bluegrass, annual crabgrass, large	foxtail, giant goosegrass witchgrass
--	--

OXYFLUORFEN 2EC may be applied to numerous conifer and deciduous species, including the following:

Conifer Species

Common Name	Scientific Name
douglas-fir	<i>Pseudotsuga menziesii</i>
fir	
grand	<i>Abies grandis</i>
fraser	<i>Abies fraseri</i>
noble	<i>Abies procera</i>
hemlock	
eastern hemlock	<i>Tsuga canadensis</i>
western hemlock	<i>Tsuga heterophylla</i>
pine	
Austrian	<i>Pinus nigra</i>
eastern white	<i>Pinus strobus</i>
jack	<i>Pinus banksiana</i>
Himalayan	<i>Pinus graffithii</i>
loblolly	<i>Pinus taeda</i>
lodgepole	<i>Pinus contorta</i>
longleaf	<i>Pinus palustris</i>
monterey	<i>Pinus radiata</i>
mugo	<i>Pinus mugo</i>
ponderosa	<i>Pinus ponderosa</i>

scotch shortleaf slash Virginia	<i>Pinus sylvestris</i> <i>Pinus echinata</i> <i>Pinus elliotii</i> <i>Pinus virginiana</i>
spruce blue dwarf Alberta Norway Sitka	<i>Picea pungens</i> <i>Picea glauca cornea</i> <i>Picea abies</i> <i>Picea sitchensis</i>
Arborvitae	<i>Thuja occidentalis</i> <i>Thuja orientalis</i>
juniper	<i>Juniperus chinensis</i> <i>Juniperus horizontalis</i> <i>Juniperus procumbens</i> <i>Juniperus sabina</i> <i>Juniperus scopulorum</i>
red cedar	<i>Juniperus virginiana</i>
yew	<i>Taxus spp.</i>

Deciduous Hardwood Species

Common Name	Scientific Name
ash	<i>Fraxinus spp.</i>
crabapple	<i>Malus spp.</i>
eucalyptus	<i>Eucalyptus spp.</i>
lilac	<i>Syringa vulgaris</i>
maple, black	<i>Acer nigrum</i>
oak, northern red	<i>Quercus rubra</i>
olive, Russian	<i>Elaeagnus angustifolia</i>
poplar (cottonwood)	<i>Populus spp.</i>
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
walnut, black	<i>Juglans nigra</i>

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Albaugh LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Albaugh MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Albaugh LLC or the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

To the extent consistent with applicable law, Albaugh takes no responsibility for unexpected results occurring due to the tank

mixing of this product with oils, surfactants, liquid fertilizers or other pesticides.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Albaugh's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Albaugh shall not be liable for losses or damages resulting from handling or use of this product unless Albaugh is promptly notified of such loss or damage in writing. To the extent consistent with applicable law, in no case shall Albaugh be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Albaugh or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.