

42750-154

1 / 13 / 2014

1/8



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

JAN 13 2014

Ms. Carolyn Miller
Albaugh, Inc.
P.O. Box 2127
Valdosta, GA 31604

Subject: Revised Label to Add Prohibition Statement Specific to the State of Oregon

Dear Ms. Miller:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated December 3, 2013 for:

EPA Registration 42750-154

Imidacloprid 0.2G

The Registration Division (RD) has conducted a review of this request of applicability under PRN 98-10 and finds that the label changes(s) requested falls within the scope of PRN 98-10. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, call me at 703 305-5409 or electronically at daniel.dani@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Dani Daniel".

Dani Daniel
Registration Division (7504P)
Insecticide/Rodenticide Branch

ALBAUGH, INC.

3/8
GEORGIA OFFICE
P.O. Box 2127
Valdosta, GA 31604-2127
229.244.3288 (Phone)
229.244.5841 (FAX)

Fed-Ex

December 3, 2013

Document Processing Desk (NOTIFY)
Ms. Venus Eagle (Team 1)
Registration Division
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard (South Bldg)
2777 South Crystal Dr.
Arlington, VA 22202

Re: Imidacloprid 0.2G
EPA Reg. No. 42750-154

Dear Ms. Eagle,

We are submitting this notification for the above referenced product under PR 98-10 to add a use prohibition statement specific to the state of Oregon.

~~One~~ ^{is} hard copies ~~are~~ enclosed, along with EPA Form 8570-1. The notification statement is on our form.

Should you have any questions, please do not hesitate to call me or e-mail. Thank you and have a great day.

Sincerely,


Carolyn Miter
Registrations Specialist
Albaugh, Inc.
P. O. Box 2127
Valdosta, GA 31604
229-244-3288
carolynm@albaughinc.com

Agri Star[®]
By Albaugh, Inc.

PREMIER SUPPLIER OF OFF-PATENT CROP PROTECTION PRODUCTS
www.albaughinc.com

4/8

NOTIFICATION

JAN 13 2014

EDITOR's NOTE: 12/4/13 label notification to add Oregon pollinator statement

IMIDACLOPRID 0.2G Lawn and Garden Insecticide

[Optional marketing statements]

Season Long grub control

Use anytime...spring through late summer

For residential use on lawns and landscaped areas.

Kills white grubs and mole crickets early, before they damage your lawn.

Apply before insects hatch to prevent turf damage.

One application controls grubs all season long.

One application stops grubs all season long.

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine 0.2%

OTHER INGREDIENTS: 99.8%

TOTAL: 100.0%

STOP - Read the label before use.

FOR OUTDOOR RESIDENTIAL USE ONLY

Not for sale, sale into, distribution and or use in Nassau, Suffolk, Kings and Queens Counties of New York State.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration. • Call a poison control center or doctor for further treatment advice.
In case of emergency call CHEMTREC toll free at 1-800-424-9300. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
Note To Physician: No specific antidote is available. Treat the patient symptomatically.	

EPA Reg. No. 42750-154

EPA Est. No. xxxxxx-xxxx

AD120413

NET CONTENTS: _____ Gallons

Manufactured For:
ALBAUGH, INC.
ANKENY, IA 50021

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

WHAT IMIDACLOPRID 0.2 G LAWN AND GARDEN INSECTICIDE DOES

White grubs can cause extensive damage to turf by feeding on grass roots. Affected turf turns yellow, then brown, and is easily pulled up in pieces with your hands. Mole crickets can seriously damage lawns by their tunneling activity. Some species also feed on the grass roots, as well. IMIDACLOPRID 0.2 G prevents turf damage by killing grubs and mole crickets when they are newly hatched and before they can cause turf damage.

IMIDACLOPRID 0.2 G works best when applied before grubs and mole crickets hatch. The active ingredient will remain in the root zone of the turf until the eggs of grubs or mole crickets hatch. These newly hatched insects are highly susceptible to IMIDACLOPRID, and are killed before they become large enough to cause turf damage. IMIDACLOPRID 0.2 G also kills grubs in landscaped ornamental areas containing flowers, ground covers, shrubs and trees.

If you have questions regarding the best time to make applications in your area, you can consult your local County University Extension Office. Be sure that you describe the product you are using when seeking such advice.

HOW TO APPLY IMIDACLOPRID 0.2 G LAWN AND GARDEN INSECTICIDE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For residential use on lawns and landscaped areas.

Do not apply this product by any application method, to linden, basswood or other *Tilia* species in the State of Oregon.

READ ALL DIRECTIONS FOR USE AND CONDITIONS OF SALE BEFORE USING

APPLICATION TO LAWNS:

Pests Controlled:

White grubs (including the larvae of Japanese beetle, Oriental beetle, Chafers, Asiatic garden beetle, May or June beetles, and black turfgrass atenius), European Crane fly (larvae) and mole crickets.

How to Apply:

Using either a drop-type or rotary-type spreader, apply IMIDACLOPRID 0.2 G uniformly over the area being treated. Avoid the use of spreaders which may apply the product in narrow, concentrated or uneven bands. Calibrate your spreader according to the spreader manufacturer's directions to ensure the proper amount of IMIDACLOPRID 0.2 G is applied. Check frequently to be sure spreader is working properly and spreading granules evenly.

6/8

After Applying:

IMIDACLOPRID 0.2 G will work best if the lawn is watered thoroughly or if rainfall occurs within 24 hours after application to move the active ingredient through the thatch and down into the soil. Avoid mowing lawn until watering or rainfall has occurred so that the uniformity of the granules is not disturbed.

When to Apply:

IMIDACLOPRID 0.2 G works best when applied before grubs and mole crickets hatch.

For best results, apply as follows:

White grubs: Apply once, anytime from April through mid-August.

Mole crickets: In Florida, apply once from early April through early May. Elsewhere, apply once from mid-May through mid-June.

Do not Apply IMIDACLOPRID 0.2 G when turf areas are waterlogged. The product may not work effectively under these conditions.

How Much to Apply:

For control of white grubs, IMIDACLOPRID 0.2 G should be applied at the rate of 2.87 pounds per 1,000 square feet. For mole cricket control, apply 3.8 pounds per 1000 square feet. The table below can be used as a guide for applying the proper rate of IMIDACLOPRID 0.2 G. However, the application rate can vary depending on the speed of application and the condition of the spreader. For best results, use a normal, steady pace when applying to obtain an accurate, even application.

SPREADER SETTING TABLE

Spreader Model	Setting*	
	To Deliver 2.87 lbs per 1,000 sq ft	To Deliver 3.8 lbs per 1,000 sq ft
Vigoro Drop	Spreader Settings to be determined	Spreader Settings to be determined
Vigoro Rotary		
Sta-Green Drop		
Sta-Green Rotary		
Bandini		
Ev-N-Spread Rotary		
Ortho Drop		
Ortho Rotary		
Precision Drop		
Precision Rotary		
Quaker Rotary		
Republic EZ Drop		
Republic EZ Grow		
Republic EZ Rotary		
Scotts Drop		
Scotts Accugreen		
Scotts Easygreen		
Scotts Evengreen		
Scotts Speedy Green		
True Temper Rotary		

Settings are approximate. The condition of your spreader, the speed at which you walk and uneven

terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.

Tip: 1,000 sq ft = 20 ft x 50 ft

Do not use a spreader that applies granules in narrow, concentrated, or uneven bands. Be sure the spreader works properly and spreads granules evenly.

APPLICATION TO LANDSCAPE PLANTINGS
(Trees, Shrubs, Evergreens, Flowers, Ground covers)

Pests Controlled:

White grubs (including larvae of Japanese beetle, Oriental beetle, chafers, Asiatic garden beetle, European Cranefly, May and June beetle, and black turfgrass atatenius).

How to Apply:

Apply as a broadcast treatment to soil, mulch or other substrate. Prior to application, remove black plastic from area to be treated to allow water to penetrate into the soil. Permeable weed barriers need not be removed prior to application.

After Applying:

IMIDACLOPRID 0.2 G will work best if material is worked into the soil around plants or if a thorough watering of rainfall occurs within 24 hours after application to move the product down through mulch and into the soil.

When to Apply:

The best control of grubs will be obtained if IMIDACLOPRID 0.2 G is applied before grubs begin to hatch. For best results, application should occur from April through mid-August.

DO NOT apply to waterlogged areas. IMIDACLOPRID may not work effectively under these conditions.

How Much to Apply:

IMIDACLOPRID 0.2 G should be applied at the rate of 2.87 pounds per 1,000 square feet. This 15 pound bag will treat 5,000 square feet of landscaped area. The table above can be used as a guide for applying the proper rate of IMIDACLOPRID 0.2 G. However, the application rate can vary depending on the speed of application and the condition of the spreader.

NOTE: DO NOT apply IMIDACLOPRID 0.2 G to vegetable gardens.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

Storage Conditions: Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.

Disposal

If empty: Do not reuse container. Place in trash or offer for recycling if available.

If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Paper and plastic bags: Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

