



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

May 21, 2025

Tasha Lott
Product Registration Manager
Albaugh, LLC
1525 NE 36th St.
Ankeny, IA 50021

Subject: Label Amendment - Registration Review Mitigation for Bifenthrin
Product Name: IMID-BIFEN 0.36 GC INSECTICIDE
EPA Registration Number: 42750-161
Application Date: February 19, 2021
Decision Number: 597031

Dear Tasha Lott:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Bifenthrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must

submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Meaghan Robertson by phone at 202-566-0169, or via email at robertson.meaghan@epa.gov.

Sincerely,

A handwritten signature in blue ink that reads "Cathryn Britton". The signature is written in a cursive, flowing style.

Cathryn Britton
Chief, Risk Management and Implementation
Branch V
Pesticide Re-evaluation Division (7508M)
Office of Pesticide Programs

ENCLOSURE: Stamped label

Bifenthrin	Group	3A	Insecticide
Imidacloprid	Group	4A	Insecticide

RESTRICTED USE PESTICIDE
Due To Toxicity To Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

IMID-BIFEN 0.36 GC
Insecticide

For general insect control in turfgrass areas of golf courses and sod farms

Do Not Use This Product On Golf Courses And Sod Farms In Nassau County Or Suffolk County, New York

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	0.20%
Bifenthrin*	0.16%
OTHER INGREDIENTS:	99.64%
TOTAL:	100.00%

*Cis isomers 97% minimum, trans isomers 3% maximum.

STOP - Read the label before use.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center. • Do not give anything by mouth to an unconscious person.
IF ON SKIN	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of emergency call CHEMTREC toll free at 1-800-424-9300.	
Note To Physician: No specific antidote is available. Treat the patient symptomatically. This product contains a pyrethroid. If large amounts have been ingested, milk, cream and other digestible fats and oils may increase absorption and so should be avoided.	


EPA Reg. No. 42750-161

EPA Est. No. xxxxxx-xx-xxx

Net Weight: 50 Lbs.

Manufactured For:
Albaugh, LLC
Ankeny, IA 50021

042750-00161.20250516.DRAFT



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long sleeve shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber (≥ 14 mils), nitrile (≥ 14 mils), neoprene rubber (≥ 14 mils), natural rubber (≥ 14 mils), polyethylene, polyvinyl chloride (PVC) (≥ 14 mils), or viton (≥ 14 mils).
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATION

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems. Run-off may be hazardous to aquatic organisms in water adjacent to treated areas.

This product contains a chemical with properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For outdoor use only.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your

State or Tribe, consult the agency responsible for pesticide regulation.

Do not formulate this product into other end-use products.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not apply or irrigate to the point of run-off.

Do not make applications during rain.

For applications at rates above 0.34 lbs bifenthrin (212.5 lbs product/A), users must water the treated area immediately after application.

For soil or foliar applications, do not apply by ground within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes or ponds, estuaries, and commercial fish farm ponds).

Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Do not apply when the wind speed is greater than 15 mph.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.)

VEGETATIVE FILTER STRIPS

Construct and maintain a vegetative filter strip, according to the width specified below, of grass or other permanent vegetation between the field edge and nearby down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing bifenthrin onto fields where a maintained vegetative filter strip of at least 25 feet exists between the field edge and where a down gradient aquatic habitat exists. This minimum required width of 25 feet may be reduced or removed under the following conditions:

- For Western irrigated agriculture, a maintained vegetative filter strip of at least 10 feet wide is required. Western irrigated agriculture is defined as irrigated farmland in the following states: WA, OR, CA, ID, NV, UT, AZ, MT, WY, CO, NM, and TX (west of I-35).
 - For Western irrigated agriculture, if a sediment control basin is present, a vegetative filter strip is not required.
- In all other areas, a vegetative filter strip with a minimum width of 25 feet is required, unless the following conditions are met. The vegetative filter strip requirement may be reduced from 25 feet to 15 feet if at least one of the following applies:
 - The area of application is considered prime farmland (as defined in 7 CFR § 657.5).
 - Conservation tillage is being implemented on the area of application. Conservation tillage is defined as any system that leaves at least 30% of the soil surface covered by residue after planting. Conservation tillage practices can include mulch-till, no-till, or strip-till.
 - A functional terrace system is maintained on the area of application.
 - Water and sediment control basins for the area of application are functional and maintained.

- The area of application is less than or equal to 10 acres.

For further guidance on vegetated filter strips, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. <https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0331-0175>.

RESISTANCE MANAGEMENT

For resistance management, please note that IMID-BIFEN 0.15 GC + FERTILIZER contains both a Group 3A and a Group 4A insecticide. Any insect/mite population may contain individuals naturally resistant to IMID-BIFEN 0.15 GC + FERTILIZER and other Group 3A or Group 4A insecticides. The resistant individuals may dominate the insect/mite population if these insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of IMID-BIFEN 0.15 GC + FERTILIZER or other Group 3A or Group 4A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, consult an Albaugh LLC representative at 1-800-247-8013.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance, it also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that Involves contact with anything that has been treated, such as plants, soil, or water, is

1. Coveralls
2. Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber (≥ 14 mils), nitrile (≥ 14 mils), neoprene rubber (≥ 14 mils), natural rubber (≥ 14 mils), polyethylene, polyvinyl chloride (PVC) (≥ 14 mils), or viton (≥ 14 mils).
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENT

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides {40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

PRODUCT INFORMATION

APPLICATION TO TURFGRASS

The use of this product will provide control of selected insect pests. IMID-BIFEN 0.36 GC can be used for the control of the following soil inhabiting pests of turfgrass:

COMMON NAME	SCIENTIFIC NAME
Northern and Southern masked chafers	<i>Cyclocephala borealis</i> , <i>C. immaculate</i> , and/or <i>C. lurida</i>
Asiatic garden beetle	<i>Maladera castanea</i>
black vine weevil	<i>Otiorhyncus sulcatus</i>
European chafer	<i>Rhizotroqus majalis</i>
green June beetle	<i>Cotinis nitida</i>
May or June beetles	<i>Phyllophaga spp.</i>
Japanese beetle	<i>Popillia japonica</i>
oriental beetle	<i>Anomala orientalis</i> , <i>Euetheola</i> , <i>humulis rugiceps</i>
billbugs	<i>Sphenophorus spp.</i>
annual bluegrass weevil	<i>Lissonotus Maculicollis</i>
black turfgrass ataenius	<i>Ataenius spretulus</i> ; <i>Aphodius spp.</i>
European crane fly	<i>Tipula paludosa</i>

frit fly	<i>Oscinella frit</i>
chinch bugs	<i>Blissus spp.</i>
fire ants	<i>Solenopsis spp.</i>
cutworms	<i>Agrotis ipsilon, Peridroma saucia, Nephelodes menians</i>
armyworms	<i>Spodoptera spp., Pseudaletia spp.</i>
sod webworm	<i>Crambus spp.</i>
mole crickets	<i>Scapteriscus spp.</i>

IMID-BIFEN 0.36 GC can be used as directed on turfgrass on golf courses and sod farms.

The active ingredients IMID-BIFEN 0.36 GC have sufficient residual activity so that applications for control of subsurface feeders can be made preceding the egg laying activity. The need for an application for control of subsurface feeders can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control of subsurface feeders will be achieved when applications are made prior to egg hatch of the target pest, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

The active ingredients in IMID-BIFEN 0.36 GC have sufficient knockdown and residual activity to provide curative and residual control of surface feeding pests. Applications for control of surface feeding pests can be made when infestations are anticipated based on historical monitoring of the site, previous records or experience, current season adult trapping or presence of insects at economic thresholds as determined by scouting and/or recommendations of local State extension personnel or other qualified specialists.

Applications should not be made when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Total amount applied cannot exceed 250 lb (0.5 lb of imidacloprid, 0.4 lb of bifenthrin) per acre per year.

NOTE: Not for use in commercial greenhouses or on grass grown for seed.

APPLICATION EQUIPMENT

Apply IMID-BIFEN 0.36 GC uniformly over the area being treated with normally used granular application equipment. Both drop-type and rotary-type spreaders may be used to apply IMID-BIFEN 0.36 GC. Avoid the use of spreaders that apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

RECOMMENDED APPLICATIONS

(Be sure to read "APPLICATION EQUIPMENT" Section of this label)

CROP	PEST	DOSAGE IMID-BIFEN 0.36 GC
------	------	------------------------------

Turfgrasses	Larvae of subsurface feeders: Asiatic garden beetle Black turfgrass ataenius Black vine weevil European chafer European crane fly Green June beetle Japanese beetle Northern masked chafer Oriental beetle Phyllophaga spp. Southern masked chafer	100 to 125 lb per acre (2.3 to 2.9 lb per 1,000 ft sq)
	Adults and/or larvae of subsurface and surface feeding pests: Centipedes Fleas Millipedes Ticks ¹⁰ Imported fire ants ⁶ Mole crickets ^{7,8} Pillbugs ⁹ Sowbugs ⁹	75 to 125 lb per acre (1.7 to 2.9 lb per 1,000 ft sq) or for heavy infestations, a single annual application of 200 to 250 lb per acre (4.6 to 5.7 lb per 1000 sq ft)
	Adults and/or larvae of surface feeders: Annual bluegrass weevil ^{1,2} Billbugs ⁴ Chinch bugs ⁵ Frit fly Leafhoppers Mealybugs Nuisance ants	75 to 125 lb per acre (1.7 to 2.9 lb per 1,000 ft sq)
	Larvae of surface feeders: Armyworm ³ Cutworm ³ Sod webworms ³	50 to 125 lb per acre (1.1 to 2.9 lb per 1,000 ft sq)

¹Annual Bluegrass Weevil, adults: Applications should be timed to control adult weevils as they leave their overwintering sites and move into grass areas. This movement generally begins when Forsythia is in full bloom and concludes when flowering dogwood (*Cornus florida*) is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

²Annual Bluegrass Weevil, larvae: For best results, applications should be made at the first sign of wilting of bluegrass. For the first generation, this wilting often occurs after full bloom of flowering dogwood (*Cornus florida*).

³Armyworms, Cutworms and Sod Webworms: If the grass is being maintained at a mowing height of greater than 1 inch, then higher application rates (up to 2.9 lb per 1000 sq ft) may be required during periods of high pest pressure.

⁴Billbug, adults: Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting billbug adults will also provide control of over-wintered chinch bugs.

⁵Chinch Bugs: Chinch bugs infest the base of grass plants and are often found in the thatch layer. Irrigation of the grass area before treatment will optimize the penetration of the insecticide to the area where the chinch bugs are located. Chinch bugs can be one of the populations that contain both nymphs and adults during the middle of the summer.

⁶Imported Fire ants: IMID-BIFEN 0.36 GC may be applied as a broadcast treatment up to 5.7 lb. per 1,000 sq. ft. For a mound treatment, apply 1/2 cup of IMID-BIFEN 0.36 GC per mound and then drench the mound with 1 to 2 gallons of water. The mounds should be treated with sufficient force to break their apex and allow the insecticide to flow into the ant tunnels.

⁷Mole Crickets, adults: Achieving acceptable control of adult mole crickets is difficult because grass areas are subject to continuous invasion during the early spring by this large active stage. For application when both adults and/or large nymphs are present and actively tunneling, irrigate prior to application if soil is dry and apply 2.9 lbs per 1000 sq. ft. Applications should be watered in with up to 0.5 inches of water immediately after treatment. Grass areas that receive pressure from adult mole crickets should be treated within one month of peak egg hatch to ensure optimum control of subsequent nymph populations (see below). For heavily infested mole cricket areas, or in areas with a history of mole cricket infestation, use a single seasonal application rate of 200 to 250 lb per acre (4.6-5.7 lb per 1000 sq ft).

⁸Mole Crickets, nymphs: Grass areas that received adult mole cricket pressure in the spring should be treated within one month of peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides. Control of larger, more damaging, nymphs later in the year will require higher application rates to maintain acceptable control. For maximum effectiveness, irrigate prior to application if soil is dry, apply as a broadcast treatment at 2.9 lb per 1000 sq. ft. Applications should be watered in with up to 0.5 inches of water after treatment. For heavily infest mole cricket areas with a history of mole cricket infestation, use a single seasonal application rate of 200 to 250 lb per acre (4.6-5.7 lb per 1000 sq ft).

⁹Pillbugs and sowbugs: For heavy infestation use 250 lbs per acre (5.7 lbs per 1,000 sq. ft.) of IMID-BIFEN 0.36 GC.

¹⁰Ticks (including ticks that may transmit Lyme disease and Rocky Mountain spotted fever): Treat the entire area where exposure to ticks may occur. Ticks may be reintroduced from surrounding areas on host animals. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed activity. Repeat application should be limited to no more than once per seven days. Deer ticks (*Ixodes* spp.) have a complicated life cycle that range over a two year period and involves four life stages. Applications should be made in the late fall and/or early spring to control adult ticks that are usually located on brush or grass above the soil surface and in mid to late spring to control larvae and nymphs that reside in the soil and leaf litter. American dog ticks (*Dermacentor variabilis*) may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These ticks commonly congregate along paths or roadways where humans are likely to be encountered. Applications should be made as necessary from mid-spring to early fall to control American dog tick larvae, nymphs and adults.

Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for more specific information regarding timing of application.

NOTE: For optimum control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. Do not apply more than 250 lb product per acre (0.5 lb of imidacloprid active ingredient, 0.4 lb bifenthrin) per acre per year. Avoid mowing turf or lawn area until after irrigation or rainfall has occurred so that uniformity of application will not be affected.

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of IMID-BIFEN 0.36 GC if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

SPREADER SETTINGS FOR IMID-BIFEN 0.36 GC

The following chart can be used as a guide in establishing an accurate rate of IMID-BIFEN 0.36 GC. Application rates will vary depending upon speed, equipment wear and condition of spreader. Carefully calibrate application equipment prior to commercial use of this product.

SPREADER	SPEED (mph)	SETTINGS	
		50 lb/A	100 lb/A
Cyclone Way	3	1.53	4
Lely	4.5	3 ¾	4 ¼
Lesco Broadcast	3	C	D
PSB Prizelawn CBR-III	3	3	3 ½
Scott's Proturf R-8A	3	H	J
Scott's ProturfSR-2000	3	I	K
Scott's Handy Green (obsolete)	3	2	3
Scott's Accupro	3	4 ½	4 ¾
Spyker	3	3.0	3.4
Vicon Model 402	4.5	11	14

RESTRICTIONS

- Avoid runoff or puddling of irrigation water following application.
- Avoid application of IMID-BIFEN 0.36 GC to areas which are waterlogged or saturated, which will not allow penetration into the root zone of the plant.
- Do not apply more than 250 lb (0.5 lb of imidacloprid and 0.4 lb of bifenthrin) per acre per year.
- Treated areas may be replanted with any crop specified on an imidacloprid and a bifenthrin label, or with any crop for which a tolerance exists for the two active ingredients.
- For crops not listed on any imidacloprid and a bifenthrin label, or for crops for which no tolerances for the active ingredients have been established, a 12-month plant-back interval should be observed.
- Do not apply by air.
- Not for use in nurseries or commercial greenhouses.
- Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If material is spilled for any reason or cause, carefully contain any spilled material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides above. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill, by incineration, or if allowed by the state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Albaugh's control it is impossible for Albaugh to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALBAUGH MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Albaugh is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Albaugh disclaims

any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ALBAUGH'S ELECTION, THE REPLACEMENT OF PRODUCT.

051625

LABEL HISTORY

Not Part of Final printed Label

File Name	Version Mark	Comment
042750-00161.20210215.DRAFT	021521	EPA Reg Review
042750-00161.20250507.DRAFT	050725	(e) Label Revisions
042750-00161.20250516.DRAFT	051625	(e) Label Revisions