



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

January 2, 2007

Mr. Morris Gaskins
Albaugh, Inc.
P.O. Box 2127
304 Janet Street, Suite H
Valdosta, GA 31604

Subject: Label Notifications for Pesticide Registration Notice 2007-4

Dear Mr. Gaskins,

The Agency is in receipt of your Applications for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated November 28, 2007 for the following products:

EPA Registration Number 42750-130	Tebuconazole 28% FL ST
EPA Registration Number 42750-167	NUCOP 3L HB
EPA Registration Number 42750-29	Weed Hoe 108
EPA Registration Number 42750-28	Weed Hoe 120
EPA Registration Number 42750-79	Picloram K-Salt Rangeland
EPA Registration Number 42750-81	Picloram K-Salt IVM


The Registration Division (RD) has conducted reviews of these requests for their applicability under PRN 2007-4 and finds that the actions requested fall within the scope of PRN 2007-4. The labels submitted with the applications have been stamped "Notification" and will be placed in our records.

With regard to your proposed label for EPA Reg. No. 42750-167, NUCOP 3L HB, it appears that in addition to revising the label to reflect the language in PRN 2007-4, you are also adding a new disease, phomopsis, to the label for use on grapes (p. 18 of your proposed label). This type of action can be handled through the notification process, but in the future please include this as a separate notification. Also, please be aware that you must have data available showing efficacy against this disease.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code to identify the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification, per PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please call me directly at 703-305-6249 or Steve Schaible of my staff at 703-308-9362.

Sincerely,

for 

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

3/28

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 42750-167	2. EPA Product Manager T. Kish	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) NU-COP 3L HB	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Albaugh Inc. P.O. Box 2127 Valdosta, GA 31604-2127 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

JAN - 2 2008

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 2.5, 30, 250 bulk		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Morris Gaskins	Title Registrations Manager	Telephone No. (Include Area Code) 229-244-3288
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date application Received (Stamped)
2. Signature 	3. Title Registrations Manager	
4. Typed Name Morris Gaskins	5. Date November 28, 2007	

4/38

NUCOP 3L HB

ACTIVE INGREDIENT:	
Copper hydroxide	37.5%
OTHER INGREDIENTS:	62.5%
TOTAL:	100.0%

CAS # 20427-59-2
(Metallic copper equivalent 24.4%)
This product contains 3 lbs. of metallic copper per gallon

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. See Inside Booklet for Additional Precautionary Statements.	

EPA Reg. No. 42750-167

EPA Est. No. 45002-MEX-2

NET CONTENTS:

NOTIFICATION

JAN - 2 2008

MANUFACTURED FOR:

Albaugh, Inc.
1525 NE 36th Street
Ankeny, IA 50021

5/38

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING – AVISO

Harmful if swallowed, absorbed through the skin, or inhaled. Causes substantial but temporary eye injury. Do not get in eyes, on skin, or on clothing. Wear goggles, face shield or safety glasses. May cause skin sensitization in certain individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Mixers, loaders, applicators and other handlers must wear the following:

- 1. Long-sleeved shirt and long pants
- 2. Chemical resistant gloves made of any waterproof material
- 3. Shoes plus socks
- 4. Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This product may contaminate water through runoff. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours provide the following instructions are followed. For at least 7 days following application, an eye-flush container, designed specifically for flushing eyes, must be available at the WPS decontamination site for workers entering the area treated with copper hydroxide.

Notify workers of the application by warning them orally that residues in the treated areas may be highly irritating to their eyes and to take precautions such as refraining from rubbing their eyes and if they get residues in their eyes they should immediately flush their eyes using the eye-flush.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep in a cool place. Do not store at temperatures below 32°F. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

~~CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.~~

CONTAINER DISPOSAL:

Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable <5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼

full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non refillable >5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container (250 gallon & bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinseate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

GENERAL INSTRUCTIONS

Use NuCop 3L HB as noted below. NuCop 3L HB is adaptable to spraying from all types of spray equipment. Depending on the equipment used and the specific crop, the volume applied per acre will differ. For dilute, high volume sprays use from 25 to 100 gallons of water per acre for most vegetable crops, 400 to 800 GPA for fruit and nut crops, and up to 1500 gallons per acre as may be required for large citrus groves. For aerial spraying, 3 to 15 GPA are commonly used. For concentrate ground sprays, apply from 5 to 20 gallons per acre for vegetable crops and 25 to 100 gallons per acre for fruit and nut crops. Add NuCop 3L HB slowly to spray tank partially filled with water. Spreader-stickers, insecticides, nutrients, etc. should be added last. NuCop 3L HB is compatible with commercially formulated spreader-stickers, oils and such insecticides as Carbaryl and other fungicides. Observe all cautions and limitations on label of all products used in mixtures.

The following specific instructions are based on general applications. The recommendations of the State Agricultural Extension Services should be closely followed as to timing, frequency, and number of sprays per season. When a range of doses are given for the use site, use the low dose when conditions are not favorable for disease development and use the high dose when conditions are favorable for disease development. Consult your State Agricultural Extension Service for guidance in determining what conditions favor diseases for the particular use site.

GENERAL CHEMIGATION INSTRUCTIONS

Do not apply this product through any irrigation system using aluminum parts or components as damage to the system may occur. Such application is prohibited regardless of whether the irrigation system is flushed with water after use of this product.

Apply this product only through one or more of the following types of systems: sprinkler including center

pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation system(s) which contain no aluminum parts or components. Do not apply this product through any other type of irrigation system.

Crop injury or lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

A. Center Pivot, Traveler, Big Gun, Motorized Lateral Move, End Tow, and Side (Wheel) Roll Irrigation Equipment: Operate system and injection equipment at normal pressures recommended by the manufacturer of injection equipment used. Fill tank or injection equipment with water. Operate system for one complete circle for center pivot or one complete run for the other recommended equipment, measuring time required, amount of water injected, and acreage contained in circle or run. Mix recommended amount of product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run, but continue to operate irrigation system until the product has been cleared from last sprinkler head. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur.

B. Solid Set and Hand Move Irrigation Equipment: Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of product for acreage to be covered into quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Provide constant mechanical agitation in the mix tank to insure that the product will remain in suspension during the injection cycle. This product can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until pesticide is cleared from last sprinkler head.

SAFETY DEVICES

- (1) The systems designated above must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- (2) All pesticide injection pipelines must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- (3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- (4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- (5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The system must functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

For additional instructions on safety precautions refer to statements (2), (3), (4), (6), and (7) in the section on SAFETY DEVICES.

RECOMMENDATIONS

FROST INJURY PROTECTION:

Bacterial Ice nucleation inhibitor - Application of NuCop 3L HB made to all crops listed on this label at rates and stages of growth indicated on this label at least 24 hours and not more than 72 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola*, and *Pseudomonas fluorescens*) and may thereby provide some protection against light frost. The degree of frost protection will vary with weather conditions and other factors. Not recommended for those geographical areas where weather conditions favor severe frost.

ALFALFA			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM RETREATMENT INTERVAL	COMMENT
Cercospora & Leptosphaerulina Leaf Spots	2/3 to 1 1/3	30 Days	Apply 10 to 14 days before each harvest or earlier if disease threatens. Apply with ground or aerial equipment. Spray injury may occur with sensitive varieties such as Lahontan.
RESTRICTIONS: Maximum single application rate is 1 1/3 pt/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 3.0 pt/A (1.12 lbs metallic copper equivalent)			

APPLES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Fall & Late Dormant: Anthracnose European Canker Pseudomonas Syringae	4 – 21 1/3	Only one dormant application allowed per season	Apply before fall rains. Use on yellow varieties may cause discoloration. To avoid, pick before spraying.
Bloom & Growing Season: Fireblight	2/3 - 1 1/3	5	Make application between silver-tip and green-tip. ATTENTION: Phytotoxicity may occur from late application (Discontinue use when green-tip is 1/2 inch.)
Crown or Collar Rot (Phytophthora cactorum)	N/A	5	Apply either in early spring or in fall after harvest each year. Do not use if soil pH is below 5.5 or copper toxicity may result. Mix 1 1/3 pint in 100 gallon of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree.
RESTRICTIONS Maximum single dormant season application rate is 21 1/3 pt/A (8.0 lbs. metallic copper equivalent) Maximum single growing season application rate is 1 1/3 pt/A (0.5 lbs. metallic copper equivalent) Maximum annual application rate is 42 2/3 pt/A (16 lbs. metallic copper equivalent)			

ALMONDS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bloom/Growing Season: Coryneum Blight Blossom Brown Rot	2/3 - 4	5	Use during the early bloom stage (popcorn). A second application in late dormant before foliage buds swell may be necessary when frequent rainfall occurs. To avoid plant injury, do not use above rate after full bloom.

ALMONDS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Dormant to Pink Bud Season: Bacterial Blast (Pseudomonas)	2/3 - 8	7	Use at dormant to early pink bud. For blast control in sprinkler irrigated orchards or where disease is severe, apply 2-4 sprays or as many as required at 2/3 to 2 pints per acre at 2 week post-bloom intervals or just before sprinkling. Slight leaf injury may occur from post-bloom spray.
RESTRICTIONS Maximum single dormant application rate is 8 pt/A (3 lbs. metallic copper equivalent) Maximum single bloom/growing application rate is 4 pt/A (1.5 lbs. metallic copper equivalent) Maximum annual application rate is 48 pt/A (18 lbs metallic copper equivalent)			

APRICOTS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Coryneum Blight (Shot Hole) Blossom Brown Rot	2/3 - 4	5	Apply at popcorn to full bloom as a full cover spray. To avoid spray injury, do not apply after bloom.
RESTRICTIONS Maximum single application rate is 4 pt/A (1.5 lbs metallic copper equivalent) Maximum annual application rate is 42 2/3 pt/A (16 lbs metallic copper equivalent)			

ATEMOYA			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracoese	2 - 4	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
RESTRICTIONS Maximum single application rate is 4 pt/A (1.5 lbs metallic copper equivalent) Maximum annual application rate is 33 2/3 pt/A (12.6 lbs metallic copper equivalent)			

AVOCADOS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Scab	2/3 - 2 2/3 pints	14	Apply when bloom buds begin to swell. Continue application at monthly intervals for 5 to 6 applications. Follow recommendations of State Agricultural Experiment Stations.
RESTRICTIONS Maximum single application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 26 2/3 pt/A (10.0 lbs metallic copper equivalent)			

BANANAS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Sigatoka	2/3 - 2 2/3	7	Apply on a 14 day schedule throughout the wet season. Apply at 21 day intervals during dry periods.
Black Pitting	1 1/3 - 2 2/3	7	Apply directly to the fruit stem and include the basal portion of the leaf crown. Apply during the first and second weeks after emergence.
RESTRICTIONS Maximum single application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 50 1/3 pt/A (18.9 lbs metallic copper equivalent)			

BEANS (Dry, Green)			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bacterial Blight (Halo & Common)	1/3 - 2	7	For protective sprays apply first application when plants are six inches high. Apply on 7 to 14 day schedule depending on local conditions.
RESTRICTIONS Maximum single application rate is 2 pt/A (0.7 lbs. metallic copper equivalent) Maximum annual application rate is 12 2/3 pt/A (4.74 lbs metallic copper equivalent)			

BLACKBERRIES (Santiams, Logans, Boysens, Marions, Auroras, Cascades, Chehalems, & Thornless Evergreens)			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Leaf & Cane Spot	1 1/3 – 5 1/3	7	Apply delayed dormant spray after training in spring. Apply again in late spring. Make fall spray application after harvest.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 26 2/3 pt/A (10.0 lbs metallic copper equivalent)			

BLUEBERRIES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bacterial Canker	1 1/3 – 5 1/3	7	Make first application before the fall rains, preferably the first week in October and a second application 4 weeks later.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 22 pt/A (8.4 lbs metallic copper equivalent)			

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARD GREENS, MUSTARD GREENS, & TURNIP GREENS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew	1/3 – 1 1/3	7	Apply in a minimum of 25 GPA at 7 day intervals. (CAUTION: A slight reddening of older leaves may occur on broccoli, and a slight flecking of wrapper leaves may occur on cabbage.)
Black Rot (Xanthomonas)	1/3	7	
Black Leaf Spot (Alternaria)	2/3 – 1 1/3		
RESTRICTIONS Maximum single application rate is 1 1/3 pt/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 14 pt/A (5.25 lbs metallic copper equivalent)			

CACAO			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Black Pod	2/3 - 6	14	<p>Begin applications at the start of the rainy season and continue while infection conditions persist.</p> <p>Sprays should be made as often as 14 to 21 days in high rainfall areas at varying rates per acre depending on disease severity.</p> <p>For drier areas where 2 to 4 applications are recommended during critical infection periods and at long intervals, use 2 - 6 pints per acre, according to disease incidence and planting density.</p>
RESTRICTIONS Maximum single application rate is 6 pt/A (2.25 lbs metallic copper equivalent) Maximum annual application rate is 42 pt/A (15.75 lbs metallic copper equivalent)			

CARAMBOLA			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	4 - 5 1/3	7	<p>Make initial application just before flowering and repeat on a weekly schedule until just before harvest.</p> <p>Apply in sufficient water for thorough coverage.</p>
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 28 pt/A (10.5 lbs metallic copper equivalent)			

CARROTS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Carrot Blight (Cercospora)	2/3 - 2 2/3	7	When Disease threatens, apply at 7 to 14 day intervals.
RESTRICTIONS Maximum single application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 13 1/3 pt/A (5.0 lbs metallic copper equivalent)			

CELERY & CELERIAC			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Early, Late & Bacterial Blights	2/3 - 2 2/3	7	Apply as soon as plants are first established in the field, then every 7 days depending on severity and weather.
RESTRICTIONS Maximum single application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 14 pt/A (5.25 lbs metallic copper equivalent)			

CHERRY			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Dormant & Late Bloom Season: Dead Bud (Pseudomonas syringae) Coryneum Blight	2 2/3 - 21 1/3	7	In orchards where the disease is severe a spray should also be applied shortly after harvest.
Bloom & Growing Season: Brown Rot Blossom	1 1/3 - 4	5	Applied at popcorn and full bloom.
RESTRICTIONS Maximum single dormant season application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent) Maximum single growing season application rate is 4 pt/A (1.5 lbs metallic copper equivalent) Maximum annual application rate is 48 pt/A (18.0 lbs metallic copper equivalent)			

CHIVES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew	1 1/3	7	Begin applications when plants are established in the field. Repeat applications every 7-10 days as dictated by disease conditions.
RESTRICTIONS Maximum single application rate is 1 1/3 pints/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 7 pints/A (2.6 lbs metallic copper equivalent)			

CITRUS (Grapefruit, Kumquat, Lemon, Orange, Pummelo, Tangelo, Tangerine & Lime)			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Melanose Scab Pink Pitting	1 1/3 – 8	7	Apply as pre-bloom and post-bloom sprays.
Greasy Spot	1/3 – 4	7	May be used in concentrate sprays at equivalent rates. For aerial application use 4 pints NuCop 3L HB per 10 gallons per acre.
Brown Rot	1 1/3 – 5 1/3	7	Apply beginning in the fall and continuing as needed. Apply to skirts of trees to a height of at least 4 feet. Apply also to bare ground one foot beyond skirt. Use higher rates when conditions favor disease. NOTE: In California, in areas subject to copper injury, add 1/3 to 1 lb. of high quality lime per gallon of NuCop 3L HB.
Citrus Canker (SUPPRESSION ONLY)	4 - 8	7	Spraying flushes 7-14 days after shoots begin to grow. Young fruit may need additional application. Number and timing of applications will depend on disease pressure. Under heavy disease pressure, each flush of new growth should be sprayed.
Phytophthora	See comment	7	Mix 2/3 pint with one gallon of water and paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May before summer rains and/or in the fall before wrapping trees for freeze protection. This treatment serves as protection for up to one year, but does not cure existing infections.

CITRUS (Grapefruit, Kumquat, Lemon, Orange, Pummelo, Tangelo, Tangerine & Lime)			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
RESTRICTIONS Maximum single application rate is 8 pt/A (3.0 lbs metallic copper equivalent) Maximum annual application rate is 33 1/3 pt/A (12.5 lbs metallic copper equivalent)			

COFFEE			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Coffee Berry Disease (Collectotrichum coffeanum)	2 - 5 1/3	14	Apply after flowering and before the start of long rains and then at 14-28 day intervals until picking. Use higher rates when rainfall is heavy and disease pressure is high.
Bacterial Blight (Pseudomonas syringae)	2 - 5 1/3	14	Begin spray program before the start of long rains and then at 21 - 28 day intervals until picking. The critical time of spraying to control disease is just before, during, and after flowering(s), especially when these times coincide with wet weather. Use higher rates when rainfall is heavy and disease pressure is high.
Iron Spot (Cercospora coffeicola) & Pink Disease (Corticium salmonicolor)	2/3 - 2 2/3	14	Begin treatment at start of wet season and continue at monthly intervals for three applications.
Leaf Rust	2/3 - 5 1/3	14	Apply before the onset of rain and then at 21 day intervals while rains continue. Use higher rates when rainfall is heavy and disease pressure is high.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 33 1/3 pt/A (12.5 lbs metallic copper equivalent)			

CRANBERRY			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Fruit Rot	2 2/3 – 5 1/3	7	One or two additional applications made at 10 to 14 day intervals may be required, depending on disease pressure. Follow the advice of the local Agricultural Extension Service.
RESTRICTIONS Maximum single application rate is 5 1/3 pints/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 16 2/3 pints/A (6.25 lb metallic copper equivalent)			

CUCURBITS (CUCUMBERS, CANTALOUPEs, HONEYDEWS, MUSKMELONS, PUMPKINS, SQUASH & WATERMELONS)			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Alternaria Leaf Spot Angular Leaf Spot Anthracnose Downy Mildew Powdery Mildew Gummy Stem Blight Watermelon Bacterial Fruit Blotch (suppression)	2/3 – 2 2/3	5	Begin application when conditions are favorable for disease development. Repeat at 5-10 day intervals. Use higher rates when conditions favor disease. NOTE: Crop injury may occur from application at higher rates and shorter intervals. Discontinue use if injury occurs.
RESTRICTIONS Maximum single application rate is 2 2/3 pints/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 14 pints/A (5.25 lbs metallic copper equivalent)			

CURRANTS & GOOSEBERRY			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose Leaf Spot	3 1/3 – 6 2/3	10	Make applications, starting after harvest, before bloom and after petal fall.
RESTRICTIONS Maximum single application rate is 6 2/3 pt/A (2.5 lbs metallic copper equivalent) Maximum annual application rate is 26 2/3 pt/A (10 lbs metallic copper equivalent)			

DILL			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Phoma Leaf Spot Rhizoctonia Foliage Blight	1 1/3 – 2	7	Begin applications when plants are first established in the field and repeat at 7-10 day intervals depending upon disease severity and environmental conditions. Use higher rates when conditions favor disease.
RESTRICTIONS Maximum single application rate is 2 pt/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 10 pt/A (3.75 lbs. metallic copper equivalent)			

DOUGLAS FIR			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Rhabdocline needlecast	1 1/3 – 5 1/3	7	Begin applications at bud break and repeat at 3 - 4 week intervals. Apply in a tank mix with another registered pesticide if moderate to severe disease pressure is present.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 53 1/3 pt/A (20.0 lbs metallic copper equivalent)			

EGGPLANT			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Alternaria Blight Anthracnose Phomopsis	2/3 – 2	7	Apply before disease appears. Repeat at 7 to 10 day intervals.
RESTRICTIONS Maximum single application rate is (2 pt/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 20 2/3 (7.75 lbs metallic copper equivalent)			

ENDIVE & ESCAROLE			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew	1/3 – 2 2/3	5	Begin treatment when disease first appears and repeat every 7 - 10 days as needed to suppress disease.
RESTRICTIONS Maximum single application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent)			

FILBERTS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bacterial Blight (Post Harvest application)	5 1/3 – 16	14	In seasons of heavy rain, make another application after the leaves have dropped. Add 1 pint of a superior type oil per 100 gallons of water.
Eastern Filbert Blight	5 1/3 – 16	14	Make initial application at budswell to budbreak in enough water to obtain thorough coverage. Additional applications should be made at intervals of 10 - 14 days depending on disease severity or when conditions favor disease pressure. Add 1 pint of superior type oil per 100 gallons of water.
RESTRICTIONS Maximum single application rate is 16 pt/A (6.0 lbs metallic copper equivalent) Maximum annual application rate is 48 pt/A (24.0 lbs metallic copper equivalent)			

21/38

GINSENG			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Alternaria Leaf Stem Blight	1 - 2 2/3	7	<p>Begin tank mix applications as a tank mix with two pounds of Iprodione 50WP in 100 gallons of water per acre as soon as plants have emerged in spring.</p> <p>Applications should be repeated every seven days until plants become dormant in fall.</p> <p>Apply fungicides at least eight hours before rain, giving the fungicides time to dry on the plants. Use of a spreader-sticker is advised.</p> <p>NOTE: Alternaria Leaf and Stem Blight is most severe in humid conditions such as those found in the dense canopies of two, three, and four year old ginseng. Complete and thorough spray is required for control.</p>
RESTRICTIONS Maximum single application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 14 pt/A (5.25 lbs metallic copper equivalent)			

GRAPES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Black Rot Powdery Mildew Downy Mildew Phomopsis	2/3 - 8	3	<p>Apply at budbreak with additional applications throughout the rainy season, depending on the disease severity.</p> <p>(Attention: Slight to severe foliage injury may occur on copper-sensitive varieties such as Concord, Delaware, Niagara, and Rosette.)</p>
RESTRICTIONS Maximum single application rate is 8 pt/A (3.0 lbs metallic copper equivalent) Maximum annual application rate is 53 1/3 pt/A (20.0 lbs metallic copper equivalent)			

GUAVA			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose Red Algae	2 – 3 1/3	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
RESTRICTIONS Maximum single application rate is 3 1/3 pt/A (1.25 lbs metallic copper equivalent) Maximum annual application rate is 12 2/3 pt/A (4.75 lbs metallic copper equivalent)			

HOPS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew	2/3 – 1 1/3	10	Apply as a fungicide crown treatment (after pruning, but before training) as needed. After training, additional fungicide treatments are needed at 10 day intervals. Discontinue use 2 weeks before harvest.
RESTRICTIONS Maximum single application rate is 1 1/3 pt/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 7 pt/A (2.6 lbs metallic copper equivalent)			

KIWI			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Blossom Blight (Bud Rot) Leaf Spot (Phomopsis)	2/3 – 5 1/3	30	Make two to three applications during dormant season. Do not apply at time of or after leaf emergence.
Pseudomonas syringae Erwinia herbicola Pseudomonas fluorescens	2 – 5 1/3	30	Make applications on a monthly basis. A maximum of 3 applications may be made.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 16 pt/A (6.0 lbs metallic copper equivalent)			

LETTUCE			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew	1/3 – 2 2/3	5	Begin treatment when disease first appears and repeat every 7 - 10 days as needed to suppress disease.
RESTRICTIONS Maximum single application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent)			

LITCHI			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	2 – 3	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
RESTRICTIONS Maximum single application rate is 3 pt/A (1.1 lbs metallic copper equivalent) Maximum annual application rate is 13 pt/A (4.9 lbs metallic copper equivalent)			

LIVE OAK			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Ball Moss	See comment *	A second application may be required after 12 months	* Mix 2 – 5 1/3 pints per 100 gallons of water. Apply in spring after heavy rain, using 1.5 gallons of spray per foot of tree height. Make sure to set tufts thoroughly. (NOTE: NuCop 3L HB may be injurious to some ornamentals grown under live oaks).
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 53 1/3 pt/A (20.0 lbs metallic copper equivalent)			

MACADAMIA NUTS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	4 – 6	7	Initiate sprays at first sign of flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage
Blossom blight Raceme blight	2 – 6	7	Apply during peak raceme development and bloom period.
RESTRICTIONS Maximum single application rate is 6 pt/A (2.25 lbs metallic copper equivalent) Maximum annual application rate is 25 pt/A (9.4 lbs metallic copper equivalent)			

MAMEY SAPOTE			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose Algal Leaf Spot	4 – 5 1/3	14	Apply when conditions favor disease development. Repeat on 14-30 day schedule as disease severity and environmental conditions dictate. Use higher rates when conditions favor disease.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent)			

MANGO			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	2 2/3 – 6 2/3	30	Apply monthly after fruit set until harvest.
RESTRICTIONS Maximum single application rate is 6 2/3 pt/A (2.5 lbs metallic copper equivalent) Maximum annual application rate is 46 2/3 pt/A (17.5 lbs metallic copper equivalent)			

OLIVES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Peacock Spot Olive Knot	2 2/3 – 8 1/3	30	Apply before winter rains fall. A second application in early spring should be made if disease is severe.
RESTRICTIONS Maximum single application rate is 8 1/3 pt/A (3.1 lbs metallic copper equivalent) Maximum annual application rate is 16 2/3 pt/A (6.25 lbs metallic copper equivalent)			

ONION & GARLIC			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Purple Blotch Downy Mildew	2/3 – 2 2/3	7	Apply when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals.
RESTRICTIONS Maximum single application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 16 pt/A (6.0 lbs metallic copper equivalent)			

PAPAYA			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	1 – 7	14	Begin application before disease is expected to appear. Repeat at 14 day intervals. Use the higher rates when conditions favor disease. The addition of a suitable spreader-sticker, such as Kinetic®, may be desirable especially during periods of heavy rains.
RESTRICTIONS Maximum single application rate is 7 pt/A (2.6 lbs metallic copper equivalent) Maximum annual application rate is 56 pt/A (21 lbs metallic copper equivalent)			

PARSLEY			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bacterial Blight (Pseudomonas sp.)	2 – 2 2/3	10	Begin applications when plants are first established in the field and repeat at 10 day intervals depending upon disease severity and environmental conditions.
RESTRICTIONS Maximum single application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent) Maximum annual application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent)			

PASSION FRUIT			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracoese	4 – 6	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
RESTRICTIONS Maximum single application rate is 6 pt/A (2.25 lbs metallic copper equivalent) Maximum annual application rate is 24 pts/A (9.0 lbs metallic copper equivalent)			

PEACHES & NECTARINES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Dormant: Leaf Curl Coryneum Blight (Shot Hole)	2 2/3 – 21 1/3	7	Apply leaf fall as dormant application. Use the higher rate when rainfall is very heavy and disease pressure is high. May be used with an agricultural spray oil.

PEACHES & NECTARINES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bloom & Growing Season: Brown Rot Blossom Blight (California)	1 1/3 – 4	5	Apply as a full cover spray at pink bud. (Application at this time also affords some control of Leaf Curl and Coryneum Blight). NOTE: Do not spray later than three weeks prior to harvest. Do not use at rates above those recommended.
Bacterial Spot	5 1/3	7	Apply as a dormant spray. Apply as a post bloom spray, 2/3 pint per acre at first and second cover sprays.
RESTRICTIONS Maximum single dormant season application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent) Maximum single growing season application rate is 4 pt/A (1.5 lbs metallic copper equivalent) Maximum annual application rate is 48 pt/A (18.0 metallic copper equivalent)			

PEANUTS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Cercospora Leaf Spot	1 – 2	7	Begin spraying 35-40 days after planting or when disease symptoms appear. Use sufficient water to get adequate coverage. Continue applications at 10 to 14 day intervals. Reduce spray interval to 7 days during humid weather. Use higher rates when conditions favor disease.
RESTRICTIONS Maximum single application rate is 2 pt/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 12 2/3 pt/A (4.75 metallic copper equivalent)			

PEARS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bloom & Growing Season: Fireblight	2/3 to 1 1/3	5	Apply at 5 day intervals throughout bloom period. Excessive dosages may cause fruit russet.
Fall & Late Dormant Season: Pseudomonas blight	4 – 21 1/3	Only one dormant application allowed per season	Apply before fall rain begins. NOTE: Excessive dosages may cause fruit russet.
RESTRICTIONS Maximum single dormant season application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent) Maximum single growing season application rate is 1 1/3 pt/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 42 2/3 pt/A (16 lbs metallic copper equivalent)			

PEAS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Powdery Mildew	1 – 2	7	Begin spray treatment when disease symptoms first appear. Repeat applications at weekly intervals.
RESTRICTIONS Maximum single application rate is 2 pt/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 10 pt/A (3.75 lbs metallic copper equivalent)			

PECANS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Shuck and Kernel rot (Phytophthora cactorum) Zonate leaf spot (Cristulariella pyramidalis)	1 1/3 – 5 1/3	14	Apply in sufficient water for good coverage at 2-4 week intervals starting at kernel growth and continuing until shucks open. Use the higher rate and shorter intervals if frequent rainfall occurs.

PECANS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Mosses Algae Lichen	See Comment	Make only one application per year	Mix 4 pts per 100 gallons spray plus spreader-sticker, such as Kinetic®, on a dilute spray basis and apply in dormant season before buds swell, thoroughly wetting limbs and mosses.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent)			

PEPPERS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bacterial Spot	2/3 – 2	3	Apply, when disease threatens, in sufficient water to provide adequate coverage. Use at 7 to 14 day intervals depending on disease severity.
RESTRICTIONS Maximum single application rate is 2 pt/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 30 pt/A (11.25 lbs metallic copper equivalent)			

PISTACHIOS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Botrytis Blight Botryosphaeria Panicle Shoot Blight Septoria Leaf Blight Late Blight (Alternaria)	1 1/3 – 5 1/3	14	Make initial application at bud swell and repeat on a 14-28 day schedule. Use higher rates when conditions favor disease.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent)			

PLUMS & PRUNES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Dormant Season: Coryneum blight (Shot hole)	2 2/3 – 21 1/3	7	Apply as a dormant spray. Use the higher rate when rainfall is heavy and/or disease pressure is high.
Bloom & Growing Season: Brown rot blossom blight	1 1/3 – 4	5	Apply full cover application at pink, red or early white bud stage. Use the higher rate when disease pressure is heavy or conditions favor disease development.
RESTRICTIONS Maximum single dormant season application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent) Maximum single growing season application rate is 4 pt/A (1.5 lbs metallic copper equivalent) Maximum annual application rate is 48 pt/A (18.0 lbs metallic copper equivalent)			

POTATOES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Early & Late Blight	2/3 – 6 2/3	5	Apply at 7 to 10 day intervals beginning when plants are 6 inches high until two weeks before harvest.
RESTRICTIONS Maximum single application rate is 6 2/3 pt/A (2.5 lbs metallic copper equivalent) Maximum annual application rate is 66 2/3 (25 lbs metallic copper equivalent)			

QUINCE			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Bloom & Growing Season: Fireblight	1 1/3	5	Apply at 5 day intervals throughout bloom period. Apply in adequate water for thorough coverage.
RESTRICTIONS Maximum single dormant season application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent) Maximum single growing season application rate is 1 1/3 pt/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 42 2/3 pt/A (16 lbs metallic copper equivalent)			

RASPBERRIES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Leaf & Cane Spot	1 1/3 – 5 1/3	7	Apply delayed dormant spray after training in spring. Make fall spray application after harvest.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 26 2/3 pt/A (10.0 lbs metallic copper equivalent)			

SPINACH			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose Cercospora Leafspot Downy Mildew White Rust	2/3 – 2	7	Begin treatment when disease first appears and repeat every 7-10 days as needed to suppress disease.
RESTRICTIONS Maximum single application rate is 2 pt/A (0.75 lbs metallic copper equivalent) Maximum annual application rate is 10 pt/A (3.75 lbs metallic copper equivalent)			

STRAWBERRIES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Downy Mildew Leaf Spot Leaf Blight	1 1/3 - 4	7	Begin application when plants are established and continue on a weekly schedule throughout season. Discontinue applications if signs of phytotoxicity appear.
RESTRICTIONS Maximum single application rate is 4 pt/A (1.5 lbs metallic copper equivalent) Maximum annual application rate is 21 1/3 pt/A (8.0 lbs metallic copper equivalent)			

SUGAR APPLE (Annona)			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	2 – 4	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
RESTRICTIONS Maximum single application rate is 4 pt/A (1.5 lbs metallic copper equivalent) Maximum annual application rate is 33 2/3 pt/A (12.6 lbs metallic copper equivalent)			

SUGAR BEETS & TABLE BEETS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Cercospora Leaf Spot	2/3 – 3 1/3	10	Start spray when disease threatens and continue for 4 to 5 applications. Spray every 10 to 14 days depending on weather conditions.
RESTRICTIONS Maximum single application rate is 3 1/3 pt/A (1.25 lbs metallic copper equivalent) Maximum annual application rate is 20 pt/A (7.5 lbs metallic copper equivalent)			

SYCAMORE			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Anthracnose	2/3 – 5 1/3	7	Make two applications as a full cover spray. Use a minimum of 100 gallons water per acre. Make first application at bud crack and second application 7 to 14 days later at 10% leaf expansion.
RESTRICTIONS Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent) Maximum annual application rate is 53 1/3 pt/A (20.0 lbs metallic copper equivalent)			

TOMATOES			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Early Blight	2/3 – 1 1/3	3	When disease threatens, apply at 7 to 10 day intervals.
Bacterial Speck	2/3 – 1 1/3	3	Apply at 10-30 day intervals beginning when the disease threatens. Use more frequent applications when disease pressure is high.
Bacterial Spot Anthracnose Gray Leaf Mold Gray Leaf Spot Septoria Leaf Spot Late Blight	2/3 – 1 1/3	3	When disease threatens, apply at 7 - 10 day intervals, more frequently when disease is severe.
RESTRICTIONS Maximum single application rate is 1 1/3 pt/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 46 pt/A (17.25 lbs metallic copper equivalent)			

TURFGRASS			
DISEASE	APPLICATION RATE (pints)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Algae	6 – 8	10	May be used as a maintenance spray as needed. May be used alone or in combination with fungicides such as dithiocarbamates. Use a minimum of 100 gallons of water per acre. Phytotoxicity may depend on varietal differences. Apply the recommended rate to a small area and observe 7 - 10 days for phytotoxicity. If phytotoxicity occurs, discontinue use.
RESTRICTIONS Maximum single application rate is 8 pt/A (3.0 lbs metallic copper equivalent) Maximum annual application rate is 24 pt/A (7.0 lbs metallic copper equivalent)			

WALNUTS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Walnut Blight	2 2/3 – 8 1/3	7	Apply first spray at early pre-bloom when catkins are partially expanded. Make three additional applications during bloom and early nutlet stages at 7 to 10 day intervals. Additional applications may be necessary when frequent rainfall occurs.
RESTRICTIONS Maximum single application rate is 8 1/3 pt/A (3.1 lbs metallic copper equivalent) Maximum annual application rate is 66 2/3 pt/A (25.0 lbs metallic copper equivalent)			

WATERCRESS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Cercospora Leaf Spot	2/3 - 1 1/3	7	Begin application when plants are first established in the field, repeating at 7-14 day intervals depending on disease severity and environmental conditions. Do not exceed 4 applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.
RESTRICTIONS Maximum single application rate is 1 1/3 pt/A (0.5 lbs metallic copper equivalent) Maximum annual application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent)			

WHEAT, BARLEY & OATS			
DISEASE	APPLICATION RATE (pints/Acre)	MINIMUM DAYS RETREATMENT INTERVAL	COMMENT
Septoria Leaf Blotch Helminthosporium Spot Blotch	2/3 – 1 1/3	10	Make first application at early heading and follow with second application 10 days later.
RESTRICTIONS Maximum single application rate is 1 1/3 pt/A (0.50 lbs metallic copper equivalent) Maximum annual application rate is 2 2/3 pt/A (1.0 lbs metallic copper equivalent)			

ORNAMENTALS

Notice to User: Plant sensitivities to NuCop 3L HB have been found to be acceptable in specific genera and species listed on this label; however, phytotoxicity may occur. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for sensitivity to NuCop 3L HB. Neither the manufacturer nor seller has determined whether or not NuCop 3L HB can be safely used on ornamental or nursery plants not listed on this label. The user should determine if NuCop 3L HB can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e., bedding plants, foliage, etc., and observe for 7-10 days for symptoms of phytotoxicity prior to commercial use.

Use NuCop 3L HB on container, bench, or bed-grown ornamentals in greenhouses or outdoor nurseries, for professional use on ornamentals grown for indoor and outdoor landscaping, and for control of bacterial and fungal diseases of foliage, flowers and stems.

Apply as a thorough coverage spray using 1 1/3 pints per 100 gallons of water. Begin application at first sign of disease and repeat at 7 to 14 day intervals as needed; use shorter interval during periods of frequent rains or when severe disease conditions persist.

NuCop 3L HB may be used as a maintenance spray alone or in combination with other fungicides such as the dithiocarbamates.

Restrictions:

Maximum single application rate is 5 1/3 pt/A (2.0 lbs metallic copper equivalent)
Maximum annual application rate is 53 1/3 pt/A (20 lbs metallic copper equivalent)

ORNAMENTAL/DISEASES:

- Althea (Rose of Sharon)/Bacterial Leaf Spot
- Aralia/Xanthomonas & Cercospora Leaf Spots, Alternaria
- Arborvitae/Alternaria Twig Blight, Cercospora Leaf Blight
- Azalea*/Cercospora Leaf Spot, Botrytis Blight, Phytophthora Dieback & Powdery Mildew
- Begonia/Xanthomonas Leaf Spot, Anthracnose
- Bougainvillea/Anthracnose, Bacterial Leaf Spot
- Bulbs (Easter Lily**, Tulip, Gladiolus)/Anthracnose, Botrytis Blight
- Camellia/Anthracnose, Bacterial Leaf Spot
- Camphor Tree/Pseudomonas Leaf Spot
- Canna/Pseudomonas Leaf Spot
- Carnation*/Alternaria Blight, Pseudomonas Leaf Spot, & Botrytis Blight
- Chinese Tallow Tree/Bacterial Leaf Spot (Xanthomonas sp., Pseudomonas sp.)
- Chrysanthemum*/Septoria Leaf Spot & Botrytis Blight
- Cotoneaster/Botrytis Blight
- Dahlia/Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
- Date Palm/Pestalotia Leaf Spot
- Dianthus/Bacterial Spot, Bacterial Soft Rot
- Dogwood/Anthracnose
- Dusty Miller/Bacterial Leaf Spot (Pseudomonas cichorii)
- Echinacea/Bacterial Leaf Spot (Pseudomonas cichorii)
- Elm "Drake"/Xanthomonas Leaf Spot
- Euonymus/Botrytis Blight & Anthracnose
- European Fan Palm/Pestalotia Leaf Spot
- Gardenia/Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot
- Geranium/Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
- Gladiolus/Alternaria Leaf Spot, Botrytis Gray Mold, Bacterial Leaf Blight

Golden Rain Tree/Bacterial Leaf Spot
 Hibiscus/Bacterial Leaf Spot
 Holly Fern/Pseudomonas Leaf Spot
 Impatiens/Bacterial Leaf Spot
 India hawthorn (greenhouse)/Entomosporium Leaf Spot
 Ivy*/Xanthomonas Leaf Spot
 Ixora/Xanthomonas Leaf Spot
 Juniper (Eastern Red Cedar)/Anthracnose
 Lantana/Bacterial Leaf Spot
 Lilac/Cercospora Leaf Spot
 Loblolly Bay/Anthracnose
 Loquat/Entomosporium maculata, Colletotrichum sp.
 Magnolia (Southern)/Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot
 Mandevillas/Anthracnose
 Marigold/Alternaria Leaf Spot, Botrytis Leaf and Flower Rot,
 Cercospora Leaf Spot
 Mulberry, Weeping/Bacterial Leaf Spot
 Oak, Laurel/Algal Leaf Spot (Cephaleuros virescens)
 Oleander/Bacterial Leaf Spot, Fungal Leaf Spot
 Pachysandra/Volutella Leaf Blight
 Pansy/Downy Mildew
 Pear (Flowering)/Fireblight, Leaf Spot
 Pentas (Egyptian Star)/Bacterial Leaf Spot (Xanthomonas sp.)
 Peony/Botrytis Blight
 Periwinkle/Phomopsis Stem Blight
 Philodendron/Bacterial Leaf Spot
 Phlox/Alternaria Leaf Spot
 Photinia (Red Tip)/Anthracnose, Entomosporium Leaf Spot
 Pistachio/Anthracnose
 Plantain Lily/Bacterial Leaf Spot
 Powder Puff Plant/Bacterial Leaf Spot
 Pyracantha/Fireblight & Scab
 Queen Palm/Exosporium Leaf Spot, Phytophthora Bud Rot
 Rhododendron/Alternaria Flower Spot
 Rose*/Powdery Mildew, Black Spot
 Verbena/Xanthomonas Leaf Spot
 Viburnum/Anthracnose
 Washingtonia Palm/Pestalotia Leaf Spot
 Weeping Willow/Anthracnose
 Yucca (Adams Needle)/Cercospora & Septoria Leaf Spot

*Discoloration of foliage and/or blooms have been noted on some varieties. To prevent residues on commercial plants, do not spray just before selling season.

**For Easter Lily, use 2 - 6 2/3 pints per 100 gallons.

Restrictions (Easter Lily):

Maximum single application rate is 6 2/3 pt/A (2.5 lbs metallic copper equivalent)

Maximum annual application rate is 200 pt/A (75 lbs metallic copper equivalent)

Kinetic® is a registered trademark of Helena Chemical Company

37
/38

CONDITIONS OF SALE

LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale – Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Albaugh, Inc. (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Albaugh's election, one of the following:

1. Refund of the purchase price paid by the buyer or user for product bought, or
2. Replacement of the product used.

To the extent allowed by law, the Company shall not be liable and any claims against the Company are waived for special, indirect, incidental, or consequential damages or expenses of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the forgoing conditions of sale and limitation of warranty, liability and remedies.

Kinetic is a registered trademark of Helena Chemical Company.

CORPORATE OFFICE

1525 NE 36th Street
 Ankeny, IA 50021
 515.964.9444 - Office
 800.247.8013 - Toll Free
 515.964.7813 - Facsimile

ALBAUGH, INC.

Valdosta Office 38/38
 P.O. Box 2127
 304 Janet Street, Suite H
 Valdosta, GA 31604
 229.244.3288 - Office
 229.244.5841 - Facsimile

FED-X

November 28, 2007

Document Processing Disk (NOTIF)
 Office of pesticide Programs (7504P)
 U.S. Environmental Protection Agency
 Room S-4900, One Potomac Yard
 2777 South Crystal Drive
 Arlington, VA 22202-4501

RE: Label notification for "Pesticide Management and Disposal; Standards for Pesticide Containers and Containment"

Dear Sirs,


The enclosed submissions are draft labels submitted in response to Pesticide Regulation Notice 2007-4 for the following Albaugh registrations:

PRODUCT	EPA REG. NO.
Tebuconazole 28% FL ST	42750-130
NUCOP 3L HB	42750-167
Weed Hoe 108	42750-29
Weed Hoe 120	42750-28
Picloram K-Salt Rangeland	42750-79
Picloram K-Salt IVM	42750-81

Changes are noted in ~~strikeout~~ for deleted text and underline/bold for added text.

Please call if you have any questions.

Regards,



Morris Gaskins
 Registrations Manager
 Albaugh, Inc.

Agri Star[®]
 By Albaugh, Inc.

PREMIER SUPPLIER OF OFF-PATENT CROP PROTECTION PRODUCTS

www.albaughinc.com