



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 5, 2017

Carolyn Miter
Registrations Specialist
Albaugh, LLC
P.O. Box 2127
Valdosta, GA 31604

Subject: Notification per PRN 98-10 – updating logo per NY request
Product Name: Myclobutanil 2% Homeowner
EPA Registration Number: 42750-196
Application Date: August 28, 2017
Decision Number: 532807

Dear Ms. Miter:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact me at 7033055410 or via email at johnson.hope@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be "H. Johnson", written over a circular scribble.

Hope Johnson, Product Manager 21
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

09/05/2017

DITOR'S NOTE: Update precautionary graphics on page 3 for New York State

MYCLOBUTANIL 2% HOMEOWNER
Multi-Purpose Systemic Fungicide

[Do-it-yourself disease control • Prevents and controls all major lawn and ornamental diseases • Controls brown patch, crown rot, powdery mildew, leaf spot, rust, Summer patch and more • Systemic action • Systemic action - Can't be washed off by rain once dried • Rainproof –Can't be washed off [away] by rain once dried • Protects for up to 2 weeks • Treats [3,000] [6,000] [12,000] sq ft • Just connect to your [garden] hose and spray • Easy-to-use spray ready nozzle • No spreading or complicated mixing • Water-based • Easy-View [Clear-View] measuring window • Guaranteed Results or your money back J*

[MYCLOBUTANIL 2% HOMEOWNER prevents and controls all major diseases on lawns and ornamentals. Because it works systemically, it can't be washed off by rain or sprinklers once dried and it works for weeks. Use this product on your lawn to keep it thriving.]

[Prevents and controls... Brown patch Dollar Spot, Powdery Mildew, Rust, Blight, Leaf Spot... and more!]

*[One 32 oz bottle of this formula will treat 3,000 sq ft]
[One 64 oz bottle of this formula will treat 6,000 sq ft]
[One gallon bottle of this formula will treat 12,000 sq ft]*

[CONTROLS] [DISEASES CONTROLLED]

[Use MYCLOBUTANIL 2% HOMEOWNER as directed to treat the following diseases in your lawn and ornamentals:]

MYCLOBUTANIL 2% HOMEOWNER prevents and *controls* anthracnose, *brown patch*, copper spot, dollar spot, leaf spot (including septoria, zonate), melting out, crown rot, leaf smuts, powdery mildew, rust, red thread, Spring dead spot, necrotic ring spot and Summer patch, roses, lawns, flowers, ornamental shrubs and trees. Protects fruits, nuts and vegetables.

Active Ingredient:	
Myclobutanil	2.0%
Other Ingredients:	98.0%
Total:	100.0%

Read all instructions inside before using.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back [booklet] for additional precautionary statements

EPA Reg. No. 42750-196

EPA Est. No. xxxxx-xx-xxx


Net Contents: [1 quart] [2 quarts] [1 gallon]

MANUFACTURED FOR:
ALBAUGH, LLC
1525 NE 36th STREET
ANKENY, IA 50021

FIRST AID	
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a Poison Control Center or doctor for treatment advice.
IF SWALLOWED:	Call a Poison Control Center or doctor immediately for treatment advice. Have person sip glass of water if able to swallow. Do not induce vomiting unless told to by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling the Poison Control Center or doctor, or going for treatment. For transportation or medical emergency information involving this product, please call 1-800-424-9300.	

[Optional Prop 65 Statements for California]

 **WARNING:** Reproductive Harm – www.P65Warnings.ca.gov]

 **WARNING:** This product can expose you to chemicals including myclobutanil which is known to the State of California to cause birth defects or other reproductive harm. For more information go to www.P65Warnings.ca.gov]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

USER SAFETY RECOMMENDATIONS

1. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
2. User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry area away from heat and open flame.

PESTICIDE DISPOSAL: If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

NOTE: This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous

area and recheck in a few hours. Do not use if any staining is observed.

[OPTIONAL PRODUCT FACT BOX]

PRODUCT FACTS

What it Does: Prevents and controls all Major Plant Diseases including Anthracnose, Brown patch, Copper Spot, Dollar Spot, Leaf Spot (including Septoria, Zonate), Melting Out, Crown Rot, Leaf Smuts, Powdery Mildew, Rust, Red Thread, Spring Dead Spot, Necrotic Ring Spot and Summer Patch

Where to Use: Outdoors on roses, flowers, ornamentals, shrubs, trees, lawns, fruit and nut trees, berries and vegetables.

When To Use: Start Treating When Grass Begins to Turn Green in Spring or When Signs of Diseases are First Noticed

Questions & Comments (phone icon) (computer icon)

Call 1-800-247-8013 or email Albaugh, LLC

STOP. READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For home use only.

Shake well before using.

Keep pets and people other than applicator off treated areas until spray has dried. *(man/stop sign/walking dog icon)*



WHEN TO USE

This product is highly effective when it is used to prevent diseases or applied at the first sign of disease. Start treating when grass begins to turn green in the spring or when signs of diseases are first noticed. For best disease control, treat every 2 weeks.

RESTRICTIONS

- Do not apply more than 13 treatments per year. In Nassau and Suffolk Counties in New York State, do not apply more than 3 times per given area per year.

- Do not treat lawns or plants during periods of stress, i.e. heat, drought, flooding, insects, excessive fertilizer, herbicides or pesticide use, etc.
- Do not apply when temperatures are above 85°F as injury to lawn can occur.
- Only use hose end sprayers for broadcast lawn application. For application to individual plants and lawn spot treatment only use conventional pump up sprayers at labeled rates.
- Apply to the lawn after mowing, or allow the area sprayed to dry completely prior to mowing.
- If it rains before product has dried, reapply.
- If disease is already present, this product will not eliminate it.

If you need more information, contact your local County Extension Service.

[TO PROTECT] LAWNS

MYCLOBUTANIL 2% HOMEOWNER prevents and cures Anthracnose, Brown patch, Copper Spot, Dollar Spot, Leaf Spot (including Septoria, Zonate), Melting Out, Crown Rot, Leaf Smuts, Powdery Mildew, Rust, Red Thread, Spring Dead Spot, Necrotic Ring Spot and Summer Patch on home lawns.

Use 2 2/3 fl oz/half-gallon of water to treat 250 sq. ft. Start treating when grass begins to turn green in the spring or when signs of diseases are first noticed. For best disease control treat every 2 weeks

HOW TO USE

Determine the size of the area (lawn) to be treated. To calculate square feet, multiply length by width.

Start at the farthest point to be treated and work backward so you don't walk over treated area. Walk at a steady pace and spray using a side-to-side [arc] motion, slightly overlapping the treated area.

The entire [32 oz] bottle will treat 3,000 sq ft of lawn. Refer to [sight gauge] [Easy-View] [Clear-View] measuring window to guide coverage. When bottle is half empty, you should have covered half the area.

ENTIRE LAWN

(A) - HOSE END SPRAYER INSTRUCTIONS [Dial Button]

1. *Shake container well before using.*
2. *Connect a garden hose to the Ready Spray nozzle. Make sure the dial on the nozzle is in the "OFF" position with the [green] safety tab in the valve notch.*
3. *Turn on water at faucet. Extend hose to the farthest area to be treated and work back toward the faucet so you don't come in contact with the treated area.*
4. *To BEGIN spraying, point nozzle toward lawn or plants to be treated and 1.) bend the [green] safety tab (located at right of [yellow] dial) back and hold, 2.) using your other hand, QUICKLY turn the dial clockwise until it stops and [green] tab engages into notch. Water will automatically mix with the product.*
5. *Spray lawn or plants thoroughly. Walk at a steady pace while spraying using an even sweeping motion, slightly overlapping treated areas. [One quart] [two quarts] [one gallon] will treat [3,000] [6,000] [12,000] sq ft. Calculate square feet by multiplying length by width. Refer to sight gauge to guide coverage (example: when one quart bottle is half empty, you should have covered 1,500 square feet).*

6. To STOP spraying, QUICKLY turn the dial in the opposite direction of "ON" (counter clockwise) until it stops and the safety tab engages into the notch. Turn water off at faucet. To relieve pressure before removing nozzle from hose, bend the safety tab back and turn dial "ON" until water stops spraying.
7. To STORE unused product, make sure the dial is in the "OFF" position with the green safety tab in the valve notch. Place in a cool area away from heat, sunlight or open flame.

[OPTIONALINSTRUCTIONS]

(B) - HOSE END SPRAYER INSTRUCTIONS

1. Attach to garden hose.
2. Turn on the water faucet.
3. Activate the sprayer by pushing the vertical tab towards the horizontal tab, which allows for resealing of partially used bottles when in the "OFF" position. The product channel is not opened until the control knob passes the vertical tab and is placed in the "ON" position.
4. When spraying is complete, or if you stop spraying at any time, move the horizontal tab back into the "OFF" position.
5. Rinse any residual product from the sprayer by passing through the "WATER" position on the sprayer.
6. Turn off the water at the faucet.
7. Remove the container from the garden hose, then rinse thoroughly and follow the directions in the Storage and Disposal section for storing or disposing of the container.

Rainproof— rain or watering 2 hours after application will not wash away effectiveness. [If rainfall occurs within 2 hours of application, the product should be reapplied.]

[OPTIONALINSTRUCTIONS]

(C) - HOSE END SPRAYER INSTRUCTIONS

READY-TO-SPRAY DIRECTIONS FOR USE OUTDOORS ON HOME LAWNS

HOW TO USE THE READY-TO-SPRAY SYSTEM

Connect

1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.

Spray

1. To begin spraying, point nozzle in the direction you want to spray.
2. [Bend small plastic tab back and] Turn knob [clockwise] to ON position.
3. Spray evenly to the area to be treated. Refer to the [sight gauge] clear view strip (graduated scale) on the side of the container to determine the amount of product sprayed.

Finish

1. To stop spraying, [turn knob counter clockwise] to OFF position.
2. Turn off water.
3. Relieve water pressure by [bending plastic tab back and] turning knob to ON position until water slows to a drip. Then turn knob back to OFF position.
4. Disconnect sprayer from hose.

[TO PROTECT] ROSES, FLOWERS, ORNAMENTAL SHRUBS & TREES

MYCLOBUTANIL 2% HOMEOWNER prevents and controls Black spot, Blight (blossom, cercospora leaf, ovulinia petal), Leaf spot (including cercospora), Powdery mildew, Rust (including needle, fusiform), Scab (including fusicladium) and other diseases on a wide variety of roses, flowers, trees and ornamental shrubs.

See table for a complete list of plants.

HOW TO TREAT

Use 3/4 fl. oz./gallon of water. Apply to the point when spray just begins to runoff the leaves. Apply once every 2 weeks throughout the season or until conditions favoring disease development (hot, moist weather) are no longer present. Allow spray to dry before watering.

USE TIPS

1. Treat roses on a 7 to 10 day schedule. If black spot is not a problem, a 10 to 14 day schedule is recommended.
2. When treating azaleas, mountain laurel or rhododendrons, begin spraying when flowers start to show color.
3. When treating irises, increase use rate to 1 1/2 fl. oz./gallon of water.
4. For Douglas fir, loblolly pine and slash pine use 2 1/4 fl. oz./gallon of water. Begin treating in early spring and continue treating every 2 to 3 weeks until threat of infection has passed.

PLANT (Number following plant refers to the specific Use Tip above.)	DISEASES
Rose(1)	Black spot, Powdery mildew, Rust
Azalea (2), Mountain laurel (2), Rhododendron (2)	Petal blight (ovulinia spp.) Powdery mildew
Abelia, Acalypha (Copper-leaf), Catalpa, Flowering cherry,	Cercospora leaf spot,
Hackberry, Honeysuckle, Privet, Trumpet creeper. Zinnia	Powdery mildew
Ageratum, Anemone, Angelica, Ash, Aster, Birch, China aster, Chokeberry, Chrysanthemum, Columbine, Cornflower, Dianthus, Flowering quince, Flowering maple (Abutilon), Four o'clock, Fuchsia, Iris (3), Marigold, Moonflower, Ninebark, Russian olive. Smoke-tree (Gothus)	Rust
Achillea (Yarrow), Alder, Amelanchier (Juneberry, Shadbush), Barberry, Carnation, Delphinium, Gaillardia, Gardenia, Seranium, Hollyhock, Mock-orange, Pansy, Petunia, Photinia, Salvia, Snap-dragon, Viburnum	Powdery mildew, Rust
African Violet, Begonia, Bittersweet, Buckeye, California poppy, Chestnut (Horse), Cosmos, Crepe myrtle, Dahlia, Elm, Euonymus, Gourd (Ornamental), Grape leaf ivy, Lilac, Locust, Maple, Oak, Polnsettia, Spirea, Sycamore, Walnut, Willow	Powdery mildew
Flowering almond	Blossom blight (Monilinia spp.)
Amorpha (False indigo), Bellflower, Buttonbush, Fleabane, Phlox, Sunflower	Cercospora leaf spot, Powdery

	mildew, Rust
Calendula, Hydrangea, Leucothoe	Cercospora leaf spot
Flowering crabapple, Flowering pear, Hawthorn	Powdery mildew, Rust, Scab
Dogwood	Anthracnose, Powdery mildew
Douglas fir (4)	Needle rust
Loblolly pine, Siasirpine (4)	Fusiform rust
Pyracantha	Fusicladium scab

[TO PROTECT] FRUITS AND NUTS

ALMOND

(blossom blight, rust, and shothole control; anthracnose suppression)

Mix 1/3 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. For anthracnose, blossom blight, and shothole, apply at pink bud stage (when about 5% of the buds have bloomed). If conditions favoring disease development (hot, moist weather) exist, reapply at full bloom and when the petals fall. For rust, apply MYCLOBUTANIL 2% HOMEOWNER about 8 weeks after flowering if conditions favoring disease development exist. Reapply every 21 days so long as disease symptoms persist.

- Do not apply within 90 days of harvest.
- Do not apply more than 3 times per season.

APPLE and MAYHAW

(powdery mildew, rusts and scab control)

Mix 1/2 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. Reapply every 7 to 10 days so long as disease symptoms persist. For powdery mildew and rusts, apply at pink bud stage when the buds are opening and the pink petals are visible. For scab, apply MYCLOBUTANIL 2% HOMEOWNER when the first green leaf tips are visible.

- Do not apply within 14 days of harvest.
- Do not apply more than 10 times per season.

STONE FRUITS

For all stone fruits, MYCLOBUTANIL 2% HOMEOWNER may be applied up to the day of harvest.

APRICOT AND NECTARINE

(brown rot blossom blight, brown rot, powdery mildew, and shothole control)

Mix 1/3 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. Apply at early red bud stage (the tips of the petals are visible and the flowers are still closed) before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist.

- Do not apply more than 7 times per season.

CHERRY

(brown rot blossom blight, brown rot, leaf spot, and powdery mildew control)

Mix 1/3 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. Apply when single flower buds are visible (but still closed), on short stalks and before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist. For leaf spot, make an additional application of MYCLOBUTANIL 2% HOMEOWNER after harvest.

- Do not apply more than 7 times per season.

PEACH

(brown rot; blossom blight, brown rot, powdery mildew, and rust control)

Mix 1/3 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. For brown rot blossom, brown rot, and powdery mildew, apply at early pink bud stage (the tips of the petals are visible and the flowers are still closed) before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist. For rust, apply MYCLOBUTANIL 2% HOMEOWNER about 8 weeks after flowering if conditions favoring disease development (hot, moist weather) exist. Reapply every 21 days so long as disease symptoms persist.

- Do not apply more than 7 times per season.

PLUM AND PRUNE

(brown rot blossom blight, brown rot, and rust control)

Mix 1/3 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the tree to point of runoff. For brown rot blossom blight and brown rot, apply when the first green leaf tips are visible and before infection occurs. Reapply every 7 to 14 days so long as disease symptoms persist. For rust, apply MYCLOBUTANIL 2% HOMEOWNER about 8 weeks after flowering if conditions favoring disease development (hot, moist weather) exist. Reapply every 21 days so long as disease symptoms persist.

- Do not apply more than 7 times per season.

STRAWBERRY

(leaf blight, leaf spot, and powdery mildew)

Mix 1 fl oz of MYCLOBUTANIL 2% HOMEOWNER E with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. MYCLOBUTANIL 2% HOMEOWNER may be applied up to the day of harvest. Apply when the disease first appears or when conditions favoring disease development (hot, moist weather) exist. Reapply every 14 to 21 days so long as disease symptoms persist.

- Do not apply more than 6 times per season.
- Do not plant a new crop in the same place as the crop that was treated within 30 days of the last application.

[TO PROTECT] BERRIES

For all berries, MYCLOBUTANIL 2% HOMEOWNER may be applied up to the day of harvest.

BLACKBERRY*, RASPBERRY*, LOGANBERRY

(cane and leaf rust, leaf spot, orange rust, powdery mildew, and yellow rust control)

Mix 1/2 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply at budbreak and reapply every 10 to 14 days so long as disease symptoms persist.

- Do not apply more than 4 times per season.

CURRANT

(powdery mildew control)

Mix 1 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply before bloom, through bloom and up to 2 weeks post-bloom.

- Do not apply more than 8 times per season.

GOOSEBERRY

(anthracnose and powdery mildew control)

Mix 1 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. For anthracnose, apply when the first leaf has completely unfolded. If conditions favoring disease development (hot, moist weather) exist, reapply every 10 to 14 days. For powdery mildew, apply MYCLOBUTANIL 2% HOMEOWNER before bloom, through bloom, and up to 2 weeks post-bloom.

- Do not apply more than 8 times per season.

*Use on the following members of the caneberry group including varieties and/or hybrids:

Bingleberry, black satin berry, boysenberry, cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, Darrowberry, dewberry, Oregon evergreen berry, phenomenal berry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, Himalayaberry, Dirkson Thornless, hullberry, Lavacaberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry

GRAPES

(anthracnose, black rot and powdery mildew control)

Mix 1 fl oz of MYCLOBUTANIL 2% HOMEOWNER with one gallon of water and uniformly apply to all parts of the plant to point of runoff.

For anthracnose and black rot, apply when new shoots are 1 to 3 inches in length. Reapply every 14 days so long as disease symptoms persist.

For powdery mildew, apply MYCLOBUTANIL 2% HOMEOWNER when shoots are 12 to 18 inches in length (before bloom). Reapply every 21 days so long as disease symptoms persist.

- Do not apply within 14 days of harvest.
- Do not apply more than 5 times per season.

PEPPERMINT AND SPEARMINT
(powdery mildew and rust control)

Mix 1 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply in the early spring when the plants have broken dormancy. Reapply every 14 to 21 days so long as disease symptoms persist.

- Do not apply within 30 days of harvest.
- Do not apply more than 3 times per season.

VEGETABLES

ASPARAGUS (rust control)

Mix 1 fl oz of MYCLOBUTANIL 2% HOMEOWNER with one gallon of water and uniformly apply to all parts of the plant to point of runoff. Do not apply to harvestable spears. In all locations, apply to the developing ferns after harvest. Reapply every 14 days so long as disease symptoms persist.

In all states except California:

- MYCLOBUTANIL 2% HOMEOWNER may be applied within 180 days of harvest.
- Do not apply more than one time per growing season.

In California only:

- MYCLOBUTANIL 2% HOMEOWNER may be applied within 30 days of harvest.
- Do not apply more than four times per year.

CUCURBITS (powdery mildew control)

Includes use on the following members of the cucurbits group:

acorn squash	Chinese waxgourd	honeydew melon	scallop squash
balsam apple	citron melon	hubbard squash	snake melon
balsam pear	crenshaw melon	mango melon	spaghetti squash
bitter melon	crookneck squash	muskmelon	straightneck squash
butternut squash	cucumber	ornamental gourd	summer squash
calabaza	edible gourd	Persian melon	vegetable marrow
cantaloupe	gherkin	pineapple melon	watermelon
casaba	golden pershaw melon	pumpkin	winter squash
Chinese cucumber	honey balls	Santa Claus melon	zucchini

Mix 1 fl oz of MYCLOBUTANIL 2% HOMEOWNER with one gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply MYCLOBUTANIL 2% HOMEOWNER when the disease first appears. Reapply every 7 to 10 days so long as disease symptoms persist.

- MYCLOBUTANIL 2% HOMEOWNER may be applied up to and including the day of harvest.
- Do not apply more than five times per season.
- Do not plant a new crop in the same place as the crop that was treated within thirty days of the last application.

SNAP BEAN

(pod tip rot and rust control)

Mix 1 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. For rust, apply when the disease first appears. For pod tip rot, apply when pods begin to develop. For both diseases, if conditions favoring disease development (hot, moist weather) exist, reapply every 7 to 10 days.

- MYCLOBUTANIL 2% HOMEOWNER may be applied up to and including the day of harvest.
- Do not apply more than 4 times per season.
- Do not plant a new crop in the same place as the crop that was treated within 30 days of the last application.

TOMATO

(powdery mildew control)

Mix 3/4 fl oz of MYCLOBUTANIL 2% HOMEOWNER with 1 gallon of water and uniformly apply to all parts of the plant to point of runoff. Apply when the disease first appears or if conditions favoring disease development (hot, moist weather) exist. Reapply every 21 days so long as disease symptoms persist

- MYCLOBUTANIL 2% HOMEOWNER may be applied up to and including the day of harvest.
- Do not apply more than 4 times per season.
- Do not plant a new crop in the same place as the crop that was treated within 30 days of the last application.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.