

42750-198

08/22/2008

1/9



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (H7505C)
 401 "M" St., S.W.
 Washington, D.C. 20460

EPA Reg. Number: 42750-198
 Date of Issuance: AUG 22 2008
 Term of Issuance: Unconditional

Name of Pesticide Product:
 Triclopyr 61% BEE Turf

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Mr. Morris Gaskins
 Albaugh Inc.
 121 NE 18th St.
 Ankeny, IA 50021

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Change the registration number to "42750-198".
3. Submit final labeling for this product **within 30 days** of the date of this letter.

Approving Official

Date

Tobi Colvin-Snyder for Jim Tompkins

AUG 22 2008

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

If you have any questions about this letter, please contact Tobi Colvin-Snyder at 703-305-7801.

Tobi Colvin-Snyder for

Jim Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7505C)

3/9

TRICLOPYR 61% BEE TURF

For the control of annual and perennial broadleaf weeds and kikuyugrass in ornamental turf.

ACTIVE INGREDIENT:	
Triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, butoxyethyl ester	61.6%
INERT INGREDIENTS:	<u>38.4%</u>
TOTAL:	100.0%

Contains petroleum distillates
Acid equivalent: triclopyr - 44.3% - 4 lb/gal

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment.

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. Call poison control or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300.


Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

ERA Reg. No. 42750-ROI

EPA Est. No. Xxxxxx-xx-xxx

Net Contents:

Manufactured By:
ALBAUGH, INC.
ANKENY, IA 50021



4/9

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing mists or vapors. Avoid contamination of food.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selections chart.

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton
3. Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying,

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

1. Coveralls
2. Chemical-resistant gloves such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: For applications to turf other than sod farms or turf grass grown for seed, do not allow entry into treated areas until sprays have dried unless coveralls, chemical resistant gloves, and shoes plus socks are worn.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store above 28°F or agitate before use.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures.

CONTAINER DISPOSAL:

Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable <5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable >5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container (250 gallon & bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

General: Consult federal, state, or local disposal authorities for approved alternative procedures.

GENERAL INFORMATION

TRICLOPYR 61% BEE TURF herbicide is recommended for the control of actively growing annual and perennial broadleaf weeds and kikuyugrass in perennial bluegrass, perennial ryegrass, or tall fescue ornamental turf including these turf grasses in sod farms and golf courses.

GENERAL USE PRECAUTIONS
(Read Carefully Before Using)

In Arizona: The state of Arizona has not approved TRICLOPYR 61% BEE TURF for use on sod farms.

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

TRICLOPYR 61% BEE TURF may injure certain turf grass species. Do not apply to bahiagrass, bentgrass, bermudagrass, centipedegrass, St. Augustine grass, or zoysiagrass, unless turf injury can be tolerated.

Do not apply TRICLOPYR 61% BEE TURF directly to, or otherwise permit it to come in contact with or permit spray mists containing TRICLOPYR 61% BEE TURF to drift onto, cotton, grapes, tobacco, vegetable crops, flowers, fruit or orchard trees, shrubs, or other desirable broadleaf plants.

Do not apply more than 8 lbs/ae/A/year of triclopyr.

Do not apply by aerial application.

Do not allow grazing to any area treated with TRICLOPYR 61% BEE TURF.

Do not apply TRICLOPYR 61% BEE TURF to exposed roots of shallow rooted trees and shrubs.

Do not apply to golf course greens.

For spot treatments, do not apply more than 2 quarts of TRICLOPYR 61% BEE TURF per acre in a single application.

AVOID INJURIOUS SPRAY DRIFT: Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Spray drift can be reduced by spraying when wind velocity is low, keeping the spray boom as low as possible, using nozzles that deliver larger spray droplets, and by using higher volume sprays and lower pressures.

Do not apply to ditches used to transport irrigation water.

Do not apply where runoff or irrigation water may flow onto susceptible crops.

CHEMIGATION: Do not apply this product through any type of irrigation system.

TRICLOPYR 61% BEE TURF is formulated as a low volatile ester. However, the combination of spray contact with impervious surfaces, such as roads and rocks, and increasing ambient temperatures, may result in an increase in the volatility potential for this herbicide, increasing the risk for off-target injury to sensitive crops such as grapes and tomatoes.

Broadleaf Weeds Controlled By TRICLOPYR 61% BEE TURF Include:

black medic	lespedeza
bull thistle	matchweed
burdock	mustard
Canada thistle	oxalis
chicory	plantain
clover	ragweed
creeping beggarweed	smartweed
curly dock	sweet clover
dandelion	vetch
field bindweed	wild carrot
Goldenrod	(Queen Annes lace)
ground ivy	wild lettuce
kikuyugrass +	wild violet
lambquarters	yarrow

+See control of kikuyugrass under "Application Directions".

MIXING INSTRUCTIONS

When **TRICLOPYR 61% BEE TURF** is mixed with water it forms an emulsion (not a solution) and separation may occur unless the spray mixture is agitated continuously.

Add about one-half the required amount of clean water to the spray tank. Start agitation and add the specified amount of **TRICLOPYR 61% BEE TURF**. Provide moderate agitation while completing the addition of water and during application.

APPLICATION DIRECTIONS

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled newly seeded turf should be mowed 2 or 3 times before being treated. When making applications to mature plants, hard to control species, or during drought conditions, use higher rates. Application under drought conditions may provide less than desirable results. Use low

pressure sprays to minimize spray drift. Do not water for 24 hours after application.

Reseeding Precaution: Do not reseed for 3 weeks after application. (This precaution does not apply when bermudagrass turf is overseeded with perennial ryegrass at a minimum reseeding rate of 400 lb per acre.)

Broadcast Treatment of Ornamental Turf

Apply 1/2 to 1 quart per acre of TRICLOPYR 61% BEE TURF in enough water to provide uniform coverage of the target area to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue.

Do not use on other turfgrass species (see "General Use Precautions") unless injury can be tolerated. To minimize turf injury, do not treat if turf is under heat or drought stress and make repeat applications at least 4 weeks apart.

Tank Mixing: To improve the spectrum of activity, TRICLOPYR 61% BEE TURF may be tank mixed at a rate of 1/2 to 1 pint per acre with specified rates of low volatile amine or ester formulations of 2,4-D, MCPP, or other labeled postemergence broadleaf herbicides provided that the tank mix product label do not prohibit such mixing. All tank mix product labels use directions, precautions, and limitations must be followed.

Spot Treatment of Ornamental Turf

Mix 3/8 to 3/4 ounces of TRICLOPYR 61% BEE TURF per 1000 square feet in enough water to provide uniform coverage of the target area and apply at any time broadleaf weeds are susceptible. Note:

Do not apply more than 2 quarts per acre or 1.5 ounces per 1000 square feet of TRICLOPYR 61% BEE TURF in a single application.

Control of Kikuyu grass

Apply TRICLOPYR 61% BEE TURF at a rate of 1/2 to 1 quart per acre. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

Suppression of Bermudagrass

Apply TRICLOPYR 61% BEE TURF at the rate of 1 quart per acre. Three to four additional applications at 4 week intervals will be required to give adequate suppression of bermudagrass and allow fescue or other desired turfgrass species to dominate.

To improve suppression and control of bermudagrass, 1 quart per acre of TRICLOPYR 61% BEE TURF may be tank mixed with a post emergence grass herbicide registered for this use pattern.

Three to four additional applications of this tank mix at 4 week intervals should be made to achieve control. Reseeding following application will accelerate the transition to cool season turf (see Reseeding Precaution above).

WARRANTY DISCLAIMER

Albaugh, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Albaugh, Inc. MAKES NO OTHER EXPRESS OR IMPLIED

WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Albaugh, Inc. or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent consistent with applicable law, The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Albaugh, Inc.' election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used

To the extent consistent with applicable law, Albaugh, Inc. shall not be liable for losses or damages resulting from handling or use of this product unless Albaugh, Inc. is promptly notified of such loss or damage in writing. To the extent consistent with applicable law, In no case shall Albaugh, Inc. be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Albaugh, Inc. or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.